

# APPENDIX A

## Emissions Inventory

## Appendix A Emissions Inventory

The 2012, 2018, 2021, and 2024 emission inventories are presented in various formats and details in this appendix.

### Appendix A-1 Estimated Forecast Summary by EIC

The Appendix A-1 (available separately in electronic file format) contains the estimated VOC and NO<sub>x</sub> forecast summaries by EIC emission categories for the Sacramento Federal Nonattainment Area (SFNA) in CEPAM: External Adjustment Reporting Tool, Section 1.a – Sacramento NAA 2016 Ozone SIP Ver. 1.04.

#### **Workbook Name: Appendix A1 Emissions By EIC**

Worksheet Name	Worksheet Description
NOX Emissions by EIC	Estimated NO <sub>x</sub> forecast summary by EIC for SFNA in CEPAM Ver. 1.04.
ROG Emissions by EIC	Estimated ROG forecast summary by EIC for SFNA in CEPAM Ver. 1.04.

### Appendix A-2 Growth and Control Data for Emission Forecasting

This Appendix A2 (available separately in electronic file format) contains the growth and control data used for emission forecasting stationary and area-wide sources in CARB's SIP planning projection model, CEPAM.

#### **Workbook Name: #DT0199 SacNA Control Profiles OZ16SIP V100 6FEB2015**

Worksheet Name	Worksheet Description
ReadME	Description of each spreadsheet
Rule_List	List of the control profiles applies to SNA
Rule_Desc	Control rule description table
Control_Data	Control data table
Rule_Desc_Field_Descriptio n	Description of the fields in the Rule_Desc table
Control_Data_Field_Descrip tion	Description of the fields in the Control_Data table

#### **Workbook Name: gap\_sacozone**

Worksheet Name	Worksheet Description
gap_sacozone	Growth Activity Profiles for SFNA

#### **Workbook Name: pad\_sacozone**

Worksheet Name	Worksheet Description
pad_sacozone	Parameter Assignment Data for SFNA

#### **Workbook Name: growthparam update**

---

Worksheet Name	Worksheet Description
Sheet1	Updated name assignments for growth parameters

### **Appendix A-3 Emission Reduction Credits (ERCs)**

This Appendix A3 contains a summary description and inventory of VOC and NO<sub>x</sub> emission reduction credits (ERCs) listed by the individual air districts. In addition, the appendix includes: 1) unused ERCs issued for reductions that occurred prior to the 2012 base year, and 2) future bankable rice burning ERCs. The VOC and NO<sub>x</sub> ERC totals were added to the emission inventory forecast years in chapter 5, Table 5-3 and Table 5-4, respectively.

#### **Unused ERCs Issued for Reductions That Occurred Prior to 2012 Base Year**

Certain pollutant emission reductions due to equipment shutdown or voluntary control may be converted to emission reduction credits (ERCs) and registered with the air districts. These ERCs may then be used as “offsets” to compensate for an increase in emissions from a new or modified major emission source regulated by the air districts. Unused ERCs are considered as potential future emissions supplemental to the forecasted emissions inventory.

The amount of unused ERCs from stationary sources that occurred prior to the 2012 base year are estimated at 4.2 tons per day of VOC and 3.1 tons per day of NO<sub>x</sub> and are summarized by air district in Table A3-1. They are included in the emissions forecasts to ensure the potential future use of these credits does not interfere with reasonable further progress and attainment goals.

#### **Future Bankable Rice Burning Emission Reduction Credits**

Emission credits from reduction in burning may not be used to comply with offset requirements at a new major stationary source or a major modification, unless they are included in an approved attainment demonstration plan. (USEPA Region IX, 2003) To meet this requirement, the impact of accounting for ERCs from reduction in rice straw burning and other agricultural burning credits are being included in this 8-hour ozone attainment and RFP demonstration plan.

California legislation in 1991 (known as the Connelly bill) required rice farmers to phase down rice field burning on an annual basis, beginning in 1992. A burn cap of 125,000 acres in the Sacramento Valley Air Basin was established, and growers with 400 acres or less were granted the option to burn their entire acreage once every four years. Since the rice burning reductions were mandated by state law, they would ordinarily not be “surplus” and eligible for banking. However, the Connelly bill included a special provision declaring that the reductions qualified for banking if they met the State and local banking rules.

The amount of future bankable rice burning ERCs for the Sacramento nonattainment area are estimated at about 0.12 tons per day of VOC and 0.13 tons per day of NO<sub>x</sub> and are listed by air district in Table A3-2. They are included in the emissions forecasts

to ensure the potential future use of these credits does not interfere with reasonable further progress and attainment goals.

Table A-1 Summary of Unused Banked ERCs in the SFNA for 2012 Baseline

Air District <sup>a</sup>	Avg. Summer Day	
	VOC (tpd)	NO <sub>x</sub> (tpd)
Sacramento Metropolitan AQMD	2.24	1.41
Yolo-Solano AQMD	0.41	0.40
Placer County APCD	0.58	0.50
Feather River AQMD (South Sutter)	0.92	0.76
<b>Total Unused Banked ERCs</b>	<b>4.16</b>	<b>3.08</b>

<sup>a</sup> There are no ERCs for El Dorado County AQMD.

Table A-2 Summary of Future Bankable Rice Burning ERCs in the SFNA

Air District <sup>a</sup>	Avg. Summer Day	
	VOC (tpd)	NO <sub>x</sub> (tpd)
Sacramento Metropolitan AQMD	0.12	0.13
Yolo-Solano AQMD	-	-
Placer County APCD	-	-
Feather River AQMD (South Sutter)	-	-
<b>Total Future Rice Burning ERCs</b>	<b>0.12</b>	<b>0.13</b>

<sup>a</sup> The only district with bankable rice ERCs is Sacramento. All other districts have already banked their rice emissions.

The VOC and NO<sub>x</sub> ERC totals from Table A3-1 and A3-2 were rounded up to 5 tons per day VOC and 4 tons per day NO<sub>x</sub> and added to the emission inventory forecast years in Chapter 5, Table 5-3 and Table 5-4, respectively.

## Reference

USEPA Region IX (Broadbent, Jack P.) Message to Larry Greene (YSAQMD) “Re: *Generating Emissions Offsets from Reductions in Rice Straw Burning in Accordance with Health and Safety Code Section 41865.*”, 30 October 2003. Print.

**Appendix A-4 Emissions Inventory Summary from CEPAM**

This Appendix A4 contains 2012, 2018, 2021, 2024, and 2025 VOC and NO<sub>x</sub> emissions inventory summary from CEPAM: External Adjustment Reporting Tool, Section 1.a – Sacramento NAA 2016 Ozone SIP Ver. 1.04.

Table A-3 2012, 2018, 2021, 2024, and 2025 VOC Inventory from CEPAM v1.04

**CEPAM: EXTERNAL ADJUSTMENT REPORTING TOOL  
Emission Projections by Summary Category**

**(Includes approved external emission adjustments)**

**Season: Summer  
Reactive Organic Gas  
Base Year: 2012**

**PRELIMINARY DRAFT: SUBJECT TO CHANGE**

[Download this data as a comma delimited file.](#)  
[Download more detail data as a comma delimited file.](#)

Sacramento NAA 2016 Ozone SIP Ver. 1.04

STATIONARY SOURCES					
SUMMARY CATEGORY NAME	2012	2018	2021	2024	2025
<b>FUEL COMBUSTION</b>					
ELECTRIC UTILITIES	0.208	0.208	0.206	0.212	0.216
COGENERATION	0.001	0.001	0.001	0.001	0.001
OIL AND GAS PRODUCTION (COMBUSTION)	0.041	0.036	0.033	0.031	0.031
MANUFACTURING AND INDUSTRIAL	0.125	0.122	0.123	0.122	0.122
FOOD AND AGRICULTURAL PROCESSING	0.220	0.110	0.097	0.085	0.082
SERVICE AND COMMERCIAL	0.069	0.070	0.071	0.071	0.071
OTHER (FUEL COMBUSTION)	0.097	0.091	0.085	0.086	0.087
* TOTAL FUEL COMBUSTION	0.761	0.637	0.617	0.609	0.609
SUMMARY CATEGORY NAME	2012	2018	2021	2024	2025
<b>WASTE DISPOSAL</b>					
SEWAGE TREATMENT	0.034	0.036	0.037	0.038	0.039
LANDFILLS	0.803	0.846	0.873	0.901	0.911
INCINERATORS	0.007	0.007	0.008	0.008	0.008
SOIL REMEDIATION	0.008	0.008	0.009	0.009	0.009
OTHER (WASTE DISPOSAL)	5.286	4.372	4.540	4.750	4.820
* TOTAL WASTE DISPOSAL	6.138	5.269	5.466	5.706	5.787
SUMMARY CATEGORY NAME	2012	2018	2021	2024	2025
<b>CLEANING AND SURFACE COATINGS</b>					
LAUNDERING	0.080	0.085	0.087	0.090	0.091

DEGREASING	1.953	2.303	2.406	2.564	2.623
COATINGS AND RELATED PROCESS SOLVENTS	2.889	3.207	3.385	3.617	3.702
PRINTING	1.292	1.436	1.466	1.495	1.505
ADHESIVES AND SEALANTS	0.732	0.919	0.975	1.003	1.013
OTHER (CLEANING AND SURFACE COATINGS)	0.231	0.252	0.261	0.274	0.279
* TOTAL CLEANING AND SURFACE COATINGS	7.177	8.200	8.579	9.042	9.213
SUMMARY CATEGORY NAME	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
PETROLEUM PRODUCTION AND MARKETING					
OIL AND GAS PRODUCTION	1.289	1.127	1.055	0.987	0.965
PETROLEUM REFINING	0.000	0.000	0.000	0.000	0.000
PETROLEUM MARKETING	4.579	4.119	3.864	3.630	3.556
OTHER (PETROLEUM PRODUCTION AND MARKETING)	0.005	0.005	0.005	0.004	0.004
* TOTAL PETROLEUM PRODUCTION AND MARKETING	5.873	5.251	4.923	4.621	4.525
SUMMARY CATEGORY NAME	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
INDUSTRIAL PROCESSES					
CHEMICAL	0.620	0.782	0.862	0.951	0.982
FOOD AND AGRICULTURE	0.577	0.655	0.688	0.717	0.726
MINERAL PROCESSES	0.246	0.303	0.318	0.335	0.341
METAL PROCESSES	0.003	0.003	0.003	0.004	0.004
WOOD AND PAPER	0.713	0.773	0.773	0.776	0.780
ELECTRONICS	0.002	0.002	0.002	0.002	0.002
OTHER (INDUSTRIAL PROCESSES)	0.215	0.415	0.462	0.507	0.523
* TOTAL INDUSTRIAL PROCESSES	2.374	2.931	3.108	3.291	3.357
** TOTAL STATIONARY	22.322	22.289	22.693	23.270	23.490
AREAWIDE SOURCES					
SUMMARY CATEGORY NAME	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
SOLVENT EVAPORATION					
CONSUMER PRODUCTS	12.410	12.273	12.632	13.008	13.137
ARCHITECTURAL COATINGS AND RELATED PROCESS SOLVENTS	8.030	8.343	8.583	8.834	8.920
PESTICIDES/FERTILIZERS	1.157	1.223	1.214	1.205	1.203
ASPHALT PAVING / ROOFING	1.001	1.367	1.482	1.538	1.559
* TOTAL SOLVENT EVAPORATION	22.597	23.205	23.911	24.586	24.819
SUMMARY CATEGORY NAME	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
MISCELLANEOUS PROCESSES					
RESIDENTIAL FUEL COMBUSTION	2.029	2.088	2.135	2.183	2.199
FARMING OPERATIONS	2.875	2.875	2.875	2.875	2.875
CONSTRUCTION AND DEMOLITION	0.000	0.000	0.000	0.000	0.000
PAVED ROAD DUST	0.000	0.000	0.000	0.000	0.000
UNPAVED ROAD DUST	0.000	0.000	0.000	0.000	0.000
FUGITIVE WINDBLOWN DUST	0.000	0.000	0.000	0.000	0.000

FIRES	0.037	0.039	0.040	0.042	0.042
MANAGED BURNING AND DISPOSAL	0.818	0.807	0.803	0.801	0.801
COOKING	0.149	0.158	0.163	0.168	0.170
OTHER (MISCELLANEOUS PROCESSES)	0.000	0.000	0.000	0.000	0.000
* TOTAL MISCELLANEOUS PROCESSES	5.908	5.967	6.017	6.070	6.087
** TOTAL AREA WIDE	28.505	29.172	29.928	30.656	30.905
<b>MOBILE SOURCES</b>					
SUMMARY CATEGORY NAME	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
<b>ON-ROAD MOTOR VEHICLES</b>					
LIGHT DUTY PASSENGER (LDA)	12.128	6.044	4.676	3.960	3.809
LIGHT DUTY TRUCKS - 1 (LDT1)	3.844	1.820	1.327	1.032	0.959
LIGHT DUTY TRUCKS - 2 (LDT2)	5.054	3.249	2.745	2.474	2.406
MEDIUM DUTY TRUCKS (MDV)	4.399	3.446	2.911	2.469	2.362
LIGHT HEAVY DUTY GAS TRUCKS - 1 (LHDV1)	1.557	1.066	0.879	0.719	0.671
LIGHT HEAVY DUTY GAS TRUCKS - 2 (LHDV2)	0.151	0.090	0.064	0.045	0.041
MEDIUM HEAVY DUTY GAS TRUCKS (MHDV)	0.589	0.160	0.111	0.083	0.077
HEAVY HEAVY DUTY GAS TRUCKS (HHDV)	0.205	0.033	0.015	0.011	0.010
LIGHT HEAVY DUTY DIESEL TRUCKS - 1 (LHDV1)	0.317	0.242	0.196	0.156	0.145
LIGHT HEAVY DUTY DIESEL TRUCKS - 2 (LHDV2)	0.076	0.059	0.049	0.042	0.040
MEDIUM HEAVY DUTY DIESEL TRUCKS (MHDV)	0.657	0.306	0.103	0.076	0.077
HEAVY HEAVY DUTY DIESEL TRUCKS (HHDV)	1.455	0.343	0.306	0.225	0.225
MOTORCYCLES (MCY)	2.790	2.530	2.488	2.439	2.422
HEAVY DUTY DIESEL URBAN BUSES (UB)	0.139	0.078	0.056	0.040	0.036
HEAVY DUTY GAS URBAN BUSES (UB)	0.054	0.036	0.028	0.023	0.020
SCHOOL BUSES - GAS (SBG)	0.043	0.007	0.006	0.005	0.005
SCHOOL BUSES - DIESEL (SBD)	0.026	0.005	0.004	0.004	0.004
OTHER BUSES - GAS (OBG)	0.046	0.029	0.024	0.019	0.019
OTHER BUSES - MOTOR COACH - DIESEL (OBC)	0.017	0.004	0.004	0.002	0.003
ALL OTHER BUSES - DIESEL (OBD)	0.026	0.004	0.003	0.002	0.002
MOTOR HOMES (MH)	0.058	0.026	0.016	0.011	0.009
* TOTAL ON-ROAD MOTOR VEHICLES	33.633	19.578	16.008	13.835	13.340
SUMMARY CATEGORY NAME	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
<b>OTHER MOBILE SOURCES</b>					
AIRCRAFT	0.483	0.500	0.514	0.526	0.531
TRAINS	0.378	0.330	0.302	0.276	0.266
OCEAN GOING VESSELS	0.003	0.004	0.004	0.004	0.004
COMMERCIAL HARBOR CRAFT	0.106	0.095	0.095	0.095	0.094
RECREATIONAL BOATS	11.722	8.619	7.330	6.181	5.833
OFF-ROAD RECREATIONAL VEHICLES	1.651	1.529	1.462	1.390	1.370
OFF-ROAD EQUIPMENT	7.955	6.305	6.095	6.012	5.991
FARM EQUIPMENT	1.686	1.223	1.047	0.916	0.880



FUEL STORAGE AND HANDLING	1.710	1.371	1.275	1.204	1.185
* TOTAL OTHER MOBILE SOURCES	25.692	19.976	18.123	16.604	16.153
** TOTAL MOBILE	59.325	39.554	34.132	30.439	29.493
GRAND TOTAL FOR SACRAMENTO NAA 2016 OZONE SIP VER. 1.04	<b>2012</b> 110.152	<b>2018</b> 91.015	<b>2021</b> 86.753	<b>2024</b> 84.364	<b>2025</b> 83.888

Notes:

- Migration ID: 2016\_SIP\_V104\_SAC
- AF Migration Table: AF\_MASTERSP16SACOZ104

Report Run time: Started: 05/08/2017 15:22:11 ; Finished: 05/08/2017 15:22:19

Table A-4 2012, 2018, 2021, 2024, and 2025 NOX Inventory from CEPAM v1.04

**CEPAM: EXTERNAL ADJUSTMENT REPORTING TOOL**  
**Emission Projections by Summary Category**

(Includes approved external emission adjustments)

**Season: Summer**  
**Oxides of Nitrogen**  
**Base Year: 2012**

**PRELIMINARY DRAFT: SUBJECT TO CHANGE**

[Download this data as a comma delimited file.](#)  
[Download more detail data as a comma delimited file.](#)

Sacramento NAA 2016 Ozone SIP Ver. 1.04

STATIONARY SOURCES					
SUMMARY CATEGORY NAME	2012	2018	2021	2024	2025
<b>FUEL COMBUSTION</b>					
ELECTRIC UTILITIES	1.237	1.318	1.347	1.401	1.431
COGENERATION	0.008	0.010	0.010	0.011	0.011
OIL AND GAS PRODUCTION (COMBUSTION)	0.069	0.060	0.056	0.053	0.052
MANUFACTURING AND INDUSTRIAL	1.629	1.370	1.426	1.455	1.471
FOOD AND AGRICULTURAL PROCESSING	2.362	1.126	0.997	0.871	0.836
SERVICE AND COMMERCIAL	1.516	1.538	1.551	1.532	1.522
OTHER (FUEL COMBUSTION)	0.682	0.576	0.496	0.498	0.499
* TOTAL FUEL COMBUSTION	7.502	5.998	5.884	5.822	5.823
SUMMARY CATEGORY NAME	2012	2018	2021	2024	2025
<b>WASTE DISPOSAL</b>					
SEWAGE TREATMENT	0.002	0.002	0.002	0.002	0.002
LANDFILLS	0.038	0.039	0.040	0.040	0.041
INCINERATORS	0.019	0.021	0.022	0.023	0.024
SOIL REMEDIATION	0.000	0.000	0.000	0.000	0.000
OTHER (WASTE DISPOSAL)	0.000	0.000	0.000	0.000	0.000
* TOTAL WASTE DISPOSAL	0.058	0.062	0.064	0.066	0.067
SUMMARY CATEGORY NAME	2012	2018	2021	2024	2025
<b>CLEANING AND SURFACE COATINGS</b>					
LAUNDERING	0.000	0.000	0.000	0.000	0.000
DEGREASING	0.000	0.000	0.000	0.000	0.000
COATINGS AND RELATED PROCESS SOLVENTS	0.008	0.011	0.011	0.012	0.013
PRINTING	0.004	0.005	0.005	0.005	0.005
ADHESIVES AND SEALANTS	0.000	0.000	0.000	0.000	0.000

OTHER (CLEANING AND SURFACE COATINGS)	0.000	0.000	0.000	0.000	0.000
* TOTAL CLEANING AND SURFACE COATINGS	0.012	0.015	0.016	0.017	0.018
<b>SUMMARY CATEGORY NAME</b>	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
<b>PETROLEUM PRODUCTION AND MARKETING</b>					
OIL AND GAS PRODUCTION	0.001	0.001	0.001	0.001	0.001
PETROLEUM REFINING	0.000	0.000	0.000	0.000	0.000
PETROLEUM MARKETING	0.010	0.011	0.010	0.010	0.010
OTHER (PETROLEUM PRODUCTION AND MARKETING)	0.000	0.000	0.000	0.000	0.000
* TOTAL PETROLEUM PRODUCTION AND MARKETING	0.012	0.012	0.011	0.011	0.010
<b>SUMMARY CATEGORY NAME</b>	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
<b>INDUSTRIAL PROCESSES</b>					
CHEMICAL	0.115	0.144	0.158	0.175	0.180
FOOD AND AGRICULTURE	0.015	0.017	0.018	0.018	0.019
MINERAL PROCESSES	0.359	0.443	0.465	0.490	0.499
METAL PROCESSES	0.008	0.009	0.009	0.010	0.010
WOOD AND PAPER	0.041	0.045	0.045	0.045	0.045
ELECTRONICS	0.000	0.000	0.000	0.000	0.000
OTHER (INDUSTRIAL PROCESSES)	0.014	0.027	0.030	0.033	0.034
* TOTAL INDUSTRIAL PROCESSES	0.553	0.684	0.725	0.771	0.787
** TOTAL STATIONARY	8.137	6.771	6.700	6.686	6.705
<b>AREAWIDE SOURCES</b>					
<b>SUMMARY CATEGORY NAME</b>	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
<b>SOLVENT EVAPORATION</b>					
CONSUMER PRODUCTS	0.000	0.000	0.000	0.000	0.000
ARCHITECTURAL COATINGS AND RELATED PROCESS SOLVENTS	0.000	0.000	0.000	0.000	0.000
PESTICIDES/FERTILIZERS	0.000	0.000	0.000	0.000	0.000
ASPHALT PAVING / ROOFING	0.000	0.000	0.000	0.000	0.000
* TOTAL SOLVENT EVAPORATION	0.000	0.000	0.000	0.000	0.000
<b>SUMMARY CATEGORY NAME</b>	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
<b>MISCELLANEOUS PROCESSES</b>					
RESIDENTIAL FUEL COMBUSTION	2.389	1.974	1.851	1.847	1.844
FARMING OPERATIONS	0.000	0.000	0.000	0.000	0.000
CONSTRUCTION AND DEMOLITION	0.000	0.000	0.000	0.000	0.000
PAVED ROAD DUST	0.000	0.000	0.000	0.000	0.000
UNPAVED ROAD DUST	0.000	0.000	0.000	0.000	0.000
FUGITIVE WINDBLOWN DUST	0.000	0.000	0.000	0.000	0.000
FIRES	0.013	0.014	0.014	0.015	0.015
MANAGED BURNING AND DISPOSAL	0.300	0.291	0.288	0.286	0.286
COOKING	0.000	0.000	0.000	0.000	0.000
OTHER (MISCELLANEOUS PROCESSES)	0.000	0.000	0.000	0.000	0.000

* TOTAL MISCELLANEOUS PROCESSES	2.702	2.279	2.153	2.148	2.144
** TOTAL AREA WIDE	2.702	2.279	2.153	2.148	2.144
<b>MOBILE SOURCES</b>					
SUMMARY CATEGORY NAME	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
<b>ON-ROAD MOTOR VEHICLES</b>					
LIGHT DUTY PASSENGER (LDA)	6.505	3.311	2.438	1.897	1.764
LIGHT DUTY TRUCKS - 1 (LDT1)	1.653	0.687	0.449	0.311	0.278
LIGHT DUTY TRUCKS - 2 (LDT2)	4.184	2.169	1.577	1.213	1.120
MEDIUM DUTY TRUCKS (MDV)	4.695	2.671	1.861	1.292	1.158
LIGHT HEAVY DUTY GAS TRUCKS - 1 (LHDV1)	1.885	1.174	0.920	0.703	0.642
LIGHT HEAVY DUTY GAS TRUCKS - 2 (LHDV2)	0.203	0.127	0.097	0.072	0.065
MEDIUM HEAVY DUTY GAS TRUCKS (MHDV)	0.564	0.281	0.194	0.134	0.120
HEAVY HEAVY DUTY GAS TRUCKS (HHDV)	0.192	0.077	0.060	0.054	0.053
LIGHT HEAVY DUTY DIESEL TRUCKS - 1 (LHDV1)	7.328	4.547	3.292	2.306	2.036
LIGHT HEAVY DUTY DIESEL TRUCKS - 2 (LHDV2)	1.654	0.918	0.610	0.381	0.320
MEDIUM HEAVY DUTY DIESEL TRUCKS (MHDV)	7.922	5.133	3.359	2.972	3.029
HEAVY HEAVY DUTY DIESEL TRUCKS (HHDV)	19.743	11.176	9.606	6.329	6.221
MOTORCYCLES (MCY)	0.535	0.480	0.465	0.455	0.453
HEAVY DUTY DIESEL URBAN BUSES (UB)	2.035	1.088	0.778	0.571	0.516
HEAVY DUTY GAS URBAN BUSES (UB)	0.110	0.082	0.067	0.056	0.052
SCHOOL BUSES - GAS (SBG)	0.042	0.012	0.009	0.007	0.006
SCHOOL BUSES - DIESEL (SBD)	0.357	0.306	0.257	0.209	0.194
OTHER BUSES - GAS (OBG)	0.122	0.071	0.053	0.040	0.038
OTHER BUSES - MOTOR COACH - DIESEL (OBC)	0.234	0.144	0.119	0.062	0.064
ALL OTHER BUSES - DIESEL (OBD)	0.311	0.141	0.108	0.061	0.062
MOTOR HOMES (MH)	0.273	0.170	0.124	0.089	0.081
* TOTAL ON-ROAD MOTOR VEHICLES	60.545	34.763	26.442	19.213	18.271
SUMMARY CATEGORY NAME	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
<b>OTHER MOBILE SOURCES</b>					
AIRCRAFT	1.409	1.430	1.507	1.579	1.605
TRAINS	6.158	6.652	6.317	5.859	5.698
OCEAN GOING VESSELS	0.086	0.074	0.067	0.062	0.061
COMMERCIAL HARBOR CRAFT	1.453	0.979	0.921	0.876	0.866
RECREATIONAL BOATS	2.260	1.937	1.818	1.722	1.691
OFF-ROAD RECREATIONAL VEHICLES	0.046	0.057	0.063	0.068	0.069
OFF-ROAD EQUIPMENT	10.027	7.856	6.747	5.878	5.524
FARM EQUIPMENT	8.322	6.599	5.618	4.674	4.402
FUEL STORAGE AND HANDLING	0.000	0.000	0.000	0.000	0.000
* TOTAL OTHER MOBILE SOURCES	29.760	25.584	23.058	20.718	19.915
** TOTAL MOBILE	90.305	60.346	49.500	39.931	38.186

GRAND TOTAL FOR SACRAMENTO NAA 2016 OZONE SIP VER. 1.04	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>	<b>2025</b>
	101.145	69.396	58.354	48.766	47.035

Notes:

- Migration ID: 2016\_SIP\_V104\_SAC
- AF Migration Table: AF\_MASTERSP16SAC0Z104

Report Run time: Started: 05/08/2017 16:20:17 ; Finished: 05/08/2017 16:20:25

### Appendix A-5 EMFAC2014 Output Data

**Appendix A5** contains the on-road motor vehicle emissions, vehicle population, and activity data generated using EMFAC2014, and includes updated activity data for Solano County from MTC.

#### **Workbook Name: Sacramento Non Attainment Area CEPAM V1.04 Emissions - May 10 2017**

Worksheet Name	Worksheet Description
ReadME	Description of each spreadsheet
El Dorado (MC)	EMFAC output files for Mountain Counties Air Basin of El Dorado County, consistent with CEPAM V1.04
Placer (MC)	EMFAC output files for Mountain Counties Air Basin of Placer County, consistent with CEPAM V1.04
Placer (SV)	EMFAC output files for Sacramento Valley Air Basin of Placer County, consistent with CEPAM V1.04
Sacramento (SV)	EMFAC output files for Sacramento Valley Air Basin of Sacramento County, consistent with CEPAM V1.04
South Sutter	EMFAC output files for the South Sutter portion of Sacramento Valley Air Basin of Sutter County, consistent with CEPAM V1.04
Yolo (SV)	EMFAC output files Sacramento Valley Air Basin of Yolo County, consistent with CEPAM V1.04
Solano (SV)	EMFAC output files for Sacramento Valley Air Basin of Solano County, consistent with CEPAM V1.04

## APPENDIX B

### Photochemical Modeling

## Appendix B Photochemical Modeling

The 2008 Ozone National Ambient Air Quality Standard (NAAQS) implementation rule (80 FR 12264) requires that an area classified as serious or higher to demonstrate attainment by means of a photochemical grid model or any other analytical method (40 CFR 51.1108).

Appendix B contains a summary and documentation regarding the photochemical grid modeling performed by the California Air Resources Board (CARB) in evaluating and supporting the attainment demonstration for the 2008 ozone NAAQS in the Sacramento Federal Nonattainment Area (SFNA). CARB prepared this appendix includes five sections:

1. summary of the modeling results,
2. conceptual modeling,
3. modeling protocol,
4. modeling attainment demonstration, and
5. modeling emissions inventory.

## References

- EPA. *Modeling and attainment demonstration requirements*. 40 CFR §51.1108.
- EPA. (80 FR 12264 - 12319) *Implementation of the 2008 National Ambient Air Quality Standards for Ozone: State Implementation Plan Requirements; Final Rule*. Federal Register, Volume 80, 6 March 2015, p. 12264 – 12319. Print.

This appendix included the following subsections:

<b>Appendix</b>	<b>Supporting Document Title</b>	<b>Description</b>
B-1	<i>Modeling 8-Hour Ozone for the Sacramento Federal Nonattainment Area's 2016 State Implementation Plan for the 75ppb 8-Hour Ozone Standard</i>	This provides a summary of the photochemical modeling results.
B-2	<i>Sacramento Federal Non-attainment Area (SFNA) 0.075 ppm 8-hour Ozone (2016)</i>	This appendix provides a description of the conceptual model for the SFNA
B-3	<i>Photochemical Modeling Protocol – Photochemical Modeling for the 8-Hour Ozone and Annual/24-hour PM<sub>2.5</sub> State Implementation Plans</i>	The modeling protocol includes the details and procedures for conducting the photochemical modeling that forms the basis of the attainment demonstration for the State Implementation Plans (SIPs) for California.
B-4	<i>Modeling Attainment Demonstration – Photochemical Modeling for the 8-Hour Ozone State Implementation Plan in the Sacramento Federal Non-attainment Area (SFNA)</i>	The modeling attainment demonstration document provides the details of the modeling results for the 2008 Ozone NAAQS in the SFNA, which forms the scientific basis for the attainment demonstration.
B-5	<i>Modeling Emission Inventory for the 8-Hour Ozone State Implementation Plan in the Sacramento Non-Attainment Area</i>	This document describes how the base and future year gridded photochemical modeling emissions inventory are prepared.



## Appendix B-1 Modeling 8-Hour Ozone for the Sacramento Federal Nonattainment Area's 2016 SIP for the 75ppb 8-Hour Ozone Standard

Photochemical modeling plays a crucial role in the SIP process to demonstrate attainment of air quality standards based on estimated future emissions and for the development of emissions targets necessary for attainment. Currently, the SFNA is designated as a severe ozone non-attainment area for the 2008 0.075 ppm (or 75 ppb) 8-hour ozone standard and is required to demonstrate attainment of this standard by 2026. Consistent with U.S. EPA guidelines for model attainment demonstrations<sup>1</sup>, photochemical modeling was used to estimate the future year 2026 ozone (O<sub>3</sub>) design values (DVs) at each monitoring site in the SFNA in order to show attainment of the standard by 2026. An additional future year 2022 was also modeled to assess progress toward the 2026 attainment deadline.

The findings of the SFNA's model attainment demonstration are summarized below. Additional information and a detailed description of the procedures employed in this modeling are available in the Modeling Attainment Demonstration Appendix (Appendix B-4) and Modeling Protocol Appendix (Appendix B-3).

The current modeling platform draws on the products of large-scale, scientific studies in the region, collaboration among technical staff of state, local, and federal regulatory agencies, as well as from participation in technical and policy groups within the region (see Modeling Protocol (Appendix B-3) for further details). In this modeling work, the Weather Research and Forecasting (WRF) numerical model version 3.6 was utilized to generate meteorological fields, while the Community Multiscale Air Quality (CMAQ) Model version 5.0.2 was used for modeling ozone in the SFNA. Other relevant information, including the modeling domain definition, chemical mechanism, initial and boundary conditions, and emissions preparation can be found in the Modeling Protocol and Modeling Emissions Inventory Appendices (Appendix B-5).

Based on U.S. EPA modeling guidance<sup>1</sup>, modeling was used in a relative sense to project observed DVs to the future. The year 2012 was chosen as the starting point for the modeling and reference (or baseline) DV calculation based on analysis regarding the conduciveness of recent years' meteorological conditions to enhanced ozone formation and the availability of the most detailed emissions inventory. These reference DVs serve as the anchor point for estimating future year projected design values. The year 2026 was the future year modeled in this attainment demonstration since that is the year for which attainment must be demonstrated. An additional future year (2022)

---

<sup>1</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM<sub>2.5</sub> and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

was also modeled to assess progress toward the attainment of the 75 ppb standard by the stipulated deadline (2026).

DVs are the three-year average of the annual 4<sup>th</sup> highest 8-hour O<sub>3</sub> mixing ratio observed at each monitor, and are used to determine compliance with the standard. In the attainment demonstration, the U.S. EPA recommends using an average of three DVs to account for the year-to-year variability in meteorology, so DVs were calculated for the three year period ending in 2012, 2013, and 2014 and then the three DVs were averaged. This average DV is called a baseline DV (see the 2<sup>nd</sup> column of Table 2 for the baseline DVs utilized in the attainment demonstration modeling).

In order to use the modeling in a relative sense, three simulations were conducted: 1) base year simulation for 2012, which was used to verify that the model reasonably reproduced the observed air quality; 2) reference year simulation for 2012, which was the same as the base year simulation, but excluded exceptional event emissions such as wildfires; 3) future year simulations for 2022 and 2026, which were the same as the reference year simulation, except that projected anthropogenic emissions for 2022 and 2026 were used in lieu of the 2012 emissions.

Table B-1 summarizes the 2012, 2022, and 2026 SFNA anthropogenic emissions used in the attainment demonstration modeling. Overall, anthropogenic NO<sub>x</sub> was projected to decrease ~45% by 2022 (from 104 tpd to 56.8 tpd) and ~55% by 2026 (from 104 tpd to 47.3 tpd) when compared to the 2012 emissions levels. In contrast, anthropogenic ROG was projected to decrease ~23 % by 2022 (from 109.7 tpd to 84.7 tpd) and ~26 % by 2026 (from 109.8 tpd to 81.7 tpd). Biogenic ROG emissions were held constant between all simulations with summer average (May – September, 2012) emissions estimated at ~693 tpd for the SFNA.

Table B-1 Summer emission inventory totals (CEPAM v1.03) for 2012, 2022 and 2026. Biogenic emission totals were averaged over May – September, 2012.

Source Category	NO <sub>x</sub>					ROG						
	2012		2022		2026		2012		2022		2026	
	[tpd]	[tpd]	% diff <sup>#</sup>	[tpd]	% diff <sup>#</sup>	[tpd]	[tpd]	% diff <sup>#</sup>	[tpd]	% diff <sup>#</sup>	[tpd]	% diff <sup>#</sup>
Stationary	9.2	7.6	-17	7.6	-17	20.6	21.9	6	22.1	7		
Area	2.7	2.1	-22	2.1	-22	28.5	29.5	4	30.4	7		
On-Road												
Mobile	62	24.5	-60	17.7	-71	35	15.6	-55	13.3	-62		
Other												
Mobile	30.1	22.6	-25	19.9	-34	25.7	17.7	-31	15.9	-38		
Total	104	56.8	-45	47.3	-55	109.8	84.7	-23	81.7	-26		
Biogenic			--				693	--	693	--		

<sup>#</sup>% diff denotes percent difference with respect to 2012 emission levels.

As part of the model attainment demonstration, the fractional changes in ozone mixing ratios between the model reference year (2012) and the two model future years (2022 and 2026) were calculated separately at each of the monitors following the U.S. EPA modeling guidance<sup>2</sup> and procedures outlined in the Modeling Protocol Appendix. These ratios, called “relative response factors” or RRFs, were calculated based on the ratio of future year modeled maximum daily average 8-hour (MDA8) ozone to modeled reference year MDA8 ozone (Equation 1).

$$\text{RRF} = \frac{\text{average MDA8 ozone}_{\text{future}}}{\text{average MDA8 ozone}_{\text{reference}}} \quad (1)$$

The site-specific RRF for each of the future years 2022 and 2026 was then multiplied by the weighted DV for the corresponding monitor to predict the future year 2022 and 2026 DVs (Table 2). The RRF approach was previously applied in the 2009 Sacramento 8-Hour Ozone SIP<sup>3</sup> where the emission targets in SFNA were appropriately characterized for attaining the 1997 federal 8-hour ozone standard of 0.08 ppm (or 84 ppb) by 2018. The RRF approach has been applied in other regions of California’s Central valley including the SJV for the 2007 8-hour Ozone SIP<sup>4</sup> and later in the 2013 1-hour Ozone SIP<sup>5</sup>. In addition, two peer-reviewed scientific publications focused primarily on areas outside of California (one from researchers at Rice University<sup>6</sup> and one from U.S. EPA scientists<sup>7</sup>), both found that the RRF approach is highly robust in its ability to predict future DVs.

Table B-2 shows that all monitoring sites in the SFNA are projected to have a future DV less than 75 ppb, so that the entire region is projected to attain the 75 ppb 8-hour O<sub>3</sub> standard by 2026 based on the substantial emission reductions from implementation of the current control program. The projected 2022 and 2026 DVs for sites in SFNA show a large decrease when compared to 2012 levels (e.g., at the Folsom monitor, the

---

<sup>2</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM2.5 and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

<sup>3</sup> 2009 Sacramento Regional 8-Hour Ozone Attainment and Reasonable Further Progress Plan, available at [http://www.airquality.org/ProgramCoordination/Documents/4\)%202013%20SIP%20Revision%20Report%201997%20Std.pdf](http://www.airquality.org/ProgramCoordination/Documents/4)%202013%20SIP%20Revision%20Report%201997%20Std.pdf)

<sup>4</sup> 2007 Plan for the 1997 8-Hour Ozone Standard available at [http://www.valleyair.org/Air\\_Quality\\_Plans/AQ\\_Final\\_Adopted\\_Ozone2007.htm](http://www.valleyair.org/Air_Quality_Plans/AQ_Final_Adopted_Ozone2007.htm)

<sup>5</sup> 2013 Plan for the Revoked 1-Hour Ozone Standard available at [http://www.valleyair.org/Air\\_Quality\\_Plans/Ozone-OneHourPlan-2013.htm](http://www.valleyair.org/Air_Quality_Plans/Ozone-OneHourPlan-2013.htm)

<sup>6</sup> Pegues, A.H., D.S. Cohan, A. Digar, C. Douglass, and R.S. Wilson (2012). Efficacy of recent state implementation plans for 8-hour ozone. *Journal of the Air & Waste Management Association*, 62, 252-261, doi: 10.1080/10473289.2011.646049.

<sup>7</sup> Foley, K., P. Dolwick, C. Hogrefe, H. Simon, B. Timin, and N. Possiel, (2015), Dynamic evaluation of CMAQ part II: Evaluation of relative response factor metrics for ozone attainment demonstrations, *Atmospheric Environment*, 103: 188–195, doi:10.1016/j.atmosenv.2014.12.039

SFNA's site with the highest baseline DV, the baseline DV declines by ~15 ppb in 2022 and ~20 ppb in 2026 compared to 2012), which is consistent with the peer-reviewed, published study conducted by the UC Berkeley researchers on the observed response of ozone to NO<sub>x</sub> reductions in the Sacramento area<sup>8</sup>. This study concluded that the region's 1-hour ozone exceedance days have been decreasing linearly with decreases in NO<sub>x</sub> suggesting that cumulative NO<sub>x</sub> controls over time have successfully transitioned the SFNA into a NO<sub>x</sub>-limited chemistry regime where NO<sub>x</sub> emission reductions have been becoming increasingly effective at reducing ozone. This is also supported by the analysis on the changes in weekday vs. weekend ozone in the SFNA presented in the Modeling Protocol and Model Attainment Demonstration Appendices, where ozone on weekends is now generally lower than ozone on weekdays (in contrast to higher weekend ozone in the past), which indicates the prevalence of a NO<sub>x</sub>-limited chemical regime in this region.

As part of the attainment demonstration, the U.S. EPA<sup>9</sup> also requires analysis of ozone levels outside of the routine monitoring network (i.e., at areas between the monitors) to ensure that all regions within the SFNA (even those without a monitor) are in attainment of the standard. This "unmonitored area" analysis combines measurement based DVs with model based RRFs and ozone spatial gradients to estimate future 2026 DVs in unmonitored areas. Details of how the unmonitored area analysis is performed can be found in the Modeling Protocol and Model Attainment Demonstration Appendices. The unmonitored area analysis in the SFNA showed that the areas with the highest future DVs were captured within the existing monitoring network and that all areas are projected to achieve the 75 ppb ozone standard.

---

<sup>8</sup> LaFranchi, B. W., Goldstein, A. H., and Cohen, R. C.: Observations of the temperature dependent response of ozone to NO<sub>x</sub> reductions in the Sacramento, CA urban plume, *Atmos. Chem. Phys.*, 11, 6945-6960, doi:10.5194/acp-11-6945-2011, 2011

<sup>9</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM<sub>2.5</sub> and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

Table B-2 Baseline Design Value, modeled RRF, and projected future year (2022 and 2026) Design Value for sites in the SFNA.

Site	Baseline 2012	Future year 2022		Future year 2026	
	Average DV (ppb)	RRF	Average DV (ppb)	RRF	Average DV (ppb)
Folsom-Natoma Street	90.0	0.8358	75	0.7857	70
Sloughhouse	84.0	0.8459	71	0.7998	67
Placerville-Gold Nugget Way	82.3	0.8259	68	0.7778	64
Roseville-N Sunrise Ave	82.3	0.8487	69	0.8055	66
Cool-Hwy193	81.3	0.8336	67	0.7882	64
Auburn - Atwood Rd	79.0	0.8180	64	0.7669	60
Sacramento-Del Paso Manor	77.3	0.8595	66	0.8162	63
North Highlands-Blackfoot Way	76.0	0.8578	65	0.8149	61
Colfax-City Hall	73.7	0.8270	60	0.7804	57
Elk Grove - Bruceville Road	71.7	0.8558	61	0.8129	58
Sacramento - 1309 T Street	70.0	0.8644	60	0.8242	57
Sacramento-Goldenland Court	70.0	0.8820	61	0.8415	58
Echo Summit	69.0	0.9411	64	0.9260	63
Woodland-Gibson Road	68.7	0.8459	58	0.7996	54
Vacaville-Ulatis Drive	67.3	0.8459	56	0.8009	53
Davis-UCD Campus	66.7	0.8495	56	0.8052	53

## Appendix B-2

## Modeling Conceptual Model

### Document Title:

Sacramento Federal Non-attainment Area (SFNA) 0.075 ppm 8-hour Ozone (2016)

### Document Description:

This document provides conceptual modeling for the SFNA. It includes the description of the history of ambient ozone field studies, ambient air monitoring network, ozone trends, and meteorological conditions that leading to SFNA ozone exceedances.

**APPENDIX: Sacramento Federal Non-attainment Area (SFNA)  
0.075 ppm 8-hour Ozone (2016)**

## Table of Contents

1. TIMELINE OF THE PLAN.....	7
2. DESCRIPTION OF THE CONCEPTUAL MODEL FOR THE NONATTAINMENT AREA .....	8
2.1 History of Field Studies in the Region .....	8
2.2 Description of the Ambient Monitoring Network .....	13
2.3 Ozone Trends and Sensitivity to Emissions Reductions .....	19
2.4 Meteorological Conditions Leading to Ozone Exceedances .....	26
REFERENCES.....	30



## LIST OF TABLES

Table 1-1 Timeline for Completion of the Plan .....	7
Table 2-1. Major Field Studies in Central California and surrounding areas. ....	11
Table 2-2. Ozone, NO <sub>x</sub> , and PAMS monitoring sites between 2012 and 2015 in the Sacramento Federal 8-hour ozone Non-attainment Area .....	18

## LIST OF FIGURES

Figure 2-1 Map of California's Central Valley and the geographical location of Sacramento Federal 8-hr Ozone Non-attainment Area (SFNA).....	14
Figure 2-2. Map of the Monitoring Sites in the Sacramento Federal 8-hour Ozone Non-attainment Area. The green, blue and magenta circle markers denote the location of NO <sub>x</sub> /NO <sub>y</sub> , ozone and PAMS monitors (top panel). The solid black line denotes the regional boundary of the SFNA, while the grey line denotes the county boundaries. The dashed brown lines (bottom panel) show the approximate regional boundaries of the Western, Central and Eastern sub-regions of SFNA.....	17
Figure 2-3. Illustrates a typical ozone isopleth plot, where each line represents ozone mixing ratio, in 10 ppb increments, as a function of initial NO <sub>x</sub> and VOC (or ROG) mixing ratio (adapted from Seinfeld and Pandis, 1998, Figure 5.15). General chemical regimes for ozone formation are shown as NO <sub>x</sub> -disbenefit (red circle), transitional (blue circle), and NO <sub>x</sub> -limited (green circle). .....	20
Figure 2-4. Trends in SFNA emissions (top), 8-hour ozone design value (middle), and number of days above the 8-hour ozone standard between 2000 and 2014.....	22
Figure 2-5. Site-specific average weekday and weekend maximum daily average 8-hour ozone for each year from 2000 to 2014 for the Western (top), Central (middle), and Eastern (bottom) sub-regions of SFNA. Points falling below the 1:1 dashed line represent a NO <sub>x</sub> -disbenefit regime, those on the 1:1 dashed line represent a transitional regime, and those above the 1:1 dashed line represent a NO <sub>x</sub> -limited regime. MDA denotes maximum daily average.....	24
Figure 2-6 Conceptual low-level wind patterns in Central California during the day (left panel) and night (right panel) for typical ozone episode conditions (adapted from Bao et al., 2008). .....	26

## ACRONYMS

ACHEX - Aerosol Characterization Experiment

ARCTAS-CARB – California portion of the Arctic Research of the Composition of the Troposphere from Aircraft and Satellites conducted in 2008

BEARPEX – Biosphere Effects on Aerosols and Photochemistry Experiment in 2007 and 2009

CABERNET – California Airborne BVOC Emission Research in Natural Ecosystem Transects in 2011

CalNex – Research at the Nexus of Air Quality and Climate Change conducted in 2010

CARB – California Air Resources Board

CARES – Carbonaceous Aerosols and Radiative Effects Study in 2010

CCOS - Central California Ozone Study

CIRPAS - Center for Interdisciplinary Remotely-Piloted Aircraft Studies

CRPAQS - California Regional PM<sub>10</sub>/PM<sub>2.5</sub> Air Quality Study

DISCOVER-AQ - Deriving Information on Surface Conditions from Column and Vertically Resolved Observations Relevant to Air Quality

DV – Design Value

IMS-95 – Integrated Monitoring Study of 1995

IONS – Intercontinental transport experiment Ozonesonde Network Study)

LIDAR – Light Detection And Ranging

MCAB – Mountain Counties Air Basin

MDA – Maximum Daily Average

NASA – National Aeronautics and Space Administration

NOAA - National Oceanic and Atmospheric Administration

NO<sub>x</sub> – Oxides of nitrogen

PAMS – Photochemical Assessment Monitoring Stations

PAN – Peroxy Acetyl Nitrate

PM<sub>2.5</sub> – Particulate Matter with aerodynamic diameter less than 2.5 micrometers

PM<sub>10</sub> – Particulate Matter with aerodynamic diameter less than 10 micrometers

ROG – Reactive Organic Gases

SAOS – Sacramento Area Ozone Study

SARMAP – SJVAQS/AUSPEX Regional Modeling Adaptation Project

SFNA – Sacramento Federal Non-attainment Area

SIP – State Implementation Plan

SJV – San Joaquin Valley

SJVAB – San Joaquin Valley Air Basin (SJVAB)

SJVAQS/AUSPEX – San Joaquin Valley Air Quality Study/Atmospheric Utilities  
Signatures Predictions and Experiments

SVAB – Sacramento Valley Air Basin (SJVAB)

SOA – Secondary Organic Aerosol

SoCAB – Southern California Air Basin

U.S. EPA – United States Environmental Protection Agency

VOC – Volatile Organic Compounds

WRF Model – Weather and Research Forecast Model

## 1. TIMELINE OF THE PLAN

Table 1-1 Timeline for Completion of the Plan

<b>Timeline</b>	<b>Action</b>
Spring 2016	Emission Inventory Completed
Summer 2016	Modeling Completed
March/April 2017	Sacramento Federal Non-attainment Area (SFNA) Governing Board Hearing to consider the Draft Plan
May 2017	ARB Board Hearing to consider the Sacramento Federal Non-attainment Area Adopted Plan
June 2017	Plan submitted to U.S. EPA

## 2. DESCRIPTION OF THE CONCEPTUAL MODEL FOR THE NONATTAINMENT AREA

### 2.1 History of Field Studies in the Region

The Sacramento Federal 8-hour ozone Non-attainment Area (SFNA) is located in the northern part of California's Central Valley (Figure 2-1), which is a 500-mile long northwest-southeast oriented valley encompassing two of the worst polluted air basins in the nation, the San Joaquin Valley and Sacramento Valley air basins. As a result, California's Central Valley is one of the most studied regions in the world, in terms of the number of publications in peer-reviewed international scientific/technical journals and other major reports. The Major Field studies that have taken place in California's Central Valley and surrounding areas are listed in Table 2-1.

The first major air quality study in the Central Valley, dubbed Project Lo-Jet, took place in 1970 and resulted in the identification of the San Joaquin Valley "Fresno" Eddy and the Sacramento "Schultz" Eddy (Lin and Jao, 1995 and references therein). The first study in the Sacramento region that formed the foundation for a State Implementation Plan (SIP) was the Sacramento Area Ozone Study (SAOS) conducted in July–August, 1990 (Roberts et al., 1990). The timing of the SAOS coincided with the San Joaquin Valley Air Quality Study/Atmospheric Utilities Signatures Predictions and Experiments (SJVAQS/AUSPEX) study, also known as SARMAP (SJVAQS/AUSPEX Regional Modeling Adaptation Project). The 1990 SAOS study was part of the technical basis for the 1-hour Extreme Ozone Attainment Demonstration Plan that was submitted to the U.S. EPA in 1994 (<https://www.arb.ca.gov/planning/sip/94sip/94sip.htm>) and was approved in 1997 (62 FR 1150). The next major study was the Integrated Monitoring Study in 1995 (IMS-95), which was the pilot study for the subsequent California Regional PM<sub>10</sub>/PM<sub>2.5</sub> Air Quality Study (CRPAQS) in 2000 (Solomon and Magliano, 1998). CRPAQS was the first annual field campaign in the Central Valley, and embedded in it was the Central California Ozone Study (CCOS) that took place during the summer of 2000 (Fujita et al., 2001). The CCOS was part of the technical basis for the 2009 Sacramento Regional 8-Hour Ozone Attainment and Reasonable Further Progress Plan for attaining the 1997 federal 8-hour ozone standard of 0.08 ppm, which was approved by EPA in 2014 (79 FR 61799). While CCOS is still very relevant to the current 8-hour O<sub>3</sub> SIP, there are five subsequent studies which are highly relevant to ozone formation in the Central Valley and surrounding regions: 1) ARCTAS-CARB 2008, 2) CalNex 2010, 3) CARES 2010, 4) BEARPEX 2007 & 2009, and 5) CABERNET 2011. Each of these studies has contributed significantly to our understanding of various atmospheric processes in the Central Valley including the SFNA region.

The ARCTAS-CARB aircraft field campaign, a joint research effort by NASA and CARB, took place from June 18 to 24, 2008 with specific objectives to improve the emission inventories of greenhouse gases/aerosols and characterize upwind boundary conditions for modeling surface ozone and PM<sub>2.5</sub>. During the ARCTAS campaign, large wildfires occurred in California, particularly in northern California. The DC-8 aircraft encountered many of the fire plumes, which helped improve the characterization of California wildfires and their chemical composition. The ARCTAS-CARB campaign provided a unique dataset for evaluating the impacts of wildfires on ozone levels through photochemical modeling studies and for evaluating the distribution of reactive nitrogen species in California (Huang et al., 2011; Cai et al., 2016).

The improved understanding of California wildfires is valuable not only from a regulatory modeling standpoint, but it can also provide helpful information relevant to the U.S. EPA's Exceptional Events Rule (72 FR 13560). For instance, the U.S. EPA approved the SFNA's request to classify 1-hour ozone exceedances for June 23, June 27 and July 10 in 2008 as Exceptional Events that were caused by wildfires prevalent in the region from June 21, 2008 through August 11, 2008 (<https://www.arb.ca.gov/desig/excevents/2008wildfires.htm>). The exclusion of these 1-hour ozone exceedance days subsequently lead to a "clean data determination" in SFNA for the revoked 1-hour ozone NAAQS based on 2007-2009 ozone monitoring data (77 FR 64036).

The CalNex May-July 2010 field campaign was organized by NOAA (NOAA, 2014) and CARB. The focus of this field study included airborne measurements using the NOAA WP-3D aircraft and the Twin Otter Remote Sensing aircraft, and surface measurements using the R/V *Atlantis* mobile platform as well as two stationary ground supersites. Overall, the CalNex study provided a comprehensive snapshot of air quality in California and indicated remarkable improvement in air quality over the past few decades. The CalNex data analysis helped in improving emissions estimates from various sources in California, including emissions from rice cultivation in the Sacramento Valley. In addition, analysis of the data collected during CalNex has shown that photochemical ozone production in the southern and central portions of the SJV have transitioned to a NO<sub>x</sub>-limited chemistry regime, where further NO<sub>x</sub> reductions are expected to lead to a more rapid reduction in ozone than was observed over the past decade or more, while the northern portion of the SJV (to the south of the SFNA) is transitioning to a NO<sub>x</sub>-limited regime (Pusede and Cohen, 2012). Studies have also shown that there is evidence for an unidentified temperature-dependent VOC emissions source on the hottest days (Pusede and Cohen, 2012; Pusede et al., 2014) and large sources of hydrocarbon compounds from petroleum extraction/processing, dairy (and other cattle) operations, and agricultural crops in the SJV (Gentner et al., 2014a,b).

The CARES field campaign coincided with CalNex, and took place to the northeast of Sacramento in June 2010. Comprehensive data sets of trace gases and aerosols were taken from the daily evolving Sacramento urban plume under relatively well-defined and regular meteorological conditions using multiple suites of ground-based and airborne instruments onboard the Gulfstream (G-1) research aircraft. The ground-based measurements were conducted at two sites: one within the Sacramento urban source area and the other in a downwind area about 70 km to the northeast in Cool, CA. A combination of measurements and model data during CARES (Fast et al., 2012) shows that emissions from the San Francisco Bay area transported by intrusions of marine air contributed a large fraction of the carbon monoxide in the vicinity of Sacramento. The study also showed that mountain venting processes contributed to aged pollutants aloft in the valley atmosphere, which can then be entrained into the growing boundary layer the following day. Overall, the CARES campaign helped in improving the current scientific understanding of the interaction between urban emissions and downwind biogenic sources in the Sacramento region.

BEARPEX was conducted at the University of California's Blodgett Forest Research Station during June-July 2007 and September-October 2009. Blodgett Forest is located 65 miles northeast of Sacramento. The project was designed to study chemistry downwind of urban areas where there is high VOC reactivity (due to biogenic emissions sources) and low  $\text{NO}_x$ , to understand the full oxidation sequence and subsequent fate of biogenic VOC and the processes leading to formation and removal of biogenic secondary organic aerosol (SOA) and the associated chemical and optical properties of SOA. A study by Bouvier-Brown et al., (2009) suggests that reactive and semi-volatile compounds, especially sesquiterpenes, significantly impact the gas- and particle-phase chemistry of the atmosphere at Blodgett Forest. An analysis of absolute PANs mixing ratios by Lafranchi et al. (2009) reveals a missing PANs sink that can be resolved by increasing the peroxy acetyl radicals + RO<sub>2</sub> rate constant by a factor of 3. At the BEARPEX field site, the sum of the individual biogenically derived nitrates account for two-thirds of the organic nitrate, confirming the importance of biogenic nitrates to the  $\text{NO}_y$  budget (Beaver et al., 2012).

The CABERNET field campaign was conducted during June 2011 in California. The objectives were to develop and evaluate new approaches for regional scale measurements of biogenic VOC emissions, quantify the response of biogenic VOC emissions to land cover change, investigate the vertical transport of isoprene and oxidation products, and evaluate biogenic emission models. Isoprene fluxes were measured on board the Center for Interdisciplinary Remotely-Piloted Aircraft Studies (CIRPAS) Twin Otter (<http://www.cirpas.org/twinOtter.html>) using the virtual disjunct eddy covariance method (Karl et al. 2012). Isoprene flux measurements from



CABERNET have formed the basis for evaluating the biogenic emissions inventory used in California’s SIP modeling (Misztal et al., 2016).

Table 2-1. Major Field Studies in Central California and surrounding areas.

<b>Year</b>	<b>Study</b>	<b>Significance</b>
1970	Project Lo-Jet	Identified summertime low-level jet and Fresno eddy
1972	Aerosol Characterization Experiment (ACHEX)	First TSP chemical composition and size distributions
1979-1980	Inhalable Particulate Network	First long-term PM2.5 and PM10 mass and elemental measurements in Bay Area, Five Points
1978	Central California Aerosol and Meteorological Study	Seasonal TSP elemental composition, seasonal transport patterns
1979-1982	Westside Operators	First TSP sulfate and nitrate compositions in western Kern County
1984	Southern SJV Ozone Study	First major characterization of O3 and meteorology in Kern County
1986-1988	California Source Characterization Study	Quantified chemical composition of source emissions
1988-1989	Valley Air Quality Study	First spatially diverse, chemical characterized, annual and 24-hour PM2.5 and PM10
July and August 1990	Sacramento Area Ozone Study	Intensive ozone measurements in the Sacramento Area
Summer 1990	San Joaquin Valley Air Quality Study/Atmospheric Utilities Signatures Predictions and Experiments (SJVAQS/AUSPEX) –	First central California regional study of O3 and PM2.5

	Also known as SARMAP (SJVAQS/AUSPEX Regional Modeling Adaptation Project)	
July – September 1990	Upper Sacramento Valley Transport Study	Measurements to study the transport of pollutants from the lower to upper Sacramento Valley
July and August 1991	California Ozone Deposition Experiment	Measurements of dry deposition velocities of O <sub>3</sub> using the eddy correlation technique made over a cotton field and senescent grass near Fresno
Winter 1995	Integrated Monitoring Study (IMS-95, the CRPAQS Pilot Study)	First sub-regional winter study
December 1999– February 2001	California Regional PM <sub>10</sub> /PM <sub>2.5</sub> Air Quality Study (CRPAQS) and Central California Ozone Study	First year-long, regional-scale effort to measure both O <sub>3</sub> and PM <sub>2.5</sub>
December 1999 to present	Fresno Supersite	First multi-year experiment with advanced monitoring technology
July 2003	NASA high-resolution lidar flights	First high-resolution airborne lidar application in SJV in the summer
February 2007	U.S. EPA Advanced Monitoring Initiative	First high-resolution airborne lidar application in SJV in the winter
August-October 2007; June-July 2009	BEARPEX (Biosphere Effects on Aerosols and Photochemistry Experiment)	Research-grade measurements to study the interaction of the Sacramento urban plume with downwind biogenic emissions
June 2008	ARCTAS - CARB	First measurement of high-time resolution (1-10s) measurements of organics and free radicals in SJV
May-July 2010	CalNex 2010 (Research at the Nexus of Air Quality and Climate Change)	Expansion of ARCTAS-CARB type research-grade measurements to multi-platform and expanded geographical area including the ocean.

June 2010	CARES (Carbonaceous Aerosols and Radiative Effects Study)	Research-grade measurements of trace gases and aerosols within the Sacramento urban plume to investigate SOA formation
May – June 2010	IONS (Intercontinental transport experiment Ozonesonde Network Study)	Daily Ozonesonde measurements from four coastal and two inland sites in California to improve the characterization of western U.S. baseline ozone
June 2011	CABERNET (California Airborne BVOC Emission Research in Natural Ecosystem Transects)	Provided the first ever airborne flux measurements of isoprene in California
January- February 2013	DISCOVER-AQ (Deriving Information of Surface Conditions from Column and Vertically Resolved Observations Relevant to Air Quality)	Research-grade measurements of trace gases and aerosols during two PM <sub>2.5</sub> pollution episodes in the SJV

## 2.2 Description of the Ambient Monitoring Network

The SFNA is located in the northern half of the California's Central Valley, which is a 500-mile long northwest-southeast oriented valley comprising the Sacramento Valley (SV) and the San Joaquin Valley (SJV) air basins (left panel of Figure 2–1). The SFNA is home to more than 2 million residents encompassing an area of 5600 square miles and is geographically located in two different air basins including the southern portion of the Sacramento Valley Air Basin (SVAB) and the northern central portion of the Mountain Counties Air Basin (MCAB) (right panel of Figure 2–1). The SFNA area occupies the southern portion of the Sacramento Valley, extending to the inland side of the California Coastal Range on the westernmost edge, and continues to the border of the Lake Tahoe air basin to the east, encompassing portions of the Sierra Nevada Mountain Range. It extends southward to the Sacramento Delta Region and northward to include the southern portion of Sutter County. In total, the SFNA comprises all of Sacramento and Yolo counties, the eastern portion of Solano County, the southern portion of Sutter County, and the portions of El Dorado and Placer counties that are not part of the Lake Tahoe Air Basin.

Due to its inland location, the climate of the Sacramento region is more extreme than that of most coastal regions, such as the San Francisco Bay Area. The winters are generally cool and wet, while the summers are hot and dry and both seasons can experience periods of high pressure and stagnation which are conducive to pollutant buildup. These climate conditions result in seasonal patterns where ozone levels are highest during the summer, while PM<sub>2.5</sub> concentrations are highest during the winter. The lack of summertime precipitation, coupled with the large extent of forested land surrounding the Central Valley, also creates conditions highly conducive to wildfires during the summer months.




Figure 2-1 Map of California's Central Valley and the geographical location of Sacramento Federal 8-hr Ozone Non-attainment Area (SFNA).

The worst ozone air quality in the SFNA typically occurs during summer months, where the interaction between geography, climate, and a mix of natural (biogenic) and anthropogenic emissions pose significant challenges to air quality progress. A combination of stable wind fields and recirculation patterns generated by daytime upslope and nighttime downslope flows from the mountains located to the west (Coast Range) and east (Sierra Nevada), tend to confine and trap emissions and the pollutants near the surface (See section 2.4 for details on meteorological conditions conducive to ozone production in SFNA). The anthropogenic NO<sub>x</sub> and ROG emissions from the urban Sacramento area and biogenic ROG emissions from the Sierra foothills coupled with the hot and dry summertime weather conditions facilitate rapid ozone production in the region. During ozone episodes within the Sacramento Metro area, the most important transport pattern is toward the northeast and the foothills within the Sacramento area itself. Due to the general daytime flow pattern from west to east, as well as the time needed for photochemical processes to occur, the highest ozone mixing ratios in the Sacramento region generally occur in the afternoon in the downwind, eastern portion of the region, near Folsom.

LaFranchi et al. (2011), and the references therein, characterized the production and evolution of ozone in the Sacramento region as a Lagrangian air parcel that produces peak ozone levels downwind of the urban city center in the eastern portion of the region (e.g., Folsom). Due to a prevailing northeast wind flow in the region (U.S. EPA, 2012), the ozone plume is diluted as it migrates farther away from the urban core and downwind into the Sierra foothills (located to the east/northeast). The transport of ozone precursor emissions from the urban Sacramento area dominates the ozone production in the downwind Sierra foothill area, where ozone levels are heavily dependent upon the proximity to the upwind urban source. When compared to the location of peak ozone production shortly downwind from the urban Sacramento area, the ozone levels are relatively lower in the downwind foothills area due to its farther proximity from the upwind anthropogenic NO<sub>x</sub> sources.

The air quality planning in the SFNA is led by the Sacramento Metro Air Quality Management District ([www.AirQuality.org](http://www.AirQuality.org)). Four other air districts also participate in air planning and management in the area. The Yolo-Solano Air Quality Management District (AQMD) ([www.ysaqmd.org](http://www.ysaqmd.org)) has jurisdiction over Yolo County and the SFNA portion of Solano County. Feather River AQMD ([www.fraqmd.org](http://www.fraqmd.org)) has jurisdiction over Sutter and Yuba counties, including the south Sutter County portion of the SFNA. Placer County Air Pollution Control District (APCD) ([www.placer.ca.gov/apcd](http://www.placer.ca.gov/apcd)) has jurisdiction over Placer County, as El Dorado County AQMD ([www.edcgov.us/AirQualityManagement](http://www.edcgov.us/AirQualityManagement)) does over its county. These five air districts along with the California Air Resources Board (CARB) operate an extensive network of

air quality monitors throughout the region to help improve and protect public health. The data collected from the SFNA regional air monitoring network is used to generate daily air quality forecasts, issue health advisories as needed, support compliance with various ambient air quality standards and serves as the basis for developing long-term attainment strategies and tracking progress toward attainment of health-based air quality standards.

Figure 2-2 shows the spatial distribution of the ozone, NO<sub>x</sub>, and PAMS (Photochemical Assessment Monitoring Stations) monitors in the SFNA (see Table 2-2 for longitude/latitude information for each monitor). There are a total of 17 monitoring sites in the region, which are strategically located to capture pollutants within the densely populated urban Sacramento metropolitan area, as well as downwind regions to measure the transport of the Sacramento urban plume to downwind sites in the foothills of eastern Sacramento, Placer, and El Dorado Counties. Finally, the network is able to provide important information on the spatial variability of pollutants, population exposure, and pollutant transport into the Sacramento area from the west/southwest and thus has been shown to sufficiently capture the highest ozone mixing ratios and the corresponding precursors under various weather conditions. A detailed discussion about the monitoring network and its adequacy can be found in the 2015 Air Monitoring Network and Assessment Plans for Sacramento (<http://www.airquality.org/air-quality-health/air-monitoring>) and other air districts that are part of the SFNA (<https://www.arb.ca.gov/aqd/amnr/amnr.htm>).

For purposes of model evaluation and analysis, the SFNA is divided into three sub regions that are characterized by distinct geography, meteorology, emissions characteristics, transport patterns, and air quality: 1) Western SFNA comprising Yolo, Solano and the southwest portion of Sacramento counties, which lies upwind of the Sacramento urban emission source and is impacted by pollutant transport from the surrounding Bay Area and SJV located on the west/southwest, 2) Central SFNA including the inland urban core, and the metropolitan areas of Sacramento county and the westernmost portion of Placer county, and 3) Eastern SFNA comprising Placer and El Dorado counties in the Sierra Nevada foothills area that is located downwind of urban Sacramento. The geographical extent of the sub-regions in SFNA and their approximate regional boundaries are shown in the bottom panel of Figure 2-2.




Figure 2-2. Map of the Monitoring Sites in the Sacramento Federal 8-hour Ozone Nonattainment Area. The green, blue and magenta circle markers denote the location of NO<sub>x</sub>/NO<sub>y</sub>, ozone and PAMS monitors (top panel). The solid black line denotes the regional boundary of the SFNA, while the grey line denotes the county boundaries. The dashed brown lines (bottom panel) show the approximate regional boundaries of the Western, Central and Eastern sub-regions of SFNA.

Table 2-2. Ozone, NO<sub>x</sub>, and PAMS monitoring sites between 2012 and 2015 in the Sacramento Federal 8-hour ozone Non-attainment Area

Site ID (AQS/ARB)	Sub Region	Site (County, Air Basin)	NO <sub>x</sub>	Ozone	PAMS	Latitude	Longitude
060170020 3196	Eastern SFNA	Cool-Hwy193 (El Dorado, MCAB)		X		38.892	-121.002
060170010 3017		Placerville-Gold Nugget Way (El Dorado, MCAB)		X		38.725	-120.822
060170012 3487		Echo Summit (El Dorado, MCAB)		X		38.812	-120.033
060610003 3789		Auburn - Atwood Rd (Placer, SVAB)		X		38.936	-121.1
060610004 3002		Colfax-City Hall (Placer, MCAB)		X		39.1	-120.954
060612002 3796*	Central SFNA	Lincoln - 1445 1st St <sup>1</sup> (Placer, SVAB)		X		38.886	-121.302
060610006 2956		Roseville- N Sunrise Ave (Placer, SVAB)	X	X		38.746	-121.265
060670012 3187		Folsom-Natoma Street (Sacramento, SVAB)	X	X	X	38.683	-121.164
060670002 2123		North Highlands- Blackfoot Way (Sacramento, SVAB)	X	X		38.712	-121.381
060670006 2731		Sacramento- Del Paso Manor (Sacramento, SVAB)	X	X		38.614	-121.368
060670014 3738		Sacramento-Goldenland Court (Sacramento, SVAB)	X	X	X	38.651	-121.507
060670010 3011		Sacramento - 1309 T Street (Sacramento, SVAB)	X	X	X	38.568	-121.493
060675003 3209		Sloughhouse (Sacramento, SVAB)		X		38.495	-121.211
060670011 2977		Western SFNA	Elk Grove - Bruceville Road (Sacramento, SVAB)	X	X	X	38.303
060953003 3678	Vacaville-Ulatis Drive (Solano, SVAB)			X		38.357	-121.95
061130004 2143	Davis-UCD Campus (Yolo, SVAB)		X	X		38.535	-121.774
061131003 3249	Woodland-Gibson Road (Yolo, SVAB)			X		38.661	-121.731

<sup>1</sup> As the Lincoln site in Placer County became operational in October 2012, the measurements were not available for calculating 8-hr ozone design values in 2012 and 2013. Hence this site was excluded from the current SIP attainment demonstration.



### 2.3 Ozone Trends and Sensitivity to Emissions Reductions

The Sacramento Federal Non-attainment Area (SFNA) is one of the most severely polluted air basins in the U.S., and is designated as a severe ozone nonattainment area for the U.S. EPA 2008 0.075 ppm 8-hour ozone standard. Anthropogenic sources of oxides of nitrogen ( $\text{NO}_x$ ) and reactive organic gases (ROG), along with natural biogenic ROG emissions, are the major precursors that lead to ozone formation in the region. The SFNA's anthropogenic emissions inventory is dominated by emissions from the urbanized areas in Sacramento, Yolo, Solano and Placer counties, while the biogenic ROG emissions in the Sierra foothills and Coast Range are the primary contributors to natural emissions in the region. Since the 1980's, the region's emission control program has substantially reduced emissions of both anthropogenic  $\text{NO}_x$  and ROG throughout the region (<https://www.arb.ca.gov/aqd/almanac/almanac.htm>). As the control program has led to changes in the relative levels of  $\text{NO}_x$  and ROG over time, it has also adapted so as to reduce ozone levels as expeditiously as possible. This adaptation within the control program is necessary because ozone formation responds differently to  $\text{NO}_x$  and ROG controls as the relative level of each pollutant in the atmosphere changes (see Figure 2-3).

Specifically, ozone formation exhibits a nonlinear dependence on  $\text{NO}_x$  and ROG precursors in the atmosphere. In general terms, under ambient conditions of high- $\text{NO}_x$  and low-ROG ( $\text{NO}_x$ -disbenefit region in Figure 2-3), ozone formation tends to exhibit a disbenefit to reductions in  $\text{NO}_x$  emissions (i.e., ozone increases with decreases in  $\text{NO}_x$ ) and a benefit to reductions in ROG emissions (i.e., ozone decreases with decreases in ROG). In contrast, under ambient conditions of low- $\text{NO}_x$  and high-ROG ( $\text{NO}_x$ -limited region in Figure 2-3), ozone formation shows a benefit to reductions in  $\text{NO}_x$  emissions, while changes in ROG emissions result in only minor decreases in ozone. These two distinct "ozone chemical regimes" are illustrated in Figure 2-3 along with a transitional regime that can exhibit characteristics of both the  $\text{NO}_x$ -disbenefit and  $\text{NO}_x$ -limited regimes. Note that Figure 2-3 is shown for illustrative purposes only, and does not represent the actual ozone sensitivity within the SFNA for a given combination of  $\text{NO}_x$  and VOC (ROG) emissions.

During the 1980's in the SFNA, ROG emission controls outpaced  $\text{NO}_x$  controls as the ROG emissions were high relative to  $\text{NO}_x$ . During the 1990's, emission controls slowly shifted to a more balanced approach between ROG and  $\text{NO}_x$ , and by the 2000's  $\text{NO}_x$  reductions began to outpace ROG reductions. For much of the 1980's through the mid-2000's, the SFNA was in a  $\text{NO}_x$ -disbenefit or transitional chemical regime and it's only been within the past decade (mid- to late-2000's) where this region began transitioning to a  $\text{NO}_x$ -limited chemical regime. This transition from a  $\text{NO}_x$ -disbenefit to a  $\text{NO}_x$ -limited chemical regime can be analyzed through the year-to-year variability in biogenic

ROG emissions, which during the summer ozone season can be many times greater than anthropogenic ROG emissions in the SFNA, as well as through the so called "weekend effect" which shows an increase in ozone on the weekend under  $\text{NO}_x$ -disbenefit conditions (and a decrease under  $\text{NO}_x$ -limited conditions).




Figure 2-3. Illustrates a typical ozone isopleth plot, where each line represents ozone mixing ratio, in 10 ppb increments, as a function of initial  $\text{NO}_x$  and VOC (or ROG) mixing ratio (adapted from Seinfeld and Pandis, 1998, Figure 5.15). General chemical regimes for ozone formation are shown as  $\text{NO}_x$ -disbenefit (red circle), transitional (blue circle), and  $\text{NO}_x$ -limited (green circle).

Area-wide summer emission trends from 2000 to 2014 for the SFNA are shown in Figure 2-4 (top) for anthropogenic  $\text{NO}_x$  and ROG, as well as biogenic ROG (biogenic

trends are for 2001 to 2014). Figure 2-4 clearly shows large decreases in both anthropogenic NO<sub>x</sub> (from 184 tpd to 91 tpd) and ROG (from 173 tpd to 101 tpd) emissions from 2000 to 2012. Over the same time period, biogenic ROG emissions exhibited large year-to-year variability, ranging from ~666 tpd in 2005 to ~1027 tpd and ~950 tpd in 2006 and 2010, respectively. Even at its lowest levels, biogenic ROG is estimated to be five times as high as the anthropogenic ROG inventory (in 2005) and upwards of eight times as high during peak biogenic years.

Over the same 2000 to 2014 time period, the ozone design value and days above the ozone standard (exceedance days) within the SFNA declined steadily (Figure 2-4 middle and bottom, respectively), but also exhibited a fair amount of variability due to year-to-year variability in meteorology and the associated changes in biogenic emissions. Overall, the area-wide design values declined by ~20 ppb from 107 ppb in 2000 to 85 ppb in 2014. However, these DVs are still substantially higher than the 2008 8-hour ozone standard of 75 ppb.

Since the area-wide DV is focused on the highest ozone values and the location of these peaks can change from year-to-year, the exceedance days, a measure of overall air quality and the frequency of ozone exposure, may be a better metric for evaluating changes in ozone chemistry when viewed in the context of changing biogenic ROG emissions. Exceedance days in the SFNA have substantially decreased over time from 61 in 2000 to 29 in 2014 (~52% lower with respect to 2000) indicating significant improvements in ozone air quality across the entire region. The decline in weekend exceedance days was slightly higher (56% decrease from 16 to 7) than the corresponding decline in weekday exceedance days (~51% decrease from 45 to 22) between the years 2000 and 2014.

Comparing the year-to-year variability in exceedance days to similar variability in the biogenic ROG emissions, shows that from 2001-2007 the two were strongly correlated (i.e., when biogenic ROG emissions increased, so did the number of exceedance days). This is consistent with the SFNA region being primarily in a NO<sub>x</sub>-disbenefit regime, where increases in ROG emissions result in enhanced ozone formation. From 2008 onwards, this correlation no longer exists and the two are actually anti-correlated for all years except 2009. Although other factors beyond chemistry, such as meteorology, play a large role in the year-to-year variability in ozone, this is suggestive of a shift from a NO<sub>x</sub>-disbenefit regime to a transitional or NO<sub>x</sub>-limited regime around the 2008 timeframe.

### Sacramento Federal Ozone Non-attainment Area Trend in 8-hr O<sub>3</sub> between 2000 and 2014




Figure 2-4. Trends in SFNA emissions (top), 8-hour ozone design value (middle), and number of days above the 8-hour ozone standard between 2000 and 2014.

Investigating the “weekend effect” and how it has changed over time is also a useful metric for evaluating the ozone chemistry regime in the SFNA. The weekend effect is a

well-known phenomenon in some major urbanized areas where ozone is observed to be lower on weekends than on weekdays. Although there are contributing factors, such as meteorology and activity patterns for various emissions sources, the general consensus is that reduced vehicle traffic (primarily diesel trucks) on the weekend results in lower  $\text{NO}_x$  emissions, while ROG emissions remain relatively unchanged. The corresponding change in ozone is an indication of the chemical regime (e.g., an increase in ozone suggests a  $\text{NO}_x$  disbenefit regime; Heuss et al., 2003). The excess  $\text{NO}_x$  in this regime not only titrates the  $\text{O}_3$  but also mutes the VOC reactivity by using peroxy radicals to terminate  $\text{NO}_2$  as  $\text{NO}_3$  radicals and subsequently  $\text{HNO}_3$ . The reduction of  $\text{NO}_x$  during the weekend would lessen the titration and increase the VOC reactivity, which in turn would lead to increased ozone levels. A lack of a weekend effect (i.e., no pronounced high  $\text{O}_3$  occurrences during weekends) suggests that the region is in a transition regime, while a reverse weekend effect (i.e., lower ozone during weekends) would suggest that the region is in a  $\text{NO}_x$ -limited chemical regime.

Murphy et al., (2007) showed that the weekend effect for ozone in the Sacramento area is strongly influenced by the region's proximity to the  $\text{NO}_x$  emission sources. Hence the trend in day-of-week dependence in the SFNA was analyzed on a sub-regional basis separately for the western (i.e. region upwind of Sacramento), central (i.e. urban Sacramento area) and eastern (i.e. Sierra foothills area downwind to the east of Sacramento) sub-regions (Figure 2-2) using observations between 2000 and 2014 (Figure 2-5). The three-panel scatter plot shown in Figure 2-5 compares the average site-specific weekday (Wednesday and Thursday) and weekend (Sunday) observed summertime (June through September) maximum daily average (MDA) 8-hr ozone value by year (2000 to 2014), separated into three sub-regions: Western SFNA (top), Central SFNA (middle), and Eastern SFNA (bottom). Different definitions of weekday and weekend days were also investigated and did not show appreciable differences from the Wednesday/Thursday and Sunday definitions.

From Figure 2-5 it can be seen that ozone levels are highest in the eastern and central regions of the SFNA, consistent with their location downwind of and within the urban Sacramento emissions source. The lowest ozone levels are seen in the western SFNA region, which is located upwind of the urban Sacramento emissions source. A key observation in Figure 2-5 is that the summertime average weekday and weekend ozone levels have steadily declined between 2000 and 2014, consistent with the decline in the area-wide DV and exceedance days shown in Figure 2-4.




Figure 2-5. Site-specific average weekday and weekend maximum daily average 8-hour ozone for each year from 2000 to 2014 for the Western (top), Central (middle), and Eastern (bottom) sub-regions of SFNA. Points falling below the 1:1 dashed line represent a NO<sub>x</sub>-disbenefit regime, those on the 1:1 dashed line represent a transitional regime, and those above the 1:1 dashed line represent a NO<sub>x</sub>-limited regime. MDA denotes maximum daily average.

Along with the declining ozone, there is a shift in the weekday and weekend ozone trends between 2000 and 2014. In the early 2000's, the central region of the SFNA exhibited roughly the same number of sites with weekend ozone greater than weekday ozone as sites with weekday ozone greater than weekend ozone, which suggests that the regions may have been in the transitional chemical regime for ozone formation. By the mid-2000's, the majority of the sites were showing weekday ozone greater than weekend ozone, which is consistent with a shift into complete NO<sub>x</sub>-limited chemistry. By 2014, however, some of the sites had shifted back towards a more equal distribution between weekday and weekend ozone. This shift though, may be explained by the relatively low level of biogenic emissions in 2014, which could cause a shift from a NO<sub>x</sub>-limited environment to a more transitional chemistry environment (e.g., Figure 2-3).

The Western SFNA region clearly experienced a greater NO<sub>x</sub>-disbenefit in the early 2000's and then moved into a transitional chemical regime in the mid-2000's and transitioned into the NO<sub>x</sub>-limited regime around the 2010/2011 timeframe. There is a shift back towards a more equal distribution between weekday and weekend ozone in 2014, similar to the Central sub-region. However, this shift occurs at low ozone levels (below 60 ppb) that are well below the 75 ppb ozone standard.

In contrast to the central and western portions (described above), the eastern portion of SFNA has been in a NO<sub>x</sub> limited regime all along, as seen from the greater weekday ozone when compared to weekend ozone. This region is in close proximity to biogenic ROG emissions sources and farther away from the anthropogenic NO<sub>x</sub> sources, such that ROG mixing ratios are relatively high compared to NO<sub>x</sub>, resulting in a NO<sub>x</sub>-limited regime. The shift towards more equal weekday/weekend ozone levels during the 2011-2014 timeframe, presumably due to the low level of biogenic emissions (Figure 2-4), highlights the important contribution of biogenic ROG emissions to ozone formation in this region.

These findings are consistent with an independent analysis by UC Berkeley researchers on the observed response of ozone between 2001 and 2007 in the Sacramento region to NO<sub>x</sub> emission reductions (LaFranchi et al. 2011). This study concluded that NO<sub>x</sub> emission reductions had been effective at reducing ozone levels at all points in the Sacramento urban plume, and by 2007 had successfully transitioned the region to a NO<sub>x</sub>-limited chemistry regime, except within the Sacramento Metropolitan Area urban core. The UC Berkeley study further predicted that the future cumulative NO<sub>x</sub> controls over time will likely transition the entire SFNA (including the urban core) to a NO<sub>x</sub> limited regime, which will make NO<sub>x</sub> emission controls extremely effective in reducing the Sacramento region's ozone levels.

## 2.4 Meteorological Conditions Leading to Ozone Exceedances

The SFNA is located in the highly complex terrain region of California's Central Valley (See Figure 2-1). Elevations in the Central Valley extend from a few feet to almost 500 feet above sea level. This long valley is surrounded by the Coastal Mountain Range on the west, the Cascade Range to the northeast, the Sierra Nevada Mountains on the east, and the Tehachapi Mountains to the south. The Coastal Range is actually a series of north/south mountain ranges that extend 800 miles from the northwest corner of Del Norte County south to the Mexican border. The San Francisco Bay Area divides the Coastal Mountain Range into northern and southern ranges. The Coastal Mountains generally form a barrier between the Pacific Ocean and the Central Valley, with occasional breaks created by low elevation passes and the small gap between the northern and southern ranges in the San Francisco Bay area known as the Carquinez Strait. Elevations in the Coastal Range generally vary between 2,000 and 4,000 feet, but can reach heights above 7,000 feet. In contrast, elevations in the Cascade Range and Sierra Mountains in northern California are typically above 5,000 feet and can exceed 10,000 feet.




Figure 2-6 Conceptual low-level wind patterns in Central California during the day (left panel) and night (right panel) for typical ozone episode conditions (adapted from Bao et al., 2008).



Weather conditions during much of the summer ozone season are dominated by an area of high pressure, known as the East Pacific Ridge, which creates a broad region of warm, descending air over Central California. Studies have shown that the strength and positioning of this ridge has a strong influence on the prevailing weather conditions and summertime ozone levels in Central California (Lehrman et al., 2004; Pun et al., 2008). Synoptic forcing under the East Pacific Ridge is typically weak, with wind flows above the planetary boundary layer from the northwest, resulting in wind flows in Central California that are primarily thermally driven and strongly influenced by orographic effects (Zhong et al., 2004). Thermal gradients between the eastern Pacific Ocean and inland in the Valley result in a strong daytime sea breeze which follows the terrain and can extend well inland through the Carquinez Strait and to a lesser extent the Altamont, Pacheco, and Cholame Passes. When meteorological conditions are favorable, polluted air masses from the Bay Area travel through the Carquinez Strait and bifurcate over the Delta region, with one branch flowing to the northeast into the southern Sacramento Valley and the other branch flowing southeast into the northern San Joaquin Valley (Figure 2-6).

At night, the sea breeze gradually weakens and can even reverse in some cases, but up-valley flow off of the Delta usually persists. Nighttime surface wind flow in the Central Valley is dominated by downslope flows, known as nocturnal drainage, off of the mountain ranges on all sides (Figure 2-6) and when combined with the continued up-valley flows from the Delta, result in low-level eddies such as the Schultz eddy in the southern Sacramento Valley and the Fresno eddy in the SJV (Lehrman et al., 2004). The dynamical conditions favorable for the formation of both the Fresno and Shultz eddies are investigated and discussed by Lin and Jao (1995).

Clustering and classification techniques have been utilized on both observed meteorology (Lehrman et al., 2001; Blanchard et al., 2008; Beaver and Palazoglu, 2009) and observed and modeled ozone (Fujita et al., 1999; Jin et al., 2011) in the Valley and the surrounding region to better understand the relationship between meteorology and elevated ozone. These various studies reveal that the position and strength of the Pacific High has a dominant influence on ozone levels throughout the Central Valley, along with the height of the marine inversion and strength of the low-level on-shore flow. Synoptic flows that weaken or break down the Pacific High result in lower ozone throughout the Central Valley, while a strong sea breeze with a deep marine boundary layer results in lower ozone levels within the Bay Area, but also an enhanced transport of polluted air masses into the Delta region. Under such conditions, elevated ozone can occur in the Sacramento and San Joaquin Valleys if the synoptic forcing is sufficiently weak so that vertical mixing is reduced and recirculation is enhanced. The highest ozone levels in the Valley occur as the thermal gradient

between off-shore and inland weakens and the high pressure system strengthens, resulting in reduced transport of polluted air masses from the Bay Area inland to the Delta, which is accompanied by a rise in temperatures inland. As the sea breeze weakens even further, conditions stagnate within the Valley and ozone levels peak and continue to remain elevated until a synoptic system moves through the area and breaks down the Pacific High.

From an air quality perspective, the Schultz eddy plays a critical role in determining the ozone levels in the Sacramento Valley. The Schultz eddy is the local counterclockwise eddy often formed to the north or northwest of Sacramento due to interaction between the northward marine up-valley inflow and the nocturnal down-valley flow. The typical air flow in the Sacramento Metro area counties is most frequently from the south-southwest, consistent with the incoming marine flow through the Carquinez Strait into the region and orientation of the river valleys extending northeast of Sacramento into the foothills and ranges of the Sierra Nevada mountain range. Instead of allowing the prevailing wind patterns to move north carrying the pollutants out of the region, the Schultz eddy causes the wind pattern and pollutants to circle back in a southeasterly flow, which serves as a mechanism to recirculate and trap air within the region thereby exacerbating the pollution levels in the area and increasing the likelihood of violating the federal and state air quality standards. The Schultz eddy also contributes to the formation of a low-level southerly jet between 500 and 1,000 ft above the surface that is capable of speeds in excess of 35 miles per hour. This jet serves as an important nighttime pollutant transport mechanism transporting air pollutants over large distances thereby impacting the air quality in the Sacramento Valley. The conditions that promote the formation of this jet within the Sacramento Valley may also limit ventilation of the region, resulting in a buildup of pollution over multiple days. The Schultz eddy normally dissipates around noon when the delta sea breeze arrives.

In summary, typical synoptic (large) and local scale weather features associated with 8-hour ozone exceedances in the SFNA generally consist of:

- Broad, upper-level high pressure over the eastern Pacific and western U.S.
- Clear skies
- Sinking motion over the region, which limits vertical mixing through the creation of a subsidence inversion
- Weak winds in most levels of the atmosphere
- Very warm to hot temperatures at the surface and aloft
- Peak warming across the western side or central portion of the Sacramento Valley, which limits the strength of the delta breeze

Synoptic and local scale weather features typically not conducive to 8-hour ozone exceedances include:

- Upper-level low pressure off the Northern California coast (onshore winds) or centered over the four-corner states of Utah, Colorado, Arizona, and New Mexico (northerly winds)
- Rising motion and moderate temperatures aloft, which allow for vertical mixing during peak afternoon heating and pollutant dispersion
- Temperatures rapidly increasing from one day to the next or extremely hot temperatures, both of which lead to a breaking of the temperature inversion
- Moderate to strong northerly winds, even if associated with hot temperatures and clear skies
- Persistent delta breeze on consecutive days with periods of strong onshore winds, which limit exceedances to the eastern-side of the region, namely the foothills, or prevent them entirely

It should be noted that nearly every summer sees both patterns occur, but the key difference is the persistence of one of the patterns, in general, over several weeks and having the pattern align with the peak ozone forming months of July, August, and early September.

## REFERENCES

Bao, J.W., Michelson, S.A., Persson, P.O.G., Djalalova, I.V., and Wilczak, J.M., 2008, Observed and WRF-simulated low-level winds in a high-ozone episode during the Central California Ozone Study, *Journal of Applied Meteorology and Climatology*, 47(9), 2372-2394.

Beaver, M.R. et al., 2012 Importance of biogenic precursors to the budget of organic nitrates: observations of multifunctional organic nitrates by CIMS and TD-LIF during BEARPEX 2009, *Atmos. Chem. Phys.*, 12, 5773-5785.

Beaver, S.; Palazoglu, A. 2009. Influence of synoptic and mesoscale meteorology on ozone pollution potential for San Joaquin Valley of California. *Atmos. Environ.* 43: 1779–1788.

Blanchard, C.L.; Tanenbaum, S.; Fujita, E.M.; Campbell, D.; Wilkinson, J. August 2008. Understanding Relationships between Changes in Ambient Ozone and Precursor Concentrations and Changes in VOC and NO<sub>x</sub> Emissions from 1990 to 2004 in Central California

Bouvier-Brown, N. C., Goldstein, A. H., Gilman, J. B., Kuster, W. C., and de Gouw, J. A.: In-situ ambient quantification of monoterpenes, sesquiterpenes, and related oxygenated compounds during BEARPEX 2007: implications for gas- and particle-phase chemistry, *Atmos. Chem. Phys.*, 9, 5505-5518, doi:10.5194/acp-9-5505-2009, 2009.

Cai C. et al., 2016 Simulating reactive nitrogen, carbon monoxide, and ozone in California during ARCTAS-CARB 2008 with high wildfire activity, *Atmos. Environ.* 128, 28-44.

Fast JD, WI Gustafson, Jr, LK Berg, WJ Shaw, MS Pekour, MKB Shrivastava, JC Barnard, R Ferrare, CA Hostetler, J Hair, MH Erickson, T Jobson, B Flowers, MK Dubey, PhD, SR Springston, BR Pirce, L Dolislager, JR Pederson, and RA Zaveri. 2012. "Transport and Mixing Patterns over Central California during the Carbonaceous Aerosol and Radiative Effects Study (CARES)." *Atmospheric Chemistry and Physics* 12(4):1759-1783. doi:10.5194/acp-12-1759-2012

Federal Register, 1997, Approval and Promulgation of Implementation Plans; California Ozone, Final Rule, January 8<sup>th</sup>, 1150-1187.

Federal Register, 2007, Treatment of Data Influenced by Exceptional Events; Final Rule", Final Rule, March 22<sup>nd</sup>, 13560-13581.

Federal Register, 2012, Determination of Attainment of the 1-Hour Ozone National Ambient Air Quality Standards in the Sacramento Metro Nonattainment Area in California, Final Rule, October 18<sup>th</sup>, 64036-64039.

Federal Register, 2014, Approval and Promulgation of Implementation Plans: State of California; Sacramento Metro Area; Attainment Plan for 1997 8-Hour Ozone Standard, Proposed Rule, October 15<sup>th</sup>, 61799-61822.

Fujita, E., D. Campbell, R. Keisler, J. Brown, S. Tanrikulu, and A. J. Ranzieri, 2001, Central California Ozone Study (CCOS)-Final report, volume III:Summary of field operations, Technical Report, California Air Resources Board, Sacramento.

Fujita, E., Keislar, R., Stockwell, W., Moosuller, H., DuBois, D., Koracin, D. and Zielinska, B. Central California Ozone Study-Volume I, Field Study Plan.Division of Atmospheric Science Desert Research Institue, 2215 Raggio Parkway, Reno, NV. 1999

Gentner, D.R. et al., 2014a, Emissions of organic carbon and methane from petroleum and dairy operations in California's San Joaquin Valley *Atmos. Chem. Phys.*, 14, 4955-4978.

Gentner, D.R., Ormeño, Fares, E.S., Ford, T.B., Weber, R., Park, J.-H., Brioude, J., Angevine, W.M., Karlik, J.F., and Goldstein, A.H., 2014b, Emissions of terpenoids, benzenoids, and other biogenic gas-phase organic compounds from agricultural crops and their potential implications for air quality *Atmos. Chem. Phys.*, 14, 5393–5413.

Heuss, J.M., Kahlbaum, D.F., and Wolff, G.T., 2003. Weekday/weekend ozone differences: What can we learn from them? *Journal of the Air & Waste Management Association* 53(7), 772-788.

Huang, M., Carmichael, G.R., Adhikary, B., Spak, S.N., Kulkarni, S., Cheng, Y.F., Wei, C., Tang, Y., Parrish, D.D., Oltmans, S.J., D'Allura, A., Kaduwela, A., Cai, C., Weinheimer, A.J., Wong, M., Pierce, R.B., Al-Saadi, J.A. Streets, D.G., and Zhang, Q., 2010, Impacts of transported background ozone on California air quality during the ARCTAS-CARB period - a multi-scale modeling study, *Atmospheric Chemistry and Physics*, 10(14), 6947-6968.

Jin, L., Harley, R.A., and Brown, N.J., 2011, Ozone pollution regimes modeled for a summer season in California's San Joaquin Valley: A cluster analysis, *Atmospheric Environment*, 45, 4707-4719.

LaFranchi, B. W., Goldstein, A. H., and Cohen, R. C.: Observations of the temperature dependent response of ozone to NO<sub>x</sub> reductions in the Sacramento, CA urban plume, *Atmos. Chem. Phys.*, 11, 6945-6960, doi:10.5194/acp-11-6945-2011, 2011

LaFranchi, B.W. et al., 2009, Closing the peroxy acetyl nitrate budget: observations of acyl peroxy nitrates (PAN, PPN, and MPAN) during BEARPEX 2007, *Atmos. Chem. Phys.*, 9, 7623–7641.

Lehrman, D., B.Knuth and D.Fairley, 2001. Characterization of the 2000 Measurement Period, Interim Report, Contract No. 01-2CCOS, Technical and Business Systems, Inc., November.

Lehrman, D., Bush, D., Knuth, B., Fairley, D., Blanchard, C., 2004. Characterization of the CCOS 2000 measurement period. Final report, California Air Resources Board, Sacramento, CA, available at <http://www.arb.ca.gov/airways/ccos/ccos.htm> (item II-8).

Lin, Y.L., and Jao, I.C., 1995, A Numerical Study of Flow Circulations in the Central Valley of California and Formation Mechanisms of the Fresno Eddy, Monthly Weather Review, 123(11), 3227-3239.

Misztal, P. K., Avise, J. C., Karl, T., Scott, K., Jonsson, H. H., Guenther, A. B., and Goldstein, A. H.: Evaluation of regional isoprene emission factors and modeled fluxes in California, Atmos. Chem. Phys., 16, 9611-9628, doi:10.5194/acp-16-9611-2016, 2016.

Murphy, J. G., Day, D. A., Cleary, P. A., Wooldridge, P. J., Millet, D. B., Goldstein, A. H., and Cohen, R. C.: The weekend effect within and downwind of Sacramento – Part 1: Observations of ozone, nitrogen oxides, and VOC reactivity, Atmos. Chem. Phys., 7, 5327-5339, doi:10.5194/acp-7-5327-2007, 2007

NOAA (2014), Synthesis of Policy Relevant Findings from the CalNex 2010 Field Study, Final report to the California Air Resources Board, available at <http://www.esrl.noaa.gov/csd/projects/calnex/synthesisreport.pdf>

Pun, B.K., J.F. Louis and C. Seigneur, 2008. A conceptual model of ozone formation in the San Joaquin Valley. Doc. No. CP049-1-98. Atmospheric and Environmental Research Inc., San Ramon, CA, 15 December.

Pusede, S. E., and R. C. Cohen, 2012, On the observed response of ozone to NO<sub>x</sub> and VOC reactivity reductions in San Joaquin Valley California 1995–present, Atmos. Chem. Phys., 12, 8323–8339.

Pusede, S. E., Gentner, D. R., Wooldridge, P. J., Browne, E. C., Rollins, A. W., Min, K.-E., Russell, A. R., Thomas, J., Zhang, L., Brune, W. H., Henry, S. B., DiGangi, J. P., Keutsch, F. N., Harrold, S. A., Thornton, J. A., Beaver, M. R., St. Clair, J. M., Wennberg, P. O., Sanders, J., Ren, X., VandenBoer, T. C., Markovic, M. Z., Guha, A., Weber, R., Goldstein, A. H., and Cohen, R. C.: On the temperature dependence of organic reactivity, nitrogen oxides, ozone production, and the impact of emission controls in San Joaquin Valley, California, Atmos. Chem. Phys., 14, 3373-3395, doi:10.5194/acp-14-3373-2014, 2014.

Roberts, P.T., C.G. Lindsey and E.M. Prins (1990). "1990 Field Operations Plan: Sacramento Area Ozone Study." Sonoma Technology Inc. Report #STI-90042-1015 prepared for Systems Applications, Inc. and the Sacramento Area Council of Governments, Santa Rosa, CA.

Seinfeld J. H. and Pandis S. N. (1998) Atmospheric Chemistry and Physics: From Air Pollution to Climate Change, 1st edition, J. Wiley, New York.

Solomon, P.A. and Magliano, K.L., 1998, The 1995-Integrated Monitoring Study (IMS95) of the California Regional PM10/PM2.5 air quality study (CRPAQS): Study overview, Atmospheric Environment, 33(29), 4747-4756.

U.S. EPA, (2012) 2008 Ground-Level Ozone Standards - Final Designations  
[https://www3.epa.gov/region9/air/ozone/pdf/R9\\_CA\\_Sacramento\\_FINAL.pdf](https://www3.epa.gov/region9/air/ozone/pdf/R9_CA_Sacramento_FINAL.pdf)

Zhong, Shiyuan, C. David Whiteman, Xindi Bian, 2004: Diurnal Evolution of Three-Dimensional Wind and Temperature Structure in California's Central Valley. J. Appl. Meteor., 43, 1679–1699.

## Appendix B-3

## Modeling Protocol

### **Document Title:**

Photochemical Modeling Protocol – Photochemical Modeling for the 8-Hour Ozone and Annual/24-hour PM<sub>2.5</sub> State Implementation Plans

### **Document Description:**

This document provides details and formalizes the procedures for conducting the photochemical modeling that forms the basis of the attainment demonstration. The protocol is intended to communicate up front how the model attainment test will be performed. In addition, the protocol discusses analyses that help corroborate the findings of the model attainment test.



# PHOTOCHEMICAL MODELING PROTOCOL

## Photochemical Modeling for the 8-Hour Ozone and Annual/24-hour PM<sub>2.5</sub> State Implementation Plans

**Prepared by**  
California Air Resources Board

**Prepared for**  
United States Environmental Protection Agency Region IX

January 27, 2017

**TABLE OF CONTENTS**

1. INTRODUCTION..... 9

    1.1 Modeling roles for the current SIP ..... 9

    1.2 Stakeholder participation..... 9

    1.3 Involvement of external scientific/technical experts and their input on the  
photochemical modeling ..... 10

    1.4 Schedule for completion of the Plan..... 11

2. DESCRIPTION OF THE CONCEPTUAL MODEL FOR THE NONATTAINMENT  
AREA ..... 11

3. SELECTION OF MODELING PERIODS..... 11

    3.1 Reference Year Selection and Justification..... 11

    3.2 Future Year Selection and Justification ..... 12

    3.3 Justification for Seasonal/Annual Modeling Rather than Episodic Modeling .... 13

4. DEVELOPMENT OF EMISSION INVENTORIES ..... 14

5. MODELS AND INPUTS ..... 14

    5.1 Meteorological Model ..... 14

        5.1.1 Meteorological Modeling Domain ..... 15

    5.2 Photochemical Model..... 18

        5.2.1 Photochemical Modeling Domain ..... 20

        5.2.2 CMAQ Model Options..... 22

        5.2.3 Photochemical Mechanism..... 22

        5.2.4 Aerosol Module..... 23

        5.2.5 CMAQ Initial and Boundary Conditions (IC/BC) and Spin-Up period..... 24

    5.3 Quality Assurance of Model Inputs..... 26

6. METEOROLOGICAL MODEL PERFORMANCE ..... 27

    6.1 Ambient Data Base and Quality of Data..... 27

    6.2 Statistical Evaluation ..... 27

    6.3 Phenomenological Evaluation ..... 29

7. PHOTOCHEMICAL MODEL PERFORMANCE ..... 29

    7.1 Ambient Data ..... 29

    7.2 Statistical Evaluation ..... 31

---

7.3	Comparison to Previous Modeling Studies .....	33
7.4	Diagnostic Evaluation.....	33
8.	ATTAINMENT DEMONSTRATION.....	34
8.1	Base Year Design Values .....	34
8.2	Base, Reference, and Future Year Simulations .....	35
8.3	Relative Response Factors .....	36
8.3.1	8-hour Ozone RRF .....	36
8.3.2	Annual and 24-hour PM <sub>2.5</sub> RRF .....	37
8.4	Future Year Design Value Calculation .....	38
8.4.1	8-hour Ozone.....	38
8.4.2	Annual and 24-hour PM <sub>2.5</sub> .....	38
8.5	Unmonitored Area Analysis.....	45
8.5.1	8-hour Ozone.....	45
8.5.2	Annual PM <sub>2.5</sub> .....	46
8.5.3	24-hour PM <sub>2.5</sub> .....	47
8.6	Banded Relative Response Factors for Ozone .....	48
9.	PROCEDURAL REQUIREMENTS .....	49
9.1	How Modeling and other Analyses will be Archived, Documented, and Disseminated.....	49
9.2	Specific Deliverables to U.S. EPA.....	49
	REFERENCES.....	50

**LIST OF FIGURES**

Figure 5-1. The three nested grids for the WRF model (D01 36km; D02 12km; and D03 4km). ..... 16

Figure 5-2. CMAQ modeling domains used in this SIP modeling platform. The outer domain (dashed black line) represents the extent of the California statewide domain (shown here with a 4 km horizontal resolution, but utilized in this modeling platform with a 12 km horizontal resolution). Nested higher resolution 4 km modeling domains are highlighted in green and red for the Northern/Central California and Southern California, respectively. The smaller SJV PM<sub>2.5</sub> 4 km domain (colored in blue) is nested within the Northern California 4 km domain..... 21

Figure 5-3. Comparison of MOZART (red) simulated CO (left), ozone (center), and PAN (right) to observations (black) along the DC-8 flight track. Shown are mean (filled symbol), median (open symbols), 10th and 90th percentiles (bars) and extremes (lines). The number of data points per 1-km wide altitude bin is shown next to the graphs. Adapted from Figure 2 in Pfister et al. (2011)..... 25

Figure 8-1. Example showing how the location of the MDA8 ozone for the top ten days in the reference and future years are chosen..... 37

**LIST OF TABLES**

Table 3-1. Future attainment year by non-attainment region and NAAQS. 0.08 ppm and 0.075 ppm refer to the 1997 and 2008 8-hour ozone standards, respectively. 15 ug/m<sup>3</sup> and 12 ug/m<sup>3</sup> refer to the 1997 and 2012 annual PM<sub>2.5</sub> standards, respectively. 35 ug/m<sup>3</sup> refers to the 2006 24-hour PM<sub>2.5</sub> standard, and 1-hr ozone refers to the revoked 1979 0.12 ppm 1-hour ozone standard. .... 13

Table 5-1. WRF vertical layer structure. .... 17

Table 5-2. WRF Physics Options. .... 18

Table 5-3. CMAQ v5.0.2 configuration and settings. .... 22

Table 7-1. Monitored species used in evaluating model performance. .... 30

Table 8-1. Illustrates the data from each year that are utilized in the Design Value calculation for that year (DV Year), and the yearly weighting of data for the weighted Design Value calculation (or DV<sub>R</sub>). “obs” refers to the observed metric (8-hr O<sub>3</sub>, 24-hour PM<sub>2.5</sub>, or annual average PM<sub>2.5</sub>). .... 35

## ACRONYMS

ARB – Air Resources Board

ARCTAS-CARB – California portion of the Arctic Research of the Composition of the Troposphere from Aircraft and Satellites conducted in 2008

BCs – Boundary Conditions

CalNex – Research at the Nexus of Air Quality and Climate Change conducted in 2010

CCOS - Central California Ozone Study

CMAQ Model – Community Multi-scale Air Quality Model

CIT – California Institute of Technology

CRPAQS – California Regional PM<sub>10</sub>/PM<sub>2.5</sub> Air Quality Study

DISCOVER-AQ - Deriving Information on Surface Conditions from Column and Vertically Resolved Observations Relevant to Air Quality

DV – Design Value

FDDA – Four-Dimensional Data Assimilation

FEM – Federal Equivalence Monitors

FRM – Federal Reference Monitors

HNO<sub>3</sub> – Nitric Acid

ICs – Initial Conditions

IMPROVE – Interagency Monitoring of Protected Visual Environments

IMS-95 – Integrated Monitoring Study of 1995

LIDAR – Light Detection And Ranging

MDA – Maximum Daily Average

MM5 – Mesoscale Meteorological Model Version 5

MOZART – Model for Ozone and Related chemical Tracers

NARR - North American Regional Reanalysis

NCAR – National Center for Atmospheric Research

NCEP – National Centers for Environmental Prediction

NH<sub>3</sub> – Ammonia

NOAA - National Oceanic and Atmospheric Administration

NO<sub>x</sub> – Oxides of nitrogen

OC – Organic Carbon

OFP - Ozone Forming Potential

PAMS – Photochemical Assessment Monitoring Stations

PAN – Peroxy Acetyl Nitrate

PM<sub>2.5</sub> – Particulate Matter with aerodynamic diameter less than 2.5 micrometers

PM<sub>10</sub> – Particulate Matter with aerodynamic diameter less than 10 micrometers

RH – Relative Humidity

ROG – Reactive Organic Gases

RRF – Relative Response Factor

RSAC – Reactivity Scientific Advisory Committee

SANDWICH – Application of the Sulfate, Adjusted Nitrate, Derived Water, Inferred Carbonaceous Material Balance Approach

SAPRC – Statewide Air Pollution Research Center

SARMAP – SJVAQS/AUSPEX Regional Modeling Adaptation Project

SCAQMD – South Coast Air Quality Management District

SIP – State Implementation Plan

SJV – San Joaquin Valley

SJVAB – San Joaquin Valley Air Basin (SJVAB)

SJVUAPCD – San Joaquin Valley Unified Air Pollution Control District

SJVAQS/AUSPEX – San Joaquin Valley Air Quality Study/Atmospheric Utilities Signatures Predictions and Experiments

SLAMS – State and Local Air Monitoring Stations

SMAQMD – Sacramento Metropolitan Air Quality Management District

SMAT – Application of the Speciated Modeled Attainment Test

SOA – Secondary Organic Aerosol

SO<sub>x</sub> – Oxides of Sulfur

STN – Speciated Trend Network

UCD – University of California at Davis

U.S. EPA – United States Environmental Protection Agency

VOC – Volatile Organic Compounds

WRF Model – Weather and Research Forecast Model



## 1. INTRODUCTION

The purpose of this modeling protocol is to detail and formalize the procedures for conducting the photochemical modeling that forms the basis of the attainment demonstration for the 8-hour ozone and annual/24-hour PM<sub>2.5</sub> State Implementation Plans (SIPs) for California. The protocol is intended to communicate up front how the model attainment test will be performed. In addition, this protocol discusses analyses that are intended to help corroborate the findings of the model attainment test.

### 1.1 Modeling roles for the current SIP

The Clean Air Act (Act) establishes the planning requirements for all those areas that routinely exceed the health-based air quality standards. These nonattainment areas must adopt and implement a SIP that demonstrates how they will attain the standards by specified dates. Air quality modeling is an important technical component of the SIP, as it is used in combination with other technical information to project the attainment status of an area and to develop appropriate emission control strategies to achieve attainment.

ARB and local Air Districts will jointly develop the emission inventories, which are an integral part of the modeling. Working closely with the Districts, the ARB will perform the meteorological and air quality modeling. Districts will then develop and adopt their local air quality plan. Upon approval by the ARB, the SIP will be submitted to U.S.EPA for approval.

### 1.2 Stakeholder participation

Public participation constitutes an integral part of the SIP development. It is equally important in all technical aspects of SIP development, including the modeling. As the SIP is developed, the Air Districts and ARB will hold public workshops on the modeling and other SIP elements. Representatives from the private sector, environmental interest groups, academia, and the federal, state, and local public sectors are invited to attend and provide comments. In addition, Draft Plan documents will be available for public review and comment at various stages of plan development and at least 30 days before Plan consideration by the Districts' Governing Boards and subsequently by the ARB Board. These documents will include descriptions of the technical aspects of the SIP. Stakeholders have the choice to provide written and in-person comments at any of the Plan workshops and public Board hearings. The agencies take the comments into consideration when finalizing the Plan.

### 1.3 Involvement of external scientific/technical experts and their input on the photochemical modeling

During the development of the modeling protocol for the 2012 SJV 24-hour PM<sub>2.5</sub> SIP (SJVUAPCD, 2012), ARB and the San Joaquin Valley Air Pollution Control District (SJVAPCD) engaged a group of experts on prognostic meteorological modeling and photochemical/aerosol modeling to help prepare the modeling protocol document.

The structure of the technical expert group was as follows:

Conveners: John DaMassa – ARB  
Samir Sheikh – SJVAPCD  
Members: Scott Bohning – U.S. EPA Region 9  
Ajith Kaduwela – ARB  
James Kelly – U.S. EPA Office of Air Quality Planning and Standards  
Michael Kleeman – University of California at Davis  
Jonathan Pleim – U.S. EPA Office of Research and Development  
Anthony Wexler – University of California at Davis

The technical consultant group provided technical consultations/guidance to the staff at ARB and SJVAPCD during the development of the protocol. Specifically, the group provided technical expertise on the following components of the protocol:

- Selection of the physics and chemistry options for the prognostic meteorological and photochemical air quality models
- Selection of methods to prepare initial and boundary conditions for the air quality model
- Performance evaluations of both prognostic meteorological and photochemical air quality models. This includes statistical, diagnostic, and phenomenological evaluations of simulated results.
- Selection of emissions profiles (size and speciation) for particulate-matter emissions.
- Methods to determine the limiting precursors for PM<sub>2.5</sub> formation.
- Application of the Sulfate, Adjusted Nitrate, Derived Water, Inferred Carbonaceous Material Balance Approach (SANDWICH) with potential modifications.
- Application of the Speciated Modeled Attainment Test (SMAT).
- Selection of methodologies for the determination of PM<sub>2.5</sub> precursor equivalency ratios.
- Preparation of Technical Support Documents.

The current approach to regional air quality modeling has not changed significantly since the 2012 SJV 24-hour PM<sub>2.5</sub> SIP (SJVUAPCD, 2012), so the expertise provided on the above components to the protocol remain highly relevant. In addition, since regional air quality modeling simulates ozone chemistry and PM chemistry/formation simultaneously, there is generally no difference in how the models are configured and simulations conducted for ozone vs. PM. Therefore, development of this modeling protocol will rely heavily on the recommendations made by this group of technical experts, as well as recently published work in peer-review journals related to regional air quality modeling.

#### **1.4 Schedule for completion of the Plan**

Final area designations kick-off the three year SIP development process. For the first two years, efforts center on updates and improvements to the Plan's technical and scientific underpinnings. These include the development of emission inventories, selection of modeling periods, model selection, model input preparation, model performance evaluation and supplemental analyses. During the last year, modeling, further supplemental analyses and control strategy development proceed in an iterative manner and the public participation process gets under way. After thorough review the District Board and subsequently the ARB Board consider the Plan. The Plan is then submitted to U.S. EPA. Table 1-1 in the Appendix corresponding to the appropriate region/standard (e.g., SJV 0.075 ppm 8-hour ozone) summarizes the overall anticipated schedule for Plan completion.

## **2. DESCRIPTION OF THE CONCEPTUAL MODEL FOR THE NONATTAINMENT AREA**

See Section 2 in the Appendix corresponding to the appropriate region/standard (e.g., SJV 0.075 ppm 8-hour ozone).

## **3. SELECTION OF MODELING PERIODS**

### **3.1 Reference Year Selection and Justification**

From an air quality and emissions perspective, ARB and the Districts have selected 2012 as the base year for design value calculation and for the modeled attainment test.

For the SJV, the PM<sub>2.5</sub> model attainment test will utilize 2013 instead of 2012. These baseline values will serve as the anchor point for estimating future year projected design values.

The selection of 2012/13 is based on the following four considerations:

- Most complete and up to date emissions inventory, which reduces the uncertainty associated with future emissions projections.
- Analysis of meteorological adjusted air quality trends to determine recent years with meteorology most conducive to ozone and PM<sub>2.5</sub> formation and buildup.
- Availability of research-grade wintertime field measurements in the Valley, which captured two significant pollution episodes during the DISCOVER-AQ field study (January-February 2013).
- The SJV PM<sub>2.5</sub> design values for year 2013 were some of the highest in recent years, making 2013 a conservative choice for attainment demonstration modeling.

Details and discussion on these analyses can be found in the Weight of Evidence Appendix.

### **3.2 Future Year Selection and Justification**

The future year modeled is determined by the year for which attainment must be demonstrated. Table 3-1 lists the year in which attainment must be demonstrated for the various ozone and PM<sub>2.5</sub> standards and non-attainment regions in California.

Table 3-1. Future attainment year by non-attainment region and NAAQS. 0.08 ppm and 0.075 ppm refer to the 1997 and 2008 8-hour ozone standards, respectively. 15 ug/m<sup>3</sup> and 12 ug/m<sup>3</sup> refer to the 1997 and 2012 annual PM<sub>2.5</sub> standards, respectively. 35 ug/m<sup>3</sup> refers to the 2006 24-hour PM<sub>2.5</sub> standard, and 1-hr ozone refers to the revoked 1979 0.12 ppm 1-hour ozone standard.

Area	Year								
	2031	2026	2025	2024	2023	2021	2020	2019	2017
<b>Southern California Modeling Domain</b>									
South Coast	0.075 ppm	--	--	--	0.08 ppm	12 ug/m <sup>3</sup>	--	--	--
Mojave/Coachella	--	0.075 ppm	--	--	--	--	--	--	0.08 ppm
Imperial County	--	--	--	--	--	12 ug/m <sup>3</sup>	--	--	0.075 ppm
Ventura County	--	--	--	--	--	--	0.075 ppm	--	--
San Diego	--	--	--	--	--	--	--	--	0.075 ppm
<b>Northern California Modeling Domain</b>									
San Joaquin Valley	0.075 ppm	--	<sup>1</sup> 12 ug/m <sup>3</sup>	35 ug/m <sup>3</sup>	--	<sup>2</sup> 12 ug/m <sup>3</sup>	15 ug/m <sup>3</sup>	35 ug/m <sup>3</sup>	1-hr ozone
Sacramento Metropolitan	--	0.075 ppm	--	--	--	--	--	--	--
Portola-Plumas County	--	--	--	--	--	12 ug/m <sup>3</sup>	--	--	--
East Kern	--	--	--	--	--	--	--	--	0.075 ppm
W. Nevada County	--	--	--	--	--	--	--	--	0.075 ppm

<sup>1</sup> Serious classification attainment date

<sup>2</sup> Moderate classification attainment date

### 3.3 Justification for Seasonal/Annual Modeling Rather than Episodic Modeling

In the past, computational constraints restricted the time period modeled for a SIP attainment demonstration to a few episodes (e.g., 2007 SJV 8-hr ozone SIP (SJVUAPCD, 2007), 2007 SC 8-hr ozone SIP (SCAQMD, 2012) and 2009 Sacramento 8-hr ozone SIP (SMAQMD, 2012)). However, as computers have become faster and

large amounts of data storage have become readily accessible, there is no longer a need to restrict modeling periods to only a few episodes. In more recent years, SIP modeling in California has covered the entire ozone or peak  $PM_{2.5}$  seasons (2012 SC 8-hour ozone and 24-hour  $PM_{2.5}$  SIP (SCAQMD, 2012), 2012 SJV 24-hour  $PM_{2.5}$  SIP (SJVUAPCD, 2012) and 2013 SJV 1-hr ozone SIP (SJVUAPCD, 2013) ), or an entire year in the case of annual  $PM_{2.5}$  ( 2008 SJV annual  $PM_{2.5}$  SIP (SJVUAPCD, 2008)) The same is true for other regulatory modeling platforms outside of California (Boylan and Russell, 2006; Morris et al., 2006; Rodriguez et al., 2009; Simon et al., 2012; Tesche et al., 2006; U.S. EPA, 2011a, b).

Recent ozone based studies, which focused on model performance evaluation for regulatory assessment, have recommended the use of modeling results covering the full synoptic cycles and full ozone seasons (Hogrefe et al., 2000; Vizuete et al., 2011). This enables a more complete assessment of ozone response to emission controls under a wide range of meteorological conditions. The same is true for modeling conducted for peak 24-hour  $PM_{2.5}$ . Consistent with the shift to seasonal or annual modeling in most regulatory modeling applications, modeling for the 8-hour ozone standard will cover the entire ozone season (May – September), modeling for the annual 24-hour  $PM_{2.5}$  standard will be conducted for the entire year, and modeling for the 24-hour  $PM_{2.5}$  standard will, at a minimum, cover the months in which peak 24-hour  $PM_{2.5}$  occurs (e.g., October – March in the SJV) and will be conducted annually whenever possible.

## **4. DEVELOPMENT OF EMISSION INVENTORIES**

For a detailed description of the emissions inventory, updates to the inventory, and how it was processed from the planning totals to a gridded inventory for modeling, see the Emissions Inventory Appendix.

## **5. MODELS AND INPUTS**

### **5.1 Meteorological Model**

Meteorological model selection is based on a need to accurately simulate the synoptic and mesoscale meteorological features observed during the selected modeling period. The main difficulties in accomplishing this are California's extremely complex terrain and its diverse climate. It is desirable that atmospheric modeling adequately represent essential meteorological fields such as wind flows, ambient temperature variation, evolution of the boundary layer, and atmospheric moisture content to properly characterize the meteorological component of photochemical modeling.

In the past, the ARB has applied prognostic, diagnostic, and hybrid models to prepare meteorological fields for photochemical modeling. There are various numerical models that are used by the scientific community to study the meteorological characteristics of an air pollution episode. For this SIP modeling platform, the Weather and Research Forecasting (WRF) model (Skaramock et al, 2005) will be used to develop the meteorological fields that drive the photochemical modeling. The U.S. EPA (2014) recommends the use of a well-supported grid-based mesoscale meteorological model for generating meteorological inputs. The WRF model is a community-based mesoscale prediction model, which represents the state-of-the-science and has a large community of model users and developers who frequently update the model as new science becomes available. In recent years, WRF has been applied in California to generate meteorological fields for numerous air quality studies (e.g., Angevine, et al., 2012; Baker et al., 2015; Ensberg et al., 2013; Fast et al., 2014; Hu et al., 2014a, 2014b; Huang et al., 2010; Kelly et al., 2014; Lu et al., 2012; Mahmud et al., 2010), and has been shown to reasonably reproduce the observed meteorology in California.

### 5.1.1 Meteorological Modeling Domain

The WRF meteorological modeling domain consists of three nested grids of 36 km, 12 km and 4 km uniform horizontal grid spacing (illustrated in Figure 5-1). The purpose of the coarse, 36 km grid (D01) is to provide synoptic-scale conditions to all three grids, while the 12 km grid (D02) is used to provide finer resolution data that feeds into the 4 km grid (D03). The D01 grid is centered at 37 °N and 120.5 °W and was chosen so that the inner two grids, D02 and D03, would nest inside of D03 and be sufficiently far away from the boundaries to minimize boundary influences. The D01 grid consists of 90 x 90 grid cells, while the D02 and D03 grids encompass 192 x 192 and 327 x 297 grid cells, respectively, with an origin at -696 km x -576 km (Lambert Conformal projection). WRF will be run for the three nested domains simultaneously with two-way feedback between the parent and the nest grids. The D01 and D02 grids are meant to resolve the larger scale synoptic weather systems, while the D03 grid is intended to resolve the finer details of the atmospheric conditions and will be used to drive the air quality model simulations. All three domains will utilize 30 vertical sigma layers (defined in Table 5-1), as well as the various physics options listed in Table 5-2 for each domain.

The initial and boundary conditions (IC/BCs) for WRF will be prepared based on 3-D North American Regional Reanalysis (NARR) data that are archived at the National Center for Atmospheric Research (NCAR). These data have a 32 km horizontal resolution. Boundary conditions to WRF are updated at 6-hour intervals for the 36 km grid (D01). In addition, surface and upper air observations obtained from NCAR will be used to further refine the analysis data that are used to generate the IC/BCs. Analysis

nudging will be employed in the outer 36km grid (D01) to ensure that the simulated meteorological fields are constrained and do not deviate from the observed meteorology.




Figure 5-1. The three nested grids for the WRF model (D01 36km; D02 12km; and D03 4km).



Table 5-1. WRF vertical layer structure.

Layer Number	Height (m)	Layer Thickness (m)	Layer Number	Height (m)	Layer Thickness (m)
30	16082	1192	14	1859	334
29	14890	1134	13	1525	279
28	13756	1081	12	1246	233
27	12675	1032	11	1013	194
26	11643	996	10	819	162
25	10647	970	9	657	135
24	9677	959	8	522	113
23	8719	961	7	409	94
22	7757	978	6	315	79
21	6779	993	5	236	66
20	5786	967	4	170	55
19	4819	815	3	115	46
18	4004	685	2	69	38
17	3319	575	1	31	31
16	2744	482	0	0	0
15	2262	403			

Note: Shaded layers denote the subset of vertical layers to be used in the CMAQ photochemical model simulations. Further details on the CMAQ model configuration and settings can be found in subsequent sections.

Table 5-2. WRF Physics Options.

Physics Option	Domain		
	D01 (36 km)	D02 (12 km)	D03 (4 km)
Microphysics	WSM 6-class graupel scheme	WSM 6-class graupel scheme	WSM 6-class graupel scheme
Longwave radiation	RRTM	RRTM	RRTM
Shortwave radiation	Dudhia scheme	Dudhia scheme	Dudhia scheme
Surface layer	Revised MM5 Monin-Obukhov	Revised MM5 Monin-Obukhov	Revised MM5 Monin-Obukhov
Land surface	Pleim-Xiu LSM	Pleim-Xiu LSM	Pleim-Xiu LSM
Planetary Boundary Layer	YSU	YSU	YSU
Cumulus Parameterization	Kain-Fritsch scheme	Kain-Fritsch scheme	None

## 5.2 Photochemical Model

The U.S. EPA modeling guidance (U.S. EPA, 2014) requires several factors to be considered as criteria for choosing a qualifying air quality model to support the attainment demonstration. These criteria include: (1) It should have received a scientific peer review; (2) It should be appropriate for the specific application on a theoretical basis; (3) It should be used with databases which are available and adequate to support its application; (4) It should be shown to have performed well in past modeling applications; and (5). It should be applied consistently with an established protocol on methods and procedures (U.S. EPA, 2014). In addition, it should be well documented with a user’s guide as well as technical descriptions. For the ozone modeled attainment test, a grid-based photochemical model is necessary to offer the best available representation of important atmospheric processes and the ability to analyze the impacts of proposed emission controls on ozone mixing ratios. In ARB’s SIP modeling platform, the Community Multiscale Air Quality (CMAQ) Modeling System has been selected as the air quality model for use in attainment demonstrations of NAAQS for ozone and PM<sub>2.5</sub>.

The CMAQ model, a state-of-the-science “one-atmosphere” modeling system developed by U.S. EPA, was designed for applications ranging from regulatory and policy analysis to investigating the atmospheric chemistry and physics that contribute to air pollution. CMAQ is a three-dimensional Eulerian modeling system that simulates ozone, particulate matter, toxic air pollutants, visibility, and acidic pollutant species throughout the troposphere (UNC, 2010). The model has undergone peer review every

few years and represents the state-of-the-science (Brown et al., 2011). The CMAQ model is regularly updated to incorporate new chemical and aerosol mechanisms, algorithms, and data as they become available in the scientific literature (e.g., Appel et al., 2013; Foley, et al., 2010; Pye and Pouliot, 2012;). In addition, the CMAQ model is well documented in terms of its underlying scientific algorithms as well as guidance on operational uses (e.g., Appel et al., 2013; Binkowski and Roselle, 2003; Byun and Ching, 1999; Byun and Schere, 2006; Carlton et al., 2010; Foley et al., 2010; Kelly, et al., 2010a; Pye and Pouliot, 2012; UNC, 2010).

The CMAQ model was the regional air quality model used for the 2008 SJV annual PM<sub>2.5</sub> SIP (SJVUAPCD, 2008), the 2012 SJV 24-hour PM<sub>2.5</sub> SIP (SJVUAPCD, 2012) and the 2013 SJV 1-hr ozone SIP (SJVUAPCD, 2013). A number of previous studies have also used the CMAQ model to study ozone and PM<sub>2.5</sub> formation in the SJV (e.g., Jin et al., 2008, 2010b; Kelly et al., 2010b; Liang and Kaduwela, 2005; Livingstone, et al., 2009; Pun et al, 2009; Tonse et al., 2008; Vijayaraghavan et al., 2006; Zhang et al., 2010). The CMAQ model has also been used for regulatory analysis for many of U.S. EPA's rules, such as the Clean Air Interstate Rule (U.S. EPA, 2005) and Light-duty and Heavy-duty Greenhouse Gas Emissions Standards (U.S. EPA, 2010, 2011a). There have been numerous applications of the CMAQ model within the U.S. and abroad (e.g., Appel, et al., 2007, 2008; Civerolo et al., 2010; Eder and Yu, 2006; Hogrefe et al., 2004; Lin et al., 2008, 2009; Marmur et al., 2006; O'Neill, et al., 2006; Philips and Finkelstein, 2006; Smyth et al., 2006; Sokhi et al., 2006; Tong et al., 2006; Wilczak et al., 2009; Zhang et al., 2004, 2006), which have shown it to be suitable as a regulatory and scientific tool for investigating air quality. Staff at the CARB has developed expertise in applying the CMAQ model, since it has been used at CARB for over a decade. In addition, technical support for the CMAQ model is readily available from the Community Modeling and Analysis System (CMAS) Center (<http://www.cmascenter.org/>) established by the U.S. EPA.

The version 5.0.2 of the CMAQ model released in May 2014, ([http://www.airqualitymodeling.org/cmaqwiki/index.php?title=CMAQ\\_version\\_5.0.2\\_%28April\\_2014\\_release%29\\_Technical\\_Documentation](http://www.airqualitymodeling.org/cmaqwiki/index.php?title=CMAQ_version_5.0.2_%28April_2014_release%29_Technical_Documentation)), will be used in this SIP modeling platform. Compared to the previous version, CMAQv4.7.1, which was used for the 2012 SJV 24-hour PM<sub>2.5</sub> SIP (SJVUAPCD, 2012) and the 2013 SJV 1-hour ozone SIP (SJVUAPCD, 2013), CMAQ version 5 and above incorporated substantial new features and enhancements to topics such as gas-phase chemistry, aerosol algorithms, and structure of the numerical code ([http://www.airqualitymodeling.org/cmaqwiki/index.php?title=CMAQ\\_version\\_5.0\\_%28February\\_2012\\_release%29\\_Technical\\_Documentation#RELEASE\\_NOTES\\_for\\_CMAQ\\_v5.0\\_.C2.A0February\\_2012](http://www.airqualitymodeling.org/cmaqwiki/index.php?title=CMAQ_version_5.0_%28February_2012_release%29_Technical_Documentation#RELEASE_NOTES_for_CMAQ_v5.0_.C2.A0February_2012)).

### 5.2.1 Photochemical Modeling Domain

Figure 5-2 shows the photochemical modeling domains used by ARB in this modeling platform. The larger domain (dashed black colored box), covering all of California, has a horizontal grid resolution of 12 km and extends from the Pacific Ocean in the west to Eastern Nevada in the east and runs from south of the U.S.-Mexico border in the south to north of the California-Oregon border in the north. The smaller 4 km Northern (green box) and Southern (red box) modeling domains are nested within the outer 12 km domain and utilized to better reflect the finer scale details of meteorology, topography, and emissions. Consistent with the WRF modeling, the 12 km and 4 km CMAQ domains are based on a Lambert Conformal Conic projection with reference longitude at -120.5°W, reference latitude at 37°N, and two standard parallels at 30°N and 60°N. The 30 vertical layers from WRF were mapped onto 18 vertical layers for CMAQ, extending from the surface to 100 mb such that the majority of the vertical layers fall within the planetary boundary layer. This vertical layer structure is based on the WRF sigma-pressure coordinates and the exact layer structure used can be found in Table 5-1. A third 4 km resolution modeling domain (blue box) is nested within the Northern California domain and covers the SJV air basin. This smaller SJV domain may be utilized for PM<sub>2.5</sub> modeling in the SJV if computational constraints (particularly for annual modeling) require the use of a smaller modeling domain. In prior work, modeling results from the smaller SJV domain were compared to results from the larger Northern California domain and no appreciable differences were noted, provided that both simulations utilized chemical boundary conditions derived from the same statewide 12 km simulation.

For the coarse portions of nested regional grids, the U.S. EPA guidance (U.S. EPA, 2014) suggests a grid cell size of 12 km if feasible but not larger than 36 km. For the fine scale portions of nested regional grids, it is desirable to use a grid cell size of ~4 km (U.S. EPA, 2014). Our selection of modeling domains and grid resolution is consistent with this recommendation. The U.S. EPA guidance (U.S. EPA, 2014) does not require a minimum number of vertical layers for an attainment demonstration, although typical applications of “one-atmosphere” models (with the model top at 50-100 mb) are anywhere from 14 to 35 vertical layers. In the ARB’s current SIP modeling platform, 18 vertical layers will be used in the CMAQ model. The vertical structure is based on the sigma-pressure coordinate, with the layers separated at 1.0, 0.9958, 0.9907, 0.9846, 0.9774, 0.9688, 0.9585, 0.9463, 0.9319, 0.9148, 0.8946, 0.8709, 0.8431, 0.8107, 0.7733, 0.6254, 0.293, 0.0788, and 0.0. As previously noted, this also ensures that the majority of the layers are in the planetary boundary layer.




Figure 5-2. CMAQ modeling domains used in this SIP modeling platform. The outer domain (dashed black line) represents the extent of the California statewide domain (shown here with a 4 km horizontal resolution, but utilized in this modeling platform with a 12 km horizontal resolution). Nested higher resolution 4 km modeling domains are highlighted in green and red for Northern/Central California and Southern California, respectively. The smaller SJV PM<sub>2.5</sub> 4 km domain (colored in blue) is nested within the Northern California 4 km domain.

## 5.2.2 CMAQ Model Options

Table 5-3 shows the CMAQv5.0.2 configuration utilized in this modeling platform. The same configuration will be used in all simulations for both ozone and PM<sub>2.5</sub>, and for all modeled years. The Intel FORTRAN compiler version 12 will be used to compile all source codes.

Table 5-3. CMAQ v5.0.2 configuration and settings.

Process	Scheme
Horizontal advection	Yamo (Yamartino scheme for mass-conserving advection)
Vertical advection	WRF-based scheme for mass-conserving advection
Horizontal diffusion	Multi-scale
Vertical diffusion	ACM2 (Asymmetric Convective Model version 2)
Gas-phase chemical mechanism	SAPRC07 gas-phase mechanism with version "C" toluene updates
Chemical solver	EBI (Euler Backward Iterative solver)
Aerosol module	Aero6 (the sixth-generation CMAQ aerosol mechanism with extensions for sea salt emissions and thermodynamics; includes a new formulation for secondary organic aerosol yields)
Cloud module	ACM_AE6 (ACM cloud processor that uses the ACM methodology to compute convective mixing with heterogeneous chemistry for AERO6)
Photolysis rate	phot_inline (calculate photolysis rates in-line using simulated aerosols and ozone)

## 5.2.3 Photochemical Mechanism

The SAPRC07 chemical mechanism will be utilized for all CMAQ simulations. SAPRC07, developed by Dr. William Carter at the University of California, Riverside, is a detailed mechanism describing the gas-phase reactions of volatile organic compounds (VOCs) and oxides of nitrogen (NO<sub>x</sub>) (Carter, 2010a, 2010b). It represents a complete update to the SAPRC99 mechanism, which has been used for previous ozone SIP plans in the SJV. The well-known SAPRC family of mechanisms have been used widely in California and the U.S. (e.g., Baker, et al., 2015; Cai et al., 2011; Chen et

al., 2014; Dennis et al., 2008; Ensberg, et al., 2013; Hakami, et al., 2004a, 2004b; Hu et al., 2012, 2014a, 2014b; Jackson, et al., 2006; Jin et al., 2008, 2010b; Kelly, et al., 2010b; Lane et al., 2008; Liang and Kaduwela, 2005; Livingstone et al., 2009; Lin et al., 2005; Napelenok, 2006; Pun et al., 2009; Tonse et al., 2008; Ying et al., 2008a, 2008b; Zhang et al., 2010; Zhang and Ying, 2011).

The SAPRC07 mechanism has been fully reviewed by four experts in the field through an ARB funded contract. These reviews can be found at <http://www.arb.ca.gov/research/reactivity/rsac.htm>. Dr. Derwent's (2010) review compared ozone impacts of 121 organic compounds calculated using SAPRC07 and the Master Chemical Mechanism (MCM) v 3.1 and concluded that the ozone impacts using the two mechanisms were consistent for most compounds. Dr. Azzi (2010) used SAPRC07 to simulate ozone formation from isoprene, toluene, m-xylene, and evaporated fuel in environmental chambers performed in Australia and found that SAPRC07 performed reasonably well for these data. Dr. Harley discussed implementing the SAPRC07 mechanism into 3-D air quality models and brought up the importance of the rate constant of  $\text{NO}_2 + \text{OH}$ . This rate constant in the SAPRC07 mechanism in CMAQv5.0.2 has been updated based on new research (Mollner et al., 2010). Dr. Stockwell (2009) compared individual reactions and rate constants in SAPRC07 to two other mechanisms (CB05 and RADM2) and concluded that SAPRC07 represented a state-of-the-science treatment of atmospheric chemistry.

### 5.2.4 Aerosol Module

The aerosol mechanism with extensions version 6 with aqueous-phase chemistry (AE6-AQ) will be utilized for all SIP modeling. When coupled with the SAPRC07 chemical mechanism, AE6-AQ simulates the formation and evaporation of aerosol and the evolution of the aerosol size distribution (Foley et al., 2010). AE6-AQ includes a comprehensive, yet computationally efficient, inorganic thermodynamic model ISORROPIA to simulate the physical state and chemical composition of inorganic atmospheric aerosols (Fountoukis and Nenes, 2007). AE6-AQ also features the addition of new  $\text{PM}_{2.5}$  species, an improved secondary organic aerosol (SOA) formation module, as well as new treatment of atmospheric processing of primary organic aerosol (Appel et al., 2013; Carlton et al., 2010; Simon and Bhawe, 2011). These updates to AE6-AQ in CMAQv5.0.2 continue to represent state-of-the-art treatment of aerosol processes in the atmosphere (Brown et al., 2011).

## 5.2.5 CMAQ Initial and Boundary Conditions (IC/BC) and Spin-Up period

Air quality model initial conditions define the mixing ratio (or concentration) of chemical and aerosol species within the modeling domain at the beginning of the model simulation. Boundary conditions define the chemical species mixing ratio (or concentration) within the air entering or leaving the modeling domain. This section discusses the initial and boundary conditions utilized in the ARB modeling system.

U.S. EPA guidance recommends using a model “spin-up” period by beginning a simulation 3-10 days prior to the period of interest (U.S. EPA, 2014). This “spin-up” period allows the initial conditions to be “washed out” of the system, so that the actual initial conditions have little to no impact on the modeling over the time period of interest, as well as giving sufficient time for the modeled species to come to chemical equilibrium. When conducting annual or seasonal modeling, it is computationally more efficient to simulate each month in parallel rather than the entire year or season sequentially. For each month, the CMAQ simulations will include a seven day spin-up period (i.e., the last seven days of the previous month) for the outer 12 km domain to ensure that the initial conditions are “washed out” of the system. Initial conditions at the beginning of the seven day spin-up period will be based on the default initial conditions that are included with the CMAQ release. The 4 km inner domain simulations will utilize a three day spin-up period, where the initial conditions will be based on output from the corresponding day of the 12 km domain simulation.

In recent years, the use of global chemical transport model (CTM) outputs as boundary conditions (BCs) in regional CTM applications has become increasingly common (Chen et al., 2008; Hogrefe et al., 2011; Lam and Fu, 2009; Lee et al., 2011; Lin et al., 2010), and has been shown to improve model performance in many cases (Appel et al., 2007; Borge et al., 2010; Tang et al., 2007, 2009; Tong and Mauzerall, 2006). The advantage of using global CTM model outputs as opposed to fixed climatological-average BCs is that the global CTM derived BCs capture spatial, diurnal, and seasonal variability, as well as provide a set of chemically consistent pollutant mixing ratios. In the ARB’s SIP modeling system, the Model for Ozone And Related chemical Tracers (MOZART; Emmons et al., 2010) will be used to define the boundary conditions for the outer 12 km CMAQ domain, while boundary conditions for the 4 km domain will be derived from the 12 km output. MOZART is a comprehensive global model for simulating atmospheric composition including both gases and bulk aerosols (Emmons et al., 2010). It was developed by the National Center for Atmospheric Research (NCAR), the Max-Planck-Institute for Meteorology (in Germany), and the Geophysical Fluid Dynamics Laboratory (GFDL) of the National Oceanic and Atmospheric Administration (NOAA), and is widely



used in the scientific community. In addition to inorganic gases and VOCs, BCs were extracted for aerosol species including elemental carbon, organic matter, sulfate, soil and nitrate. MOZART has been extensively peer-reviewed and applied in a range of studies that utilize its output in defining BCs for regional modeling studies within California and other regions of the U.S. (e.g., Avise et al., 2008; Chen et al., 2008, 2009a, 2009b; Fast et al., 2014; Jathar et al., 2015).




Figure 5-3. Comparison of MOZART (red) simulated CO (left), ozone (center), and PAN (right) to observations (black) along the DC-8 flight track. Shown are mean (filled symbol), median (open symbols), 10th and 90th percentiles (bars) and extremes (lines). The number of data points per 1-km wide altitude bin is shown next to the graphs. Adapted from Figure 2 in Pfister et al. (2011).

In particular, MOZART version 4 (MOZART-4) was recently used in a study characterizing summertime air masses entering California from the Pacific Ocean (Pfister et al., 2011). In their work, Pfister et al. (2011) compared MOZART-4 simulation results to measurements of CO, ozone, and PAN made off the California coast during the ARCTAS-CARB airborne field campaign (Jacob et al., 2010) and showed good agreement between the observations and model results (see Figure 5-3). The specific MOZART simulations to be utilized in this modeling platform are the MOZART4-GEOS5 simulations by Louisa Emmons (NCAR) for the years 2012 and 2013, which are available for download at <http://www.acom.ucar.edu/wrf-chem/mozart.shtml>. These simulations are similar to those of Emmons et al. (2010), but with updated meteorological fields. Boundary condition data will be extracted from the MOZART-4 output and processed to CMAQ model ready format using the “mozart2camx” code developed by the Rambol-Environ Corporation (available at <http://www.camx.com/download/support-software.aspx>). The final BCs represent day-specific mixing ratios, which vary in both space (horizontal and vertical) and time (every six hours).

Per U.S. EPA guidance, the same MOZART derived BCs for the 12 km outer domain will be used for all simulations (e.g., Base Case, Reference, Future, and any sensitivity simulation).

### **5.3 Quality Assurance of Model Inputs**

In developing the IC/BCs and Four Dimensional Data Assimilation (FDDA) datasets for WRF, quality control is performed on all associated meteorological data. Generally, all surface and upper air meteorological data are plotted in space and time to identify extreme values that are suspected to be “outliers”. Data points are also compared to other, similar surrounding data points to determine whether there are any large relative discrepancies. If a scientifically plausible reason for the occurrence of suspected outliers is not known, the outlier data points are flagged as invalid and may not be used in the modeling analyses.

In addition, the model-ready emissions files used in CMAQ will be evaluated and compared against the planning inventory totals. Although deviations between the model-ready and planning inventories are expected due to temporal adjustments (e.g., month-of-year and day-of-week) and adjustments based on meteorology (e.g., evaporative emissions from motor vehicles and biogenic sources), any excessive deviation will be investigated to ensure the accuracy of the temporal and meteorology based adjustments. If determined to be scientifically implausible, then the adjustments which led to the deviation will be investigated and updated based on the best available science.

Similar to the quality control of the modeling emissions inventory, the chemical boundary conditions derived from the global CTM model will be evaluated to ensure that no errors were introduced during the processing of the data (e.g., during vertical interpolation of the global model data to the regional model vertical structure or mapping of the chemical species). Any possible errors will be evaluated and addressed if they are determined to be actual errors and not an artifact of the spatial and temporal dynamics inherent in the boundary conditions themselves.

## 6. METEOROLOGICAL MODEL PERFORMANCE

The complex interactions between the ocean-land interface, orographic induced flows from the mountain-valley topography, and the extreme temperature gradients between the ocean, delta region, valley floor, and mountain ranges surrounding the valley, make the SJV one of the most challenging areas in the country to simulate using prognostic meteorological models. Although there is a long history of prognostic meteorological model applications in California (e.g., Bao et al., 2008; Hu et al., 2010; Jackson et al., 2006; Jin et al., 2010a, 2010b; Livingstone et al., 2009; Michelson et al., 2010; Seaman, Stauffer, and Lario-Gibbs, 1995; Stauffer et al., 2000; Tanrikulu et al., 2000), there is no single model configuration that works equally well for all years and/or seasons, which makes evaluation of the simulated meteorological fields critical for ensuring that the fields reasonably reproduce the observed meteorology for any given time period.

### 6.1 Ambient Data Base and Quality of Data

Observed meteorological data used to evaluate the WRF model simulations will be obtained from the Air Quality and Meteorological Information System (AQMIS) database, which is a web-based source for real-time and official air quality and meteorological data ([www.arb.ca.gov/airqualitytoday/](http://www.arb.ca.gov/airqualitytoday/)). This database contains surface meteorological observations from 1969-2016, with the data through 2013 having been fully quality assured and deemed official. In addition ARB also has quality-assured upper-air meteorological data obtained using balloons, aircraft, and profilers.

### 6.2 Statistical Evaluation

Statistical analyses will be performed to evaluate how well the WRF model captured the overall structure of the observed atmosphere during the simulation period, using wind speed, wind direction, temperature, and humidity. The performance of the WRF model against observations will be evaluated using the METSTAT analysis tool (Emery et al., 2001) and supplemented using statistical software tools developed at ARB. The model output and observations will be processed, and data points at each observational site for wind speed, wind direction, temperature, and moisture data will be extracted. The following values will be calculated: Mean Obs, Mean Model, Mean Bias (MB), Mean (Gross) Error (ME/MGE), Normalized Mean Bias (NMB), Root Mean Squared error (RMSE), and the Index Of Agreement (IOA) when applicable. Additional statistical analysis may also be performed.

The mathematical expressions for these quantities are:

$$MB = \frac{1}{N} \sum_1^N (\text{Model} - \text{Obs}) \quad (6-1)$$

$$ME = \frac{1}{N} \sum_1^N |\text{Model} - \text{Obs}| \quad (6-2)$$

$$NMB = \frac{\sum_1^N (\text{Model} - \text{Obs})}{\sum_1^N \text{Obs}} \times 100\%, \quad (6-3)$$

$$RSME = \sqrt{\frac{\sum_1^N (\text{Model} - \text{Obs})^2}{N}} \quad (6-4)$$

$$IOA = 1 - \frac{\sum_1^N (\text{Model} - \text{Obs})^2}{\sum_1^N [(\text{Model} - \text{Obs}) + (\text{Model} + \text{Obs})]^2}, \quad (6-5)$$

where, “*Model*” is the simulated values, “*Obs*” is the observed value, and *N* is the number of observations. These values will be tabulated and plotted for all monitoring sites within the air basin of interest, and summarized by subregion when there are distinct differences in the meteorology within the basin. Statistics may be compared to other prognostic model applications in California to place the current model performance within the context of previous studies. In addition to the statistics above, model performance may also be evaluated through metrics such as frequency distributions, time-series analysis, and wind-rose plots. Based on previous experience with meteorological simulations in California, it is expected that the analysis will show wind speed to be overestimated at some stations with a smaller difference at others. The diurnal variations of temperature and wind direction at most stations are likely to be captured reasonably well. However, the model will likely underestimate the larger magnitudes of temperature during the day and smaller magnitudes at night.

### 6.3 Phenomenological Evaluation

In addition to the statistical evaluation described above, a phenomenological based evaluation can provide additional insights as to the accuracy of the meteorological modeling. A phenomenological evaluation may include analysis such as determining the relationship between observed air quality and key meteorological parameters (e.g., conceptual model) and then evaluating whether the simulated meteorology and air quality is able to reproduce those relationships. Another possible approach would be to generate geopotential height charts at 500 and 850 mb using the simulated results and compare those to the standard geopotential height charts. This would reveal if the large-scale weather systems at those pressure levels were adequately simulated by the regional prognostic meteorology model. Another similar approach is to identify the larger-scale meteorological conditions associated with air quality events using the National Centers for Environmental Prediction (NCEP) Reanalysis dataset. These can then be visually compared to the simulated meteorological fields to determine whether those large-scale meteorological conditions were accurately simulated and whether the same relationships observed in the NCEP reanalysis are present in the simulated data.

## 7. PHOTOCHEMICAL MODEL PERFORMANCE

### 7.1 Ambient Data

Air quality observations are routinely made at state and local monitoring stations. Gas species and PM species are measured on various time scales (e.g., hourly, daily, weekly). The U.S. EPA guidance recommends model performance evaluations for the following gaseous pollutants: ozone ( $O_3$ ), nitric acid ( $HNO_3$ ), nitric oxide (NO), nitrogen dioxide ( $NO_2$ ), peroxyacetyl nitrate (PAN), volatile organic compounds (VOCs), ammonia ( $NH_3$ ),  $NO_y$  (sum of  $NO_x$  and other oxidized compounds), sulfur dioxide ( $SO_2$ ), carbon monoxide (CO), and hydrogen peroxide ( $H_2O_2$ ). The U.S. EPA recognizes that not all of these species are routinely measured (U.S. EPA, 2014) and therefore may not be available for evaluating every model application. Recognizing that  $PM_{2.5}$  is a mixture, U.S. EPA recommends model performance evaluation for the following individual  $PM_{2.5}$  species: sulfate ( $SO_4^{2-}$ ), nitrate ( $NO_3^-$ ), ammonium ( $NH_4^+$ ), elemental carbon (EC), organic carbon (OC) or organic mass (OM), crustal, and sea salt constituent (U.S. EPA, 2014).

Table 7-1 lists the species for which routine measurements are generally available in 2012 and 2013. When quality assured data are available and appropriate for use, model performance for each species will be evaluated. Observational data will be

obtained from the Air Quality and Meteorological Information System (AQMIS), which is a web-based source for real-time and official air quality and meteorological data ([www.arb.ca.gov/airqualitytoday/](http://www.arb.ca.gov/airqualitytoday/)). This database contains surface air quality observations from 1980-2016, with the data through 2014 having been fully quality assured and deemed official.

Table 7-1. Monitored species used in evaluating model performance.

<b>Species</b>	<b>Sampling frequency</b>
O <sub>3</sub>	1 hour
NO	1 hour
NO <sub>2</sub>	1 hour
NO <sub>x</sub>	1 hour
CO	1 hour
SO <sub>2</sub>	1 hour
Selected VOCs from the PAMS measurement	3 hours (not every day)
PM <sub>2.5</sub> measured using FRM <sup>1</sup>	24 hours (daily to one in six days)
PM <sub>2.5</sub> measured using FEM	Continuously
PM <sub>2.5</sub> Speciation sites	24 hours (not every day)
Sulfate ion	24 hours (not every day)
Nitrate ion	24 hours (not every day)
Ammonium ion	24 hours (not every day)
Organic carbon	24 hours (not every day)
Elemental carbon	24 hours (not every day)
Sea salt constituents	24 hours (not every day)

<sup>1</sup> Direct comparison between modeled and FRM PM<sub>2.5</sub> may not be appropriate because of various positive and negative biases associated with FRM measurement procedures.

These species cover the majority of pollutants of interest for evaluating model performance as recommended by the U.S. EPA. Other species such as H<sub>2</sub>O<sub>2</sub>, HNO<sub>3</sub>, NH<sub>3</sub>, and PAN are not routinely measured. During the DISCOVER-AQ field campaign, which took place in January and February 2013 in the SJV, aircraft sampling provided daytime measurements for a number of species (including HNO<sub>3</sub>, NH<sub>3</sub>, PAN, alkyl nitrates, and selected VOC species) that are not routinely measured. Modeled concentrations will be compared to aircraft measurements for these species, except for the gaseous HNO<sub>3</sub> measurements, which were contaminated by particulate nitrate (Dr. Chris Cappa, personal communication).

## 7.2 Statistical Evaluation

As recommended by U.S. EPA, a number of statistical metrics will be used to evaluate model performance for ozone, speciated and total PM<sub>2.5</sub>, as well as other precursor species. These metrics may include mean bias (MB), mean error (ME), mean fractional bias (MFB), mean fractional error (MFE), normalized mean bias (NMB), normalized mean error (NME), root mean square error (RMSE), correlation coefficient (R<sup>2</sup>), mean normalized bias (MNB), and mean normalized gross error (MNGE). The formulae for estimating these metrics are given below.

$$MB = \frac{1}{N} \sum_1^N (\text{Model} - \text{Obs}) \quad (7-1)$$

$$ME = \frac{1}{N} \sum_1^N |\text{Model} - \text{Obs}| \quad (7-2)$$

$$MFB = \frac{2}{N} \sum_1^N \left( \frac{\text{Model} - \text{Obs}}{\text{Model} + \text{Obs}} \right) \times 100\%, \quad (7-3)$$

$$MFE = \frac{2}{N} \sum_1^N \left( \frac{|\text{Model} - \text{Obs}|}{\text{Model} + \text{Obs}} \right) \times 100\%, \quad (7-4)$$

$$NMB = \frac{\sum_1^N (\text{Model} - \text{Obs})}{\sum_1^N \text{Obs}} \times 100\%, \quad (7-5)$$

$$NME = \frac{\sum_1^N |\text{Model} - \text{Obs}|}{\sum_1^N \text{Obs}} \times 100\%, \quad (7-6)$$

$$RSME = \sqrt{\frac{\sum_1^N (\text{Model} - \text{Obs})^2}{N}} \quad (7-7)$$

$$R^2 = \left( \frac{\sum_1^N ((\text{Model} - \overline{\text{Model}}) \times (\text{Obs} - \overline{\text{Obs}}))}{\sqrt{\sum_1^N (\text{Model} - \overline{\text{Model}})^2 \sum_1^N (\text{Obs} - \overline{\text{Obs}})^2}} \right)^2 \quad (7-8)$$

$$MNB = \frac{1}{N} \sum_1^N \left( \frac{\text{Model} - \text{Obs}}{\text{Obs}} \right) \times 100\%, \quad (7-9)$$

$$MNGE = \frac{1}{N} \sum_1^N \left( \frac{|\text{Model} - \text{Obs}|}{\text{Obs}} \right) \times 100\%. \quad (7-10)$$

where, "Model" is the simulated mixing ratio, " $\overline{\text{Model}}$ " is the simulated mean mixing ratio, "Obs" is the observed value, " $\overline{\text{Obs}}$ " is the mean observed value, and "N" is the number of observations.

In addition to the above statistics, various forms of graphics will also be created to visually examine and compare the model predictions to observations. These will include time-series plots comparing the predictions and observations, scatter plots for



comparing the magnitude of the simulated and observed mixing ratios, box plots to summarize the time series data across different regions and averaging times, as well as frequency distributions. For  $PM_{2.5}$  the so called “bugle plots” of MFE and MFB from Boylan and Russell (2006) will also be generated. The plots described above will be created for paired observations and predictions over time scales dictated by the averaging frequencies of observations (i.e., hourly, daily, monthly, seasonally) for the species of interest. Together, they will provide a detailed view of model performance during different time periods, in different sub-regions, and over different concentrations and mixing ratio levels.

### 7.3 Comparison to Previous Modeling Studies

Previous U.S. EPA modeling guidance (U.S. EPA, 1991) utilized “bright line” criteria for the performance statistics that distinguished between adequate and inadequate model performance. In the latest modeling guidance from U.S. EPA (U.S. EPA, 2014) it is now recommended that model performance be evaluated in the context of similar modeling studies to ensure that the model performance approximates the quality of those studies. The work of Simon et al. (2012) summarized photochemical model performance for studies published in the peer-reviewed literature between 2006 and 2012 and this work will form the basis for evaluating the modeling utilized in the attainment demonstration.

### 7.4 Diagnostic Evaluation

Diagnostic evaluations are useful for investigating whether the physical and chemical processes that control ozone and  $PM_{2.5}$  formation are correctly represented in the modeling. These evaluations can take many forms, such as utilizing model probing tools like process analysis, which tracks and apportions ozone mixing ratios in the model to various chemical and physical processes, or source apportionment tools that utilize model tracers to attribute ozone formation to various emissions source sectors and/or geographic regions. Sensitivity studies (either “brute-force” or the numerical Direct Decoupled Method) can also provide useful information as to the response exhibited in the modeling to changes in various input parameters, such as changes to the emissions inventory or boundary conditions. Due to the nature of this type of analysis, diagnostic evaluations can be very resource intensive and the U.S. EPA modeling guidance acknowledges that air agencies may have limited resources and time to perform such analysis under the constraints of a typical SIP modeling application. To the extent possible, some level of diagnostic evaluation will be included in the model attainment demonstration for this SIP.

In addition to the above analysis, the 2013 DISCOVER-AQ field campaign in the SJV offers a unique dataset for additional diagnostic analysis that is not available in other areas, in particular, the use of indicator ratios in determining the sensitivity of secondary  $PM_{2.5}$  to its limiting precursors. As an example, the ratio between free ammonia (total ammonia – 2 x sulfate) and total nitrate (gaseous + particulate) was proposed by Ansari and Pandis (1998) as an indicator of whether ammonium nitrate formation is limited by  $NO_x$  or ammonia emissions. The DISCOVER-AQ dataset will be utilized to the extent possible to investigate  $PM_{2.5}$  precursor sensitivity in the SJV as well as analysis of upper measurements and detailed ground level AMS measurements (Young et al., 2016).

## 8. ATTAINMENT DEMONSTRATION

The U.S. EPA modeling guidance (U.S. EPA, 2014) outlines the approach for utilizing models to predict future attainment of the 0.075 ppm 8-hour ozone standard. Consistent with the previous modeling guidance (U.S. EPA, 2007) utilized in the most recent 8-hour ozone (2007), annual  $PM_{2.5}$  (2008), and 24-hour  $PM_{2.5}$  (2012) SIPs, the current guidance recommends utilizing modeling in a relative sense. A detailed description of how models are applied in the attainment demonstration for both ozone and  $PM_{2.5}$ , as prescribed by U.S. EPA modeling guidance, is provided below.

### 8.1 Base Year Design Values

The starting point for the attainment demonstration is with the observational based design value (DV), which is used to determine compliance with the standard at any given monitor. The DV for a specific monitor and year represents the three-year average of the annual 4<sup>th</sup> highest 8-hour ozone mixing ratio, 98<sup>th</sup> percentile of the 24-hour  $PM_{2.5}$  concentration, or annual average  $PM_{2.5}$  concentration, depending on the standard, observed at the monitor. For example, the 8-hr  $O_3$  DV for 2012 is the average of the observed 4<sup>th</sup> highest 8-hour ozone mixing ratio from 2010, 2011, and 2012.

The U.S. EPA recommends using an average of three DVs to better account for the year-to-year variability inherent in meteorology. Since 2012 has been chosen as the base year for projecting DVs to the future, site-specific DVs will be calculated for the three three-year periods ending in 2012, 2013, and 2014 and then these three DVs will be averaged. This average DV is called a weighted DV (in the context of this SIP, the weighted DV will also be referred to as the reference year DV or  $DV_R$ ). Table 8-1 illustrates how the weighted DV is calculated.

Table 8-1. Illustrates the data from each year that are utilized in the Design Value calculation for that year (DV Year), and the yearly weighting of data for the weighted Design Value calculation (or DV<sub>R</sub>). “obs” refers to the observed metric (8-hr O<sub>3</sub>, 24-hour PM<sub>2.5</sub>, or annual average PM<sub>2.5</sub>).

DV Year	Years Averaged for the Design Value (4 <sup>th</sup> highest observed 8-hr O <sub>3</sub> , 98 <sup>th</sup> percentile 24-hour PM <sub>2.5</sub> , or annual average PM <sub>2.5</sub> )				
2012	2010	2011	2012		
2013		2011	2012	2013	
2014			2012	2013	2014
Yearly Weightings for the Weighted Design Value Calculation					
2012-2014 Average	$DV_R = \frac{obs_{2010} + (2)obs_{2011} + (3)obs_{2012} + (2)obs_{2013} + obs_{2014}}{9}$				

## 8.2 Base, Reference, and Future Year Simulations

Projecting the weighted DVs to the future requires three photochemical model simulations as described below:

### 1. Base Year Simulation

The base year simulation for 2012 or 2013 is used to assess model performance (i.e., to ensure that the model is reasonably able to reproduce the observed ozone mixing ratios). Since this simulation will be used to assess model performance, it is essential to include as much day-specific detail as possible in the emissions inventory, including, but not limited to hourly adjustments to the motor vehicle and biogenic inventories based on observed local meteorological conditions, known wildfire and agricultural burning events, and exceptional events such as the Chevron refinery fire in 2012.

### 2. Reference Year Simulation

The reference year simulation is identical to the base year simulation, except that certain emissions events which are either random and/or cannot be projected to the future are removed from the emissions inventory. These include wildfires and events such as the 2012 Chevron refinery fire.

### 3. Future Year Simulation

The future year simulation is identical to the reference year simulation, except that the projected future year anthropogenic emission levels are used rather than the reference year emission levels. All other model inputs (e.g., meteorology, chemical boundary conditions, biogenic emissions, and calendar

for day-of-week specifications in the inventory) are the same as those used in the reference year simulation.

The base year simulation is solely used for evaluating model performance, while the reference and future year simulations are used to project the weighted DV to the future as described in subsequent sections of this document.

### 8.3 Relative Response Factors

As part of the model attainment demonstration, the fractional change in ozone or PM<sub>2.5</sub> between the model future year and model reference year are calculated for each monitor location. These ratios, called “relative response factors” or RRFs, are calculated based on the ratio of modeled future year ozone or PM<sub>2.5</sub> to the corresponding modeled reference year ozone or PM<sub>2.5</sub> (Equation 8-1).

$$\text{RRF} = \frac{\text{average } (O_3 \text{ or } PM_{2.5})_{\text{future}}}{\text{average } (O_3 \text{ or } PM_{2.5})_{\text{reference}}} \quad (8-1)$$

#### 8.3.1 8-hour Ozone RRF

For 8-hour ozone, the modeled maximum daily average 8-hour (MDA8) ozone is used in calculating the RRF. These MDA8 ozone values are based on the maximum simulated ozone within a 3x3 array of cells surrounding the monitor (Figure 8-1). The future and base year ozone values used in RRF calculations are paired in space (i.e., using the future year MDA8 ozone value at the same grid cell where the MDA8 value for the reference year is located within the 3x3 array of cells). The days used to calculate the average MDA8 for the reference and future years are inherently consistent, since the same meteorology is used to drive both simulations.

Not all modeled days are used to calculate the average MDA8 ozone from the reference and future year simulations. The form of the 8-hour ozone NAAQS is such that it is geared toward the days with the highest mixing ratios in any ozone season (i.e., the 4<sup>th</sup> highest MDA8 ozone). Therefore, the modeled days used in the RRF calculation should also reflect days with the highest ozone levels. As a result, the current U.S. EPA guidance (U.S. EPA, 2014) suggests using the top 10 modeled days when calculating the RRF. Since the relative sensitivity to emissions changes (in both the model and real world) can vary from day-to-day due to meteorology and emissions (e.g., temperature dependent emissions or day-of-week variability) using the top 10 days ensures that the

calculated RRF is robust and stable (i.e., not overly sensitive to any single day used in the calculation).

When choosing the top 10 days, the U.S. EPA recommends beginning with all days in which the simulated reference MDA8 is  $\geq 60$  ppb and then calculating RRFs based on the top 10 high ozone days. If there are fewer than 10 days with MDA8 ozone  $\geq 60$  ppb then all days  $\geq 60$  ppb are used in the RRF calculation, as long as there are at least 5 days used in the calculation. If there are fewer than 5 days  $\geq 60$  ppb, an RRF cannot be calculated for that monitor. To ensure that only modeled days which are consistent with the observed ozone levels are used in the RRF calculation, the modeled days are further restricted to days in which the reference MDA8 ozone is within  $\pm 20\%$  of the observed value at the monitor location.




Figure 8-1. Example showing how the location of the MDA8 ozone for the top ten days in the reference and future years are chosen.

### 8.3.2 Annual and 24-hour $PM_{2.5}$ RRF

The U.S. EPA (2014) guidance requires RRFs for both the annual and 24-hour  $PM_{2.5}$  attainment tests be calculated on a quarterly basis (January-March, April-June, July-September, and October-December) and for each  $PM_{2.5}$  component (sulfate, nitrate, ammonium, organic carbon, elemental carbon, particle bound water, salt, and other primary inorganic components).

For annual  $PM_{2.5}$ , the quarterly RRFs are based on modeled quarterly mean concentrations for each component, where the concentrations are averaged over the 9 model grid cells within the 3x3 array of grid cells surrounding each monitor. For the 24-hour  $PM_{2.5}$  attainment test, the quarterly RRFs are calculated based on the average for

each component over the top 10% of modeled days (or the top nine days per quarter) with the highest total 24-hour average PM<sub>2.5</sub> concentration. Peak PM<sub>2.5</sub> values are selected and averaged using the PM<sub>2.5</sub> concentration simulated at the single grid cell containing the monitoring site for calculating the 24-hour PM<sub>2.5</sub> RRF (as opposed to the 3x3 array average used in the annual PM<sub>2.5</sub> RRF calculation).

## 8.4 Future Year Design Value Calculation

### 8.4.1 8-hour Ozone

For 8-hour ozone, a future year DV at each monitor is calculated by multiplying the corresponding reference year DV by the site-specific RRF from Equation 8-1 (Equation 8-2).

$$DV_F = DV_R \times RRF \quad (8-2)$$

where,

DV<sub>F</sub> = future year design value,

DV<sub>R</sub> = reference year design value, and

RRF = the site specific RRF from Equation 8-1

The resulting future year DVs are then compared to the 8-hour ozone NAAQS to demonstrate whether attainment will be reached under the future emissions scenario utilized in the future year modeling. A monitor is considered to be in attainment of the 8-hour ozone standard if the estimated future design value does not exceed the level of the standard.

### 8.4.2 Annual and 24-hour PM<sub>2.5</sub>

#### 8.4.2.1 Sulfate, Adjusted Nitrate, Derived, Water, Inferred Carbonaceous Material Balance Approach (SANDWICH) and Potential Modifications

Federal Reference Method (FRM) PM<sub>2.5</sub> mass measurements provide the basis for the attainment/nonattainment designations. For this reason it is recommended that the FRM data be used to project future air quality and progress towards attainment. However, given the complex physicochemical nature of PM<sub>2.5</sub>, it is necessary to consider individual PM<sub>2.5</sub> species as well. While the FRM measurements give the mass

of the bulk sample, a method for apportioning this bulk mass to individual  $PM_{2.5}$  components is the first step towards determining the best emissions controls strategies to reach NAAQS levels in a timely manner.

The FRM measurement protocol finds its roots in the past epidemiological studies of health effects associated with  $PM_{2.5}$  exposure. It is upon these studies that the NAAQS are based. The FRM protocol is sufficiently detailed so that results might be easily reproducible and involves the measurement of filter mass before and after sampling together with equilibrating at narrowly defined conditions. Filters are equilibrated for more than 24 hours at a standard relative humidity between 30 and 40% and temperature between 20 and 23 °C. Due to the sampler construction and a lengthy filter equilibration period, FRM measurements are subjected to a number of known positive and negative artifacts. FRM measurements do not necessarily capture the  $PM_{2.5}$  concentrations in the atmosphere and can differ substantially from what is measured by speciation monitors including the Speciation Trends Network (STN) monitors (see <http://www.epa.gov/ttnamti1/specgen.html> for more details). Nitrate and semi-volatile organic mass can be lost from the filter during the equilibration process, and particle bound water associated with hygroscopic species like sulfate provides a positive artifact. These differences present an area for careful consideration when one attempts to utilize speciated measurements to apportion the bulk FRM mass to individual species. Given that (1) attainment status is currently dependent upon FRM measurements and (2) concentrations of individual  $PM_{2.5}$  species need to be considered in order to understand the nature of and efficient ways to ameliorate the  $PM_{2.5}$  problem in a given region, a method has been developed to speciate bulk FRM  $PM_{2.5}$  mass with known FRM limitations in mind. This method is referred to as the measured **Sulfate, Adjusted Nitrate, Derived Water, Inferred Carbonaceous** material balance approach or “SANDWICH” (Frank, 2006). SANDWICH is based on speciated measurements from other (often co-located) samplers, such as those from STN, and the known sampling artifacts of the FRM. The approach strives to provide mass closure, reconciliation between speciated and bulk mass concentration measurements, and the basis for a connection between observations, modeled  $PM_{2.5}$  concentrations, and the air quality standard (U.S. EPA, 2014).

The main steps in estimating the  $PM_{2.5}$  composition are as follows:

- (1) Calculate the nitrate retained on the FRM filter using hourly relative humidity and temperature together with the STN nitrate measurements,**

The FRM does not retain all of the semi-volatile  $PM_{2.5}$  mass, and at warmer temperatures, loss of particulate nitrate from filters has been commonly observed (Chow et al., 2005). In order to estimate how much nitrate is retained on the FRM filter,

simple thermodynamic equilibrium relations may be used. Necessary inputs include 24-hour average nitrate measurements and hourly temperature and relative humidity data. Frank (2006) suggests the following methodology for estimating retained nitrate. For each hour  $i$  of the day, calculate the dissociation constant,  $K_i$  from ambient temperature and relative humidity (RH).

For  $RH < 61\%$ :

$$\ln(K_i) = 118.87 - (24084/T_i) - 6.025 \times \ln(T_i),$$

where,  $T_i$  is the hourly temperature in Kelvins and  $K_i$  is in nanobars.

For  $RH \geq 61\%$ ,  $K_i$  is replaced by:

$$K'_i = [P_1 - P_2(1 - a_i) + P_3(1 - a_i)^2] \times (1 - a_i)^{1.75} \times K_i,$$

where,  $a_i$  is "fractional" relative humidity and

$$\ln(P_1) = -135.94 + 8763/T_i + 19.12 \times \ln(T_i),$$

$$\ln(P_2) = -122.65 + 9969/T_i + 16.22 \times \ln(T_i),$$

$$\ln(P_3) = -182.61 + 13875/T_i + 24.46 \times \ln(T_i).$$

Using this information, calculate the nitrate retained on the filter as:

$$\text{Retained Nitrate} = \text{STN nitrate} - 745.7/T_R \times (\kappa - \gamma) \times \frac{1}{24} \sum_{i=1}^{24} \sqrt{K_i},$$

where,  $T_R$  is the daily average temperature for the sampled air volume in Kelvin,  $K_i$  is the dissociation constant for  $\text{NH}_4\text{NO}_3$  at ambient temperature for hour  $i$ , and  $(\kappa - \gamma)$  relates to the temperature rise of the filter and vapor depletion from the inlet surface and is assumed to have a value equal to one (Hering and Cass, 1999).

**(2) Calculate quarterly averages for retained nitrate, sulfate, elemental carbon, sea salt, and ammonium,**



**(3) Calculate particle bound water using the concentrations of ammonium, sulfate, and nitrate, using an equilibrium model like the Aerosol Inorganic Model (AIM) or a polynomial equation derived from model output**

Under the FRM filter equilibration conditions, hygroscopic aerosol will retain its particle bound water (PBW) and be included in the observed FRM PM<sub>2.5</sub> mass. PBW can be calculated using an equilibrium model like the Aerosol Inorganics Model (AIM). AIM requires the concentrations of ammonium, nitrate, sulfate, and estimated H<sup>+</sup> as inputs. In addition to inorganic concentrations, the equilibration conditions are also necessary model inputs. In this case, a temperature of 294.15 K and 35% RH is recommended. Alternatively, for simplification, a polynomial regression equation may be constructed by fitting the calculated water concentration from an equilibrium model and the concentrations of nitrate, ammonium, and sulfate. The AIM model will be used for more accurate calculation of PBW.

**(4) Add 0.5 µg/m<sup>3</sup> as blank mass, and**

**(5) Calculate organic carbon mass (OCMmb) by difference, subtracting all inorganic species (including blank mass) from the PM<sub>2.5</sub> mass.**

Other components that may be represented on the FRM filter include elemental carbon, crustal material, sea salt, and passively collected mass. Depending on location certain species may be neglected (e.g., sea salt for inland areas).

While carbonaceous aerosol may make up a large portion of airborne aerosol, speciated measurements of carbonaceous PM are considered highly uncertain. This is due to the large number of carbon compounds in the atmosphere and the measurement uncertainties associated with samplers of different configurations. In the SANDWICH approach, organic carbonaceous mass is calculated by difference. The sum of all nonorganic carbon components will be subtracted from the FRM PM<sub>2.5</sub> mass to estimate the mass of organic carbon.

After having calculated the species concentrations as outlined above, we will calculate the percentage contribution of each species to the measured FRM mass (minus the blank concentration of 0.5 µg/m<sup>3</sup>) for each quarter of the years represented by the speciated data. Note that blank mass is kept constant at 0.5 µg/m<sup>3</sup> between the base and future years, and future year particle bound water needs to be calculated for the future year values of nitrate, ammonium, and sulfate.

#### **8.4.2.2 Estimation of Species Concentrations at Federal Reference Method (FRM) Monitors that Lack Speciation Data**

Speciation data from available STN (speciation) sites will be used to speciate the FRM mass for all FRM sites. For those sites not collocated with STN monitors, surrogate speciation sites will be determined based on proximity and evaluation of local emissions or based on similarity in speciation profiles if such data exists (e.g., such as the speciated data collected in the SJV during CRPAQS (Solomon and Magliano, 1998)).

#### **8.4.2.3 Speciated Modeled Attainment Test (SMAT)**

Following U.S. EPA modeling guidance (U.S. EPA, 2014), the model attainment test for the annual  $PM_{2.5}$  standard will be performed with the following steps.

Step 1: For each year used in the design value calculation, determine the observed quarterly mean  $PM_{2.5}$  and quarterly mean composition for each monitor by multiplying the monitored quarterly mean concentration of FRM derived  $PM_{2.5}$  by the fractional composition of  $PM_{2.5}$  species for each quarter.

Step 2: Calculate the component specific RRFs at each monitor for each quarter as described in section 8.3.2.

Step 3: Apply the component specific RRFs to the quarterly mean concentrations from Step 1 to obtain projected quarterly species estimates.

Step 4: Calculate future year annual average  $PM_{2.5}$  estimates by summing the quarterly species estimates at each monitor and then compare to the annual  $PM_{2.5}$  NAAQS. If the projected average annual arithmetic mean  $PM_{2.5}$  concentration is  $\leq$  the NAAQS, then the attainment test is passed.

For the 24-hour  $PM_{2.5}$  standard, the attainment test is performed with the following steps (U.S. EPA, 2014):

Step 1: Determine the top eight days with the highest observed 24-hour  $PM_{2.5}$  concentration (FRM sites) in each quarter and year used in the design value calculation (a total of 32 days per year), and calculate the 98<sup>th</sup> percentile value for each year.

Step 2: Calculate quarterly ambient species fractions on “high” PM<sub>2.5</sub> days for each of the major PM<sub>2.5</sub> component species (i.e., sulfate, nitrate, ammonium, elemental carbon, organic carbon, particle bound water, salt, and blank mass). The “high” days are represented by the top 10% of days in each quarter. Depending on the sampling frequency, the number of days captured in the top 10% would range from three to nine. The species fractions of PM<sub>2.5</sub> are calculated using the “SANDWICH” approach which was described previously. These quarter-specific fractions along with the FRM PM<sub>2.5</sub> concentrations are then used to calculate species concentrations for each of the 32 days per year determined in Step 1.

Step 3: Apply the component and quarter specific RRF, described in Section 8.3.2, to observed daily species concentrations from Step 2 to obtain future year concentrations of sulfate, nitrate, elemental carbon, organic carbon, salt, and other primary PM<sub>2.5</sub>.

Step 4: Calculate the future year concentrations for the remaining PM<sub>2.5</sub> components (i.e., ammonium, particle bound water, and blank mass). The future year ammonium is calculated based on the calculated future year sulfate and nitrate, using a constant value for the degree of neutralization of sulfate from the ambient data. The future year particle bound water is calculated from the AIM model.

Step 5: Sum the concentration of each of the species components to calculate the total PM<sub>2.5</sub> concentration for each of the 32 days per year and at each site. Sort the 32 days for each site and year, and calculate the 98<sup>th</sup> percentile value corresponding to each year.

Step 6: Calculate the future design value at each site based on the 98<sup>th</sup> percentile concentrations calculated in Step 5 and following the standard protocol for calculating design values (see Table 8-1). Compare the future-year 24-hour design values to the NAAQS. If the projected design value is ≤ the NAAQS, then the attainment test is passed.

#### 8.4.2.4 Sensitivity Analyses

Model sensitivity analysis may be conducted if the model attainment demonstration does not show attainment of the applicable standard with the baseline future inventory, or for determining precursor sensitivities and inter-pollutant equivalency ratios. For both ozone and PM<sub>2.5</sub>, the sensitivity analysis will involve domain wide fractional reductions of the appropriate anthropogenic precursor emissions using the future year baseline emissions scenario as a starting point. In the event that the model attainment demonstration does not show attainment for the applicable standard, it is important to know the precursor limitation to assess the level of emissions controls needed to attain the standard.

In order to identify what combinations of precursor emissions reductions is predicted to lead to attainment, a series of modeling sensitivity simulations with varying degrees of precursor reductions from anthropogenic sources are typically performed. These sensitivity simulations are identical to the baseline future year simulation discussed earlier except that domain-wide fractional reductions are applied to future year anthropogenic precursor emission levels and a new future year design value is calculated. The results of these sensitivity simulations are plotted on isopleth diagrams, which are also referred to as carrying capacity diagrams. The isopleths provide an estimate of the level of emissions needed to demonstrate attainment and thereby inform the development of a corresponding control strategy.

For ozone, this would likely entail reducing anthropogenic NO<sub>x</sub> and VOC emissions in 25% increments including cross sensitivities (e.g., 0.75 x NO<sub>x</sub> + 1.00 x VOC; 1.00 x NO<sub>x</sub> + 0.75 x VOC; 0.75 x NO<sub>x</sub> + 0.75 x VOC; 0.5 x NO<sub>x</sub> + 1.00 x VOC; ...). Typically, a full set of sensitivities would include simulations for 25%, 50%, and 75% reduction in NO<sub>x</sub> and VOC, along with the cross sensitivities (for a total of 16 simulations including the future base simulation). After design values are calculated for each new sensitivity simulation, an ozone isopleth (or carrying capacity diagram) as a function of NO<sub>x</sub> and VOC emissions is generated and used to estimate the additional NO<sub>x</sub> and VOC emission reductions needed to attain the standard. The approach for PM<sub>2.5</sub> is similar, except that additional precursor emissions must be considered. Typically, the precursors considered for PM<sub>2.5</sub> would include anthropogenic NO<sub>x</sub>, SO<sub>x</sub>, VOCs, NH<sub>3</sub>, as well as direct PM<sub>2.5</sub> emissions (Chen et al., 2014). Cross sensitivities for generating PM<sub>2.5</sub> carrying capacity diagrams would be conducted with respect to NO<sub>x</sub>, which would include the following precursor pairs: NO<sub>x</sub> vs. primary PM<sub>2.5</sub>, NO<sub>x</sub> vs. VOC, NO<sub>x</sub> vs. NH<sub>3</sub>, and NO<sub>x</sub> vs. SO<sub>x</sub>.

In addition to the PM<sub>2.5</sub> carrying capacity simulations, precursor sensitivity modeling may be conducted for determining the significant precursors to PM<sub>2.5</sub> formation and for

developing inter-pollutant equivalency ratios. These simulations would follow a similar approach to the carrying capacity simulations described above, but would involve only a single sensitivity simulation for each precursor, where emissions of that precursor are reduced between 30% and 70% from the future base year. The “effectiveness” of reducing a given species can be quantified at each FRM monitor as the change in  $\mu\text{g PM}_{2.5}$  (i.e., change in design value) per ton of precursor emissions (corresponding to the 15% change in emissions). Equivalency ratios between  $\text{PM}_{2.5}$  precursors (i.e.,  $\text{NO}_x$ ,  $\text{SO}_x$ , VOCs, and  $\text{NH}_3$ ) and primary  $\text{PM}_{2.5}$  will be determined by dividing primary  $\text{PM}_{2.5}$  effectiveness by the precursors’ effectiveness.

## 8.5 Unmonitored Area Analysis

The unmonitored area analysis is used to ensure that there are no regions outside of the existing monitoring network that could exceed the NAAQS if a monitor was present at that location (U.S. EPA, 2014). The U.S. EPA recommends combining spatially interpolated design value fields with modeled gradients for the pollutant of interest (e.g. Ozone and  $\text{PM}_{2.5}$ ) and grid-specific RRFs in order to generate gridded future year gradient adjusted design values. The spatial Interpolation of the observed design values is done only within the geographic region constrained by the monitoring network, since extrapolating to outside of the monitoring network is inherently uncertain. This analysis can be done using the Model Attainment Test Software (MATS) (Abt, 2014); however this software is not open source and comes as a precompiled software package. To maintain transparency and flexibility in the analysis, in-house R codes (<https://www.r-project.org/>) developed at ARB will be utilized in this analysis. The basic steps followed in the unmonitored area analysis for 8-hour ozone and annual/24-hour  $\text{PM}_{2.5}$  are described below.

### 8.5.1 8-hour Ozone

In this section, the specific steps followed in 8-hr ozone unmonitored area analysis are described briefly:

Step 1: At each grid cell, the top-10 modeled maximum daily average 8-hour ozone mixing ratios from the reference year simulation will be averaged, and a gradient in this top-10 day average between each grid cell and grid cells which contain a monitor will be calculated.

Step 2: A single set of spatially interpolated 8-hr ozone DV fields will be generated based on the observed 5-year weighted base year 8-hr ozone DVs from the available monitors. The interpolation is done using normalized inverse

distance squared weightings for all monitors within a grid cell's Voronoi Region (calculated with the R tripack library; <https://cran.r-project.org/web/packages/tripack/README>), and adjusted based on the gradients between the grid cell and the corresponding monitor from Step 1.

Step 3: At each grid cell, the RRFs are calculated based on the reference- and future-year modeling following the same approach outlined in Section 8.3, except that the +/- 20% limitation on the simulated and observed maximum daily average 8-hour ozone is not applicable because observed data do not exist for grid cells in unmonitored areas.

Step 4: The future year gridded 8-hr ozone DVs are calculated by multiplying the gradient-adjusted interpolated 8-hr ozone DVs from Step 2 with the gridded RRFs from Step 3

Step 5: The future-year gridded 8-hr ozone DVs (from Step 4) are examined to determine if there are any peak values higher than those at the monitors, which could potentially cause violations of the applicable 8-hr ozone NAAQS.

### 8.5.2 Annual PM<sub>2.5</sub>

The unmonitored area analysis for the annual PM<sub>2.5</sub> standard will include the following steps:

Step 1: At each grid cell, the quarterly average PM<sub>2.5</sub> (total and by species) will be calculated from the reference year simulation, and a gradient in these quarterly averages between each grid cell and grid cells which contain a monitor will be calculated.

Step 2: Interpolated spatial fields, based on the observed PM<sub>2.5</sub> (FRM) and each component species of PM<sub>2.5</sub>, will be generated for each quarter using normalized inverse distance squared weightings for all monitors within a grid cell's Voronoi Region. The ambient interpolated spatial fields are then adjusted based on the gradients in predicted quarterly mean concentrations from Step 1.

Step 3: The component specific RRFs are calculated at each grid cell for each quarter as described in section 8.3.2.

Step 4: The quarterly mean concentrations from Step 2 are then multiplied by the corresponding component specific RRF (from Step 3) to obtain the corresponding projected quarterly species estimates.

Step 5: The future year annual average  $PM_{2.5}$  estimates are calculated by summing the quarterly species estimates at each grid cell and then compared to the annual  $PM_{2.5}$  NAAQS to determine compliance.

### 8.5.3 24-hour $PM_{2.5}$

The unmonitored area analysis for the 24-hour  $PM_{2.5}$  standard will include the following steps:

Step 1: At each grid cell, the quarterly average of the top 10% of the modeled days for 24-hour  $PM_{2.5}$  (total and by species for the same top 10% of days) will be calculated from the reference year simulation, and a gradient in these quarterly averages between each grid cell and grid cells which contain a monitor will be calculated.

Step 2: The top 8 days with observed high  $PM_{2.5}$  (FRM) are identified for each quarter and for each of the five years (a total of 32 days per year), used in the base year DV calculation. The speciated  $PM_{2.5}$  (FRM) values are then interpolated for each of the "high"  $PM_{2.5}$  days (identified above) using normalized inverse distance squared weightings for all monitors within a grid cell's Voronoi Region. These ambient interpolated spatial fields are then adjusted based on the appropriate gradients in predicted concentrations from Step 1.

Step 3: The component specific RRFs are calculated at each grid cell for each quarter as described in section 8.3.2.

Step 4: The observed daily species concentrations from Step 2 are multiplied by the component and quarter specific RRF (from Step 3) to estimate the future year concentration of each  $PM_{2.5}$  species using the method outlined in section 8.4.2.3

Step 5: The concentration of each of the component  $PM_{2.5}$  species is summed to calculate the total  $PM_{2.5}$  concentration for each of the 32 days per year (8 days per quarter) and at each grid cell. For each year, the 98<sup>th</sup> percentile value is calculated by the sorting the 32 days for that particular year at each grid cell.

Step 6: The future design value at each grid cell is calculated based on the 98<sup>th</sup> percentile concentrations calculated in Step 5 and following the standard protocol for calculating design values (see Table 8-1). The future-year 24-hour design values are then compared to the 24-hour PM<sub>2.5</sub> NAAQS to determine compliance with that standard.

The R codes used in this analysis will be made available upon request.

## 8.6 Banded Relative Response Factors for Ozone

The “Band-RRF” approach expands upon the standard “Single-RRF” approach for 8-hour ozone to account for differences in model response to emissions controls at varying ozone levels. The most recent U.S. EPA modeling guidance (U. S. EPA, 2014) accounts for some of these differences by focusing on the top ten modeled days, but even the top ten days may contain a significant range of ozone mixing ratios. The Band-RRF approach accounts for these differences more explicitly by grouping the simulated ozone into bands of lower, medium, and higher ozone mixing ratios. Specifically, daily peak 8-hour ozone mixing ratios for all days meeting model performance criteria (+/- 20% with the observations) can be stratified into 5 ppb increments from 60 ppb upwards (bin size and mixing ratio range may vary under different applications). A separate RRF is calculated for each ozone band following a similar approach as the standard Single-RRF. A linear regression is then fit to the data resulting in an equation relating RRF to ozone band. Similar to the Single-RRF, this equation is unique to each monitor/location.

The top ten days for each monitor, based on observed 8-hour ozone, for each year that is utilized in the design value calculation (see Table 8-1) is then projected to the future using the appropriate RRF for the corresponding ozone band. The top ten future days for each year are then re-sorted, the fourth highest 8-hour ozone is selected, and the future year design value is calculated in a manner consistent with the base/reference year design value calculation. More detailed information on the Band-RRF approach can be found in Kulkarni et al. (2014) and the 2013 SJV 1-hour ozone SIP (SJVUAPCD, 2013).



## 9. PROCEDURAL REQUIREMENTS

### 9.1 How Modeling and other Analyses will be Archived, Documented, and Disseminated

The computational burden of modeling the entire state of California and its sub-regions requires a significant amount of computing power and large data storage requirements. For example, there are over half a million grid cells in total for each simulation based on the Northern CA domain (192 x 192 cells in the lateral direction and 18 vertical layers). The meteorological modeling system has roughly double the number of grid cells since it has 30 vertical layers. Archiving of all the inputs and outputs takes several terabytes (TB) of computer disk space (for comparison, one single-layer DVD can hold roughly 5 gigabytes (GB) of data, and it would require ~200 DVDs to hold one TB). Please note that this estimate is for simulated surface-level pollutant output only. If three-dimensional pollutant data are needed, it would add a few more TB to this total. Therefore, transferring the modeling inputs/outputs over the internet using file transfer protocol (FTP) is not practical.

Interested parties may send a request for model inputs/outputs to Mr. John DaMassa, Chief of the Modeling and Meteorology Branch at the following address.

John DaMassa, Chief  
Modeling and Meteorology Branch  
Air Quality Planning and Science Division  
Air Resources Board  
California Environmental Protection Agency  
P.O. Box 2815  
Sacramento, CA 95814, USA

The requesting party will need to send an external disk drive(s) to facilitate the data transfer. The requesting party should also specify what input/output files are requested so that ARB can determine the capacity of the external disk drive(s) that the requester should send.

### 9.2 Specific Deliverables to U.S. EPA

The following is a list of modeling-related documents that will be provided to the U.S. EPA.

- The modeling protocol

- Emissions preparation and results
- Meteorology
  - Preparation of model inputs
  - Model performance evaluation
- Air Quality
  - Preparation of model inputs
  - Model performance evaluation
- Documentation of corroborative and weight-of-evidence analyses
- Predicted future year Design Values
- Access to input data and simulation results

## REFERENCES

Abt, 2014. Modeled Attainment Test Software: User's Manual. MATS available at:  
[http://www.epa.gov/scram001/modelingapps\\_mats.htm](http://www.epa.gov/scram001/modelingapps_mats.htm)

Angevine, W.M., Eddington, L., Durkee, K., Fairall, C., Bianco, L., and Brioude, J., 2012, Meteorological model evaluation for CalNex 2010, Monthly Weather Review, 140, 3885-3906.

Ansari, A.S., and Pandis, S.N., 1998, Response of inorganic PM to precursor concentrations, Environmental Science & Technology, 32, 2706-2714.

Appel, K. W., Pouliot, G. A., Simon, H., Sarwar, G., Pye, H. O. T., Napelenok, S. L., Akhtar, F., and Roselle, S. J., 2013, Evaluation of dust and trace metal estimates from the Community Multiscale Air Quality (CMAQ) model version 5.0, Geoscientific Model Development, 6, 883-899, doi:10.5194/gmd-6-883-2013, 2013.

Appel, W. K., Gilliland, A.B., Sarwar, G., and Gilliam, R.C., 2007, Evaluation of the Community Multiscale Air Quality (CMAQ) model version 4.5: Sensitivities impacting model performance: Part I – Ozone, Atmospheric Environment, 41, 9603-9615.

Appel, W.K., Bhave, P.V., Gilliland, A.B., Sarwar, G., and Roselle, S.J., 2008, Evaluation of the Community Multiscale Air Quality (CMAQ) model version 4.5: Sensitivities impacting model performance; Part II – Particulate Matter, Atmospheric Environment, 42, 6057-6066.

Avise, J., Chen, J., Lamb, B., Wiedinmyer, C., Guenther, A., Salathe, E., and Mass, C., 2009, Attribution of projected changes in summertime US ozone and PM<sub>2.5</sub> concentrations to global changes, *Atmospheric Chemistry and Physics*, 9, 1111-1124.

Azzi, M., White, S.J., Angove, D.E., Jamie, I. M., and Kaduwela, A., 2010, Evaluation of the SAPRC-07 mechanism against CSIRO smog chamber data, *Atmospheric Environment*, 44, 1707-1713.

Baker, K. R., Carlton, A. G., Kleindienst, T. E., Offenberg, J. H., Beaver, M. R., Gentner, D. R., Goldstein, A. H., Hayes, P. L., Jimenez, J. L., Gilman, J. B., de Gouw, J. A., Woody, M. C., Pye, H. O. T., Kelly, J. T., Lewandowski, M., Jaoui, M., Stevens, P. S., Brune, W. H., Lin, Y.-H., Rubitschun, C. L., and Surratt, J. D.: Gas and aerosol carbon in California: comparison of measurements and model predictions in Pasadena and Bakersfield, *Atmos. Chem. Phys.*, 15, 5243-5258, doi:10.5194/acp-15-5243-2015, 2015.

Bao, J.W., Michelson, S.A., Persson, P.O.G., Djalalova, I.V., and Wilczak, J.M., 2008, Observed and WRF-simulated low-level winds in a high-ozone episode during the Central California Ozone Study, *Journal of Applied Meteorology and Climatology*, 47(9), 2372-2394.

Binkowski, F.S. and Roselle, S.J., 2003, Models-3 Community Multiscale Air Quality (CMAQ) model aerosol component, 2. Model description, *Journal of Geophysical Research*, 108, D6, doi:10.1029/2001jd001409.

Borge, R., Lopez, J., Lumberras, J., Narros, A., and Rodriguez, E., 2010, Influence of boundary conditions on CMAQ simulations over the Iberian Peninsula, *Atmospheric Environment*, 44, 2681-2695 (doi:10.1016/j.atmosenv.2010.04.044).

Boylan, J.W., Russell, A.G., (2006), PM and light extinction model performance metrics, goals, and criteria for three-dimensional air quality models. *Atmospheric Environment* 40, 4946-4959.

Brown, N., Allen, D.T., Amar, P., Kallos, G., McNider, R., Russell, A.G., and Stockwell, W.R., 2011, Final report: Fourth peer review of the CMAQ model, Submitted to Community Modeling and Analysis System Center, The University of North Carolina at Chapel Hill.

Byun, D.W. and Ching, J.K.S., 1999, Science Algorithms of the EPA Models-3 Community Multiscale Air Quality (CMAQ) Modeling System, EPA/600/R-99/030, available at <http://www.epa.gov/AMD/CMAQ/CMAQscienceDoc.html>

Byun, D.W. and Schere, K.L., 2006, Review of the governing equations, computational algorithms, and other components of the Models-3 Community Multiscale Air Quality (CMAQ) modeling system, *Applied Mechanics Review*, 59, 51-77.

- Cai, C., Kelly, J.T., Avise, J.C., Kaduwela, A.P., and Stockwell, W.R., 2011, Photochemical modeling in California with two chemical mechanisms: Model intercomparison and response to emission reductions, *Journal of the Air & Waste Management Association*, 61, 559-572.
- Carlton, A.G., Bhave, P., Napelenok, S.L., Edney, E.O., Sarwar, G., Pinder, R.W., Pouliot, G.A., and Houyoux, M., 2010, Model representation of secondary organic aerosol in CMAQv4.7, *Environmental Science Technology*, 44, 8553-8560.
- Carter, W.P.L., 2010a, Development of the SAPRC-07 chemical mechanism, *Atmospheric Environment*, 44(40), 5324-5335.
- Carter, W.P.L., 2010b, Development of a condensed SAPRC-07 chemical mechanism, *Atmospheric Environment*, 44(40), 5336-5345.
- Chen, J., Vaughan, J., Avise, J., O'Neill, S., and Lamb, B., 2008, Enhancement and evaluation of the AIRPACT ozone and PM<sub>2.5</sub> forecast system for the Pacific Northwest, *Journal of Geophysical Research*, 113, D14305, doi:10.1029/2007JD009554.
- Chen, J., Avise, J., Guenther, A., Wiedinmyer, C., Salathe, E., Jackson, R.B., and Lamb, B., 2009a, Future land use and land cover influences on regional biogenic emissions and air quality in the United States, *Atmospheric Environment*, 43, 5771-5780.
- Chen, J., Avise, J., Lamb, B., Salathe, E., Mass, C., Guenther, A., Wiedinmyer, C., Lamarque, J.-F., O'Neill, S., McKenzie, D., and Larkin, N., 2009b, The effects of global changes upon regional ozone pollution in the United States, *Atmospheric Chemistry and Physics*, 9, 1125-1141.
- Chen, J., Lu, J., Avise, J.C., DaMassa, J.A., Kleeman, M.J., and Kaduwela, A.P., 2014, Seasonal modeling of PM<sub>2.5</sub> in California's San Joaquin Valley, *Atmospheric Environment*, 92, 182-190.
- Chow J.C., Watson, J.G., Lowenthal, D.H., and Magliano, K., 2005, Loss of PM<sub>2.5</sub> nitrate from filter samples in Central California, *Journal of Air & Waste Management Association*, 55, 1158-1168.
- Civerolo, K., Hogrefe, C., Zalewsky, E., Hao, W., Sistla, G., Lynn, B., Rosenzweig, C., and Kinney, P., 2010, Evaluation of an 18-year CMAQ simulation: Seasonal variations and long-term temporal changes in sulfate and nitrate, *Atmospheric Environment*, 44, 3745-3752.
- Dennis, R.L., Bhave, P., and Pinder, R.W., 2008, Observable indicators of the sensitivity of PM<sub>2.5</sub> nitrate to emission reductions – Part II: Sensitivity to errors in total ammonia and total nitrate of the CMAQ-predicted non-linear effect of SO<sub>2</sub> emission reductions, *Atmospheric Environment*, 42, 1287-1300.

Derwent, R. G., M. E. Jenkin, M. J. Pilling, W.P.L. Carter, and A. Kaduwela, 2010, Reactivity scales as comparative tools for chemical mechanisms, *Journal of the Air & Waste Management Association*, 60, 914-924.

Eder, B., Yu, S., 2006, A performance evaluation of the 2004 release of Models-3 CMAQ, *Atmospheric Environment*, 40, 4811-4824.

Emery, C., Tai, E., and Yarwood, G., 2001, Enhanced Meteorological Modeling and Performance Evaluation for Two Texas Ozone Episodes, Final report submitted to the Texas Natural Resources Conservation Commission.

Emmons, L.K., Walters, S., Hess, P.G., Lamarque, J.F., Pfister, G.G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S.L., and Kloster, S., 2010, Description and evaluation of the Model for Ozone and Related chemical Tracers, Version 4 (MOZART-4), *Geoscientific Model Development*, 3, 43-67.

Ensberg, J. J., et al., 2013, Inorganic and black carbon aerosols in the Los Angeles Basin during CalNex, *Journal of Geophysical Research - Atmosphere*, 118, 1777-1803, doi:10.1029/2012JD018136.

Fast, J.D., et al., 2014, Modeling regional aerosol and aerosol precursor variability over California and its sensitivity to emissions and long-range transport during the 2010 CalNex and CARES campaigns, *Atmospheric Chemistry Physics*, 14, 10013-10060.

Foley, K.M., Roselle, S.J., Appel, K.W., Bhawe, P.V., Pleim, J.E., Otte, T.L., Mathur, R., Sarwar, G., Young, J.O., Gilliam, R.C., Nolte, C.G., Kelly, J.T., Gilliland, A.B., and Bash, J.O., 2010, Incremental testing of the Community Multiscale Air Quality (CMAQ) modeling system version 4.7, *Geoscientific Model Development*, 3, 205-226.

Fountoukis, C. and Nenes, A., 2007, ISORROPIA II: a computationally efficient thermodynamic equilibrium model for  $K^+$ - $Ca^{2+}$ - $Mg^{2+}$ - $NH_4^+$ - $Na^+$ - $SO_4^{2-}$ - $NO_3^-$ - $Cl^-$ - $H_2O$  aerosols, *Atmospheric Chemistry Physics*, 7, 4639-4659.

Frank, N.H., 2006, Retained nitrate, hydrated sulfates, and carbonaceous mass in federal reference method fine particulate matter for six eastern U.S. cities, *Journal of Air & Waste Management Association*, 56, 500-511.

Hakami, A., Bergin, M.S., and Russell, A.G., 2004a, Ozone formation potential of organic compounds in the eastern United States: A comparison of episodes, inventories, and domain, *Environmental Science & Technology*, 38, 6748-6759.

Hakami, A., Harley, R.A., Milford, J.B., Odman, M.T., and Russell, A.G., 2004b, Regional, three-dimensional assessment of the ozone formation potential of organic compounds, *Atmospheric Environment*, 38, 121-134.

Hering, S. and Cass, G. 1999, The magnitude of bias in measurement of PM<sub>2.5</sub> arising from volatilization of particulate nitrate from Teflon filters, *Journal of Air & Waste Management Association*, 49, 725-733.

Hogrefe, C., Hao, W., Zalewsky, E.E., Ku, J.Y., Lynn, B., Rosenzweig, C., Schultz, M.G., Rast, S., Newchurch, M.J., Wang, L., Kinney, P.L., and Sistla, G., 2011, An analysis of long-term regional-scale ozone simulations over the Northeastern United States: variability and trends, *Atmospheric Chemistry and Physics*, 11, 567-582.

Hogrefe, C., Biswas, J., Lynn, B., Civerolo, K., Ku, J.Y., Rosenthal, J., Rosenzweig, C., Goldberg, R., and Kinney, P.L., 2004, Simulating regional-scale ozone climatology over the eastern United States: model evaluation results, *Atmospheric Environment*, 38, 2627-2638.

Hogrefe, C., S. T. Rao, I. G. Zurbenko, and P. S. Porter, 2000 *Interpreting Information in Time Series of Ozone Observations and Model Predictions Relevant to Regulatory Policies in the Eastern United States*. *Bull. Amer. Met. Soc.*, 81, 2083 – 2106

Hu, J., Howard, C.J., Mitloehner, F., Green, P.G., and Kleeman, M.J., 2012, Mobile source and livestock feed contributions to regional ozone formation in Central California, *Environmental Science & Technology*, 46, 2781-2789.

Hu, J., Ying, Q., Chen, J., Mahmud, A., Zhao, Z., Chen, S.H., and Kleeman, M.J., 2010, Particulate air quality model predictions using prognostic vs. diagnostic meteorology in central California, *Atmospheric Environment*, 44, 215-226.

Hu, J., Zhang, H., Chen, S., Ying, Q., Wiedinmyer, C., Vandenberghe, F., and Kleeman, M.J., 2014a, Identifying PM<sub>2.5</sub> and PM<sub>0.1</sub> sources for epidemiological studies in California, *Environmental Sciences & Technology*, 48, 4980-4990.

Hu, J., Zhang, H., Ying, Q., Chen, S.-H., Vandenberghe, F., and Kleeman, M. J., 2014b, Long-term particulate matter modeling for health effects studies in California – Part 1: Model performance on temporal and spatial variations, *Atmospheric Chemistry Physics Discussion*, 14, 20997-21036.

Huang, M., Carmichael, G.R., Adhikary, B., Spak, S.N., Kulkarni, S., Cheng, Y.F., Wei, C., Tang, Y., Parrish, D.D., Oltmans, S.J., D'Allura, A., Kaduwela, A., Cai, C., Weinheimer, A.J., Wong, M., Pierce, R.B., Al-Saadi, J.A. Streets, D.G., and Zhang, Q., 2010, Impacts of transported background ozone on California air quality during the ARCTAS-CARB period - a multi-scale modeling study, *Atmospheric Chemistry and Physics*, 10(14), 6947-6968.

Jackson, B., Chau, D., Güreş, K., and Kaduwela, A, 2006, Comparison of ozone simulations using MM5 and CALMET/MM5 hybrid meteorological fields for the July/August 2000 CCOS episode, *Atmospheric Environment*, 40, 2812-2822.

Jacob, D.J., Crawford, J.H., Maring, H., Clarke, A.D., Dibb, J.E., Emmons, L.K., Ferrare, R.A., Hostetler, C.A., Russell, P.B., Singh, H.B., Thompson, A.M., Shaw, G.E., McCauley, E., Pederson, J.R., and Fisher, J.A., 2010, The Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) Mission: Design, Execution, and First Results, *Atmospheric Chemistry and Physics*, 10(11), 5191-5212.

Jathar, S. H., Cappa, C. D., Wexler, A. S., Seinfeld, J. H., and Kleeman, M. J.: Multi-generational oxidation model to simulate secondary organic aerosol in a 3-D air quality model, *Geosci. Model Dev.*, 8, 2553-2567, doi:10.5194/gmd-8-2553-2015, 2015.

Jin L., Brown, N.J., Harley, R.A., Bao, J.W., Michelson, S.A., and Wilczak, J.M., 2010b, Seasonal versus episodic performance evaluation for an Eulerian photochemical air quality model, *Journal of Geophysical Research*, 115, D09302, doi:10.1029/2009JD012680.

Jin, L. Brown, N. and Harley, R.A. A Seasonal Perspective on Regional Air Quality in Central California, Draft Final Report, Lawrence Berkeley National Laboratory, Berkeley, CA, January, 2010a.

Jin, L., Tonse, S., Cohan, D.S., Mao, X., Harley, R.A., and Brown, N.J., 2008, Sensitivity analysis of ozone formation and transport for a central California air pollution episode, *Environmental Science Technology*, 42, 3683-3689.

Kelly, J. T., et al., 2014, Fine-scale simulation of ammonium and nitrate over the South Coast Air Basin and San Joaquin Valley of California during CalNex-2010, *Journal of Geophysical Research - Atmosphere*, 119, 3600–3614, doi:10.1002/2013JD021290.

Kelly, J.T., Avise, J., Cai, C., and Kaduwela, A., 2010b, Simulating particle size distributions over California and impact on lung deposition fraction, *Aerosol Science & Technology*, 45, 148-162.

Kelly, J.T., Bhave, P., Nolte, C.G., Shankar, U., and Foley, K.M., 2010a, Simulating emission and chemical evolution of coarse sea-salt particles in the Community Multiscale Air Quality (CMAQ) model, *Geoscientific Model Development*, 3, 257-273.

Kulkarni, S., Kaduwela, A.P., Avise, J.C., DaMassa, J.A., and Chau, D. , 2014, An extended approach to calculate the ozone relative response factors used in the attainment demonstration for the National Ambient Air Quality Standards, *Journal of the Air & Waste Management Association*, 64, 1204-1213.

Lam, Y.F. and Fu, J.S., 2009, A novel downscaling technique for the linkage of global and regional air quality modeling, *Atmospheric Chemistry and Physics*, 9, 9169-9185.

Lane, T.E., Donahue, N.M., and Pandis, S.N., 2008, Simulating secondary organic aerosol formation using the volatility basis-set approach in a chemical transport model, *Atmospheric Environment*, 42, 7439-7451.

Lee, S. H., Kim, S.W., Trainer, M., Frost, G.J., McKeen, S.A., Cooper, O.R., Flocke, F., Holloway, J.S., Neuman, J.A., Ryerson, T., Senff, C.J., Swanson, A.L., and Thompson, A.M., 2011, Modeling ozone plumes observed downwind of New York City over the North Atlantic Ocean during the ICARTT field campaign, *Atmospheric Chemistry and Physics*, 11, 7375-7397, doi:10.5194/acp-11-7375-2011.

Liang, J. and Kaduwela, A., 2005, Micro-development of CMAQ for California Regional Particulate Matter Air Quality Study, Proceedings of the 4<sup>th</sup> Annual CMAQ Models-3 User's Conference, Chapel Hill, NC.

Lin, C. J., Ho, T. C., Chu, H. W., Yang, H., Chandru, S., Krishnarajanagar, N., Chiou, P., Hopper J. R., June 2005, Sensitivity analysis of ground-level ozone concentration to emission changes in two urban regions of southeast Texas, *Journal of Environ. Manage.*, 75 315-323, <http://dx.doi.org/10.1016/j.jenvman.2004.09.012>.

Lin, M., Holloway, T., Carmichael, G.R., and Fiore, A.M., 2010, Quantifying pollution inflow and outflow over East Asia in spring with regional and global models, *Atmospheric Chemistry and Physics*, 10, 4221-4239, doi:10.5194/acp-10-4221-2010.

Lin, M., Holloway, T., Oki, T., Streets, D. G., and Richter, A., 2009, Multi-scale model analysis of boundary layer ozone over East Asia, *Atmospheric Chemistry Physics*, 9, 3277-3301, 2009.

Lin, M., Oki, T., Holloway, T., Streets, D.G., Bengtsson, M., and Kanai, S., 2008, Long-range transport of acidifying substances in East Asia - Part I: Model evaluation and sensitivity studies, *Atmospheric Environment*, 42, 5939-5955.

Livingstone, P.L., Magliano, K., Guerer, K., Allen, P.D., Zhang, K.M., Ying, Q., Jackson, B.S., Kaduwela, A., Kleeman, M., Woodhouse, L.F., Turkiewicz, K., Horowitz, L.W., Scott, K., Johnson, D., Taylor, C., O'Brien, G., DaMassa, J., Croes, B.E., Binkowski, F., and Byun, D., 2009, Simulating PM concentration during a winter episode in a subtropical valley: Sensitivity simulations and evaluation methods, *Atmospheric Environment*, 43, 5971-5977.

Lu, W., Zhong, S., Charney, J.J., Bian, X., and Liu, S., 2012, WRF simulation over complex terrain during a southern California wildfire event, *Climate and Dynamics*, 117, D05125, doi:10.1029/2011JD017004.

Mahmud, A., Hixson, M., Hu, J., Zhao, Z., Chen, S.H., and Kleeman, M.J., 2010, Climate impact on airborne particulate matter concentrations in California using seven year analysis periods, *Atmospheric Chemistry Physics*, 10, 11097-11114.



Marmur, A., Park, S.K., Mulholland, J.A., Tolbert, P.E., and Russell, A.G., 2006, Source apportionment of PM<sub>2.5</sub> in the southeastern United States using receptor and emissions-based models: Conceptual differences and implications for time-series health studies, *Atmospheric Environment*, 40, 2533-2551.

Michelson, S.A., Djalalova, I.V., and Bao, J.W., 2010, Evaluation of the Summertime Low-Level Winds Simulated by MM5 in the Central Valley of California, *Journal of Applied Meteorology and Climatology*, 49(11), 2230-2245.

Mollner, A.K., Valluvadasan, S., Feng, L., Sprague, M.K., Okumura, M., Milligan, D.B., Bloss, W.J., Sander, S.P., Martien, P.T., Harley, R.A., McCoy, A.B., and Carter, W.P.L., 2010, Rate of Gas Phase Association of Hydroxyl Radical and Nitrogen Dioxide, *Sciences*, 330, 646-649.

Morris, R.E., Koo, B., Guenther, A., Yarwood, G., McNally, D., Tesche, T.W., Tonnesen, G., Boylan, J., Brewer, P., (2006), Model sensitivity evaluation for organic carbon using two multi-pollutant air quality models that simulate regional haze in the southeastern United States, *Atmospheric Environment* 40, 4960-4972.

Napelenok, S.L., Cohan, D.S., Hu, Y., and Russell, A.G., 2006, Decoupled direct 3D sensitivity analysis for particulate matter (DDM-3D/PM), *Atmospheric Environment*, 40, 6112-6121.

O'Neill, S.M., Lamb, B.K., Chen, J., Claiborn, C., Finn, D., Otterson, S., Figueroa, C., Bowman, C., Boyer, M., Wilson, R., Arnold, J., Aalbers, S., Stocum, J., Swab, C., Stoll, M., Dubois, M., and Anderson, M., 2006, Modeling ozone and aerosol formation and transport in the Pacific Northwest with the Community Multi-Scale Air Quality (CMAQ) Modeling System, *Environmental Science Technology*, 40, 1286 – 1299.

Seinfeld J. H. and Pandis S. N. (1998) *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 1st edition, J. Wiley, New York

.Pfister, G.G., Parrish, D.D., Worden, H., Emmons, L.K., Edwards, D.P., Wiedinmyer, C., Diskin, G.S., Huey, G., Oltmans, S.J., Thouret, V., Weinheimer, A., and Wisthaler, A., 2011, Characterizing summertime chemical boundary conditions for airmasses entering the US West Coast, *Atmospheric Chemistry and Physics*, 11(4), 1769-1790.

Philips, S.B., Finkelstein, P.L., 2006, Comparison of spatial patterns of pollutant distribution with CMAQ predictions, *Atmospheric Environment*, 40, 4999-5009.

Pun, B.K., Balmori, R.T.F., Seigneur, C., 2009, Modeling wintertime particulate matter formation in central California, *Atmospheric Environment*, 43, 402-409.

Pye, H.O.T. and Pouliot, G.A., 2012, Modeling the role of alkanes, polycyclic aromatic hydrocarbons, and their oligomers in secondary organic aerosol formation, *Environmental Science & Technology*, 46, 6041-6047.

Rodriguez, M.A., Barna, M.G., Moore, T., (2009), Regional Impacts of Oil and Gas Development on Ozone Formation in the Western United States. J. Air Waste Manage. Assoc. 59, 1111-1118.

SCAQMD 8-hr ozone SIP, 2007, available at <http://www.aqmd.gov/home/library/clean-air-plans/air-quality-mgt-plan/2007-air-quality-management-plan>

SCAQMD 8-hr ozone and 24-hour PM<sub>2.5</sub> SIP, 2012, available at <http://www.aqmd.gov/home/library/clean-air-plans/air-quality-mgt-plan/final-2012-air-quality-management-plan>

Seaman, N.L., Stauffer, D.R., and Lario-Gibbs, A.M., 1995, A Multiscale Four-Dimensional Data Assimilation System Applied in the San Joaquin Valley during SARMAP. Part I: Modeling Design and Basic Performance Characteristics, Journal of Applied Meteorology 34(8), 1739-1761.

Simon, H., and Bhave, P.V., 2011, Simulating the degree of oxidation in atmospheric organic particles, Environmental Science & Technology, 46, 331-339.

Simon, H., Baker, K.R., and Phillips, S., 2012, Compilation and interpretation of photochemical model performance statistics published between 2006 and 2012, Atmospheric Environment, 61, 124-139.

SJVUAPCD 8-hour ozone Plan, 2007, available at [http://www.valleyair.org/Air\\_Quality\\_Plans/AQ\\_Final\\_Adopted\\_Ozone2007.htm](http://www.valleyair.org/Air_Quality_Plans/AQ_Final_Adopted_Ozone2007.htm)

SJVUAPCD Annual PM<sub>2.5</sub> Plan, 2008, available at [http://www.valleyair.org/Air\\_Quality\\_Plans/AQ\\_Final\\_Adopted\\_PM25\\_2008.htm](http://www.valleyair.org/Air_Quality_Plans/AQ_Final_Adopted_PM25_2008.htm)

SJVUAPCD 24-hour PM<sub>2.5</sub> Plan, 2012, available at [http://www.valleyair.org/Air\\_Quality\\_Plans/PM25Plans2012.htm](http://www.valleyair.org/Air_Quality_Plans/PM25Plans2012.htm)

SJVUAPCD 1-hour ozone Plan, 2013, available at [http://www.valleyair.org/Air\\_Quality\\_Plans/Ozone-OneHourPlan-2013.htm](http://www.valleyair.org/Air_Quality_Plans/Ozone-OneHourPlan-2013.htm)

Skamarock, W. C., J. B. Klemp, J. Dudhia, D. O. Gill, D. M. Barker, W. Wang, and J. G. Powers, 2005: A description of the Advanced Research WRF Version 2. NCAR Tech Notes-468+STR

SMAQMD 8-hour ozone Plan, 2009, available at <http://airquality.org/plans/federal/ozone/8hr1997/index.shtml>

Smyth, S.C., Jiang, W., Yin, D., Roth, H., and Giroux, E., 2006, Evaluation of CMAQ O<sub>3</sub> and PM<sub>2.5</sub> performance using Pacific 2001 measurement data, Atmospheric Environment, 40, 2735-2749.

Sokhi, R.S., Jose, R.S., Kitwiroon, N., Fragkoua, E., Perez, J.L., and Middleton, D.R., 2006, Prediction of ozone levels in London using the MM5–CMAQ modeling system. *Environmental Modeling & Software*, 21, 566–576.

Solomon, P.A. and Magliano, K.L., 1998, The 1995-Integrated Monitoring Study (IMS95) of the California Regional PM10/PM2.5 air quality study (CRPAQS): Study overview, *Atmospheric Environment*, 33(29), 4747-4756.

Stauffer, D.R., Seaman, N.L. Hunter, G.K., Leidner, S.M., Lario-Gibbs, A., and Tanrikulu, S., 2000, A field-coherence technique for meteorological field-program design for air quality studies. Part I: Description and interpretation, *Journal of Applied Meteorology*, 39(3), 297-316.

Stockwell, W. R., 2009, Peer review of the SAPRC-07 chemical mechanism of Dr. William Carter, Report to the California Air Resources Board, March 9.

Tang, Y., Carmichael, G.R., Thongboonchoo, N., Chai, T.F., Horowitz, L.W., Pierce, R.B., Al-Saadi, J.A., Pfister, G., Vukovich, J.M., Avery, M.A., Sachse, G.W., Ryerson, T.B., Holloway, J.S., Atlas, E.L., Flocke, F.M., Weber, R.J., Huey, L.G., Dibb, J.E., Streets, D.G., and Brune, W.H., 2007, Influence of lateral and top boundary conditions on regional air quality prediction: A multiscale study coupling regional and global chemical transport models, *Journal of Geophysical Research* 112, D10S18, doi:10.1029/2006JD007515.

Tang, Y.H., Lee, P., Tsidulko, M., Huang, H.C., McQueen, J.T., DiMego, G.J., Emmons, L.K., Pierce, R.B., Thompson, A.M., Lin, H.M., Kang, D.W., Tong, D., Yu, S.C., Mathur, R., Pleim, J.E., Otte, T.L., Pouliot, G., Young, J.O., Schere, K.L., Davidson, P.M., and Stajner, I., 2009, The impact of chemical lateral boundary conditions on CMAQ predictions of tropospheric ozone over the continental United States, *Environmental Fluid Mechanics*, 9, 43-58, doi:10.1007/s10652-008-9092-5.

Tanrikulu, S., Stauffer, D.R., Seaman, N.L., and Ranzieri, A.J., 2000, A Field-Coherence Technique for Meteorological Field-Program Design for Air Quality Studies. Part II: Evaluation in the San Joaquin Valley, *Journal of Applied Meteorology*, 39(3), 317-334.

Tesche, T.W., Morris, R., Tonnesen, G., McNally, D., Boylan, J., Brewer, P., 2006. CMAQ/CAMx annual 2002 performance evaluation over the eastern US. *Atmospheric Environment* 40, 4906-4919.

Tong, D.Q., and Mauzerall, D.L., 2006, Spatial variability of summertime tropospheric ozone over the continental United States: Implications of an evaluation of the CMAQ model, *Atmospheric Environment*, 40, 3041-3056.

Tonse, S.R., Brown, N.J., Harley, R.A., and Jin, L. 2008, A process-analysis based study of the ozone weekend effect, *Atmospheric Environment*, 42, 7728-7736.

U.S. EPA, 2005, Technical Support Document for the Final Clean Air Interstate Rule, Air Quality Modeling, prepared by the U.S. EPA Office of Air Quality Planning and Standards, RTP, NC.

U.S. EPA, 2007, Guidance on the Use of Models and Other Analyses for Demonstrating Attainment of Air Quality Goals for Ozone, PM<sub>2.5</sub>, and Regional Haze, EPA-454/B07-002.

U.S. EPA, 2010, Air Quality Modeling Technical Support Document: Light-Duty Vehicle Greenhouse Gas Emission Standards Final Rule, EPA Report 454/4-10-003.

U.S. EPA, (2011a), Air Quality Modeling Final Rule Technical Support Document, <http://www.epa.gov/airquality/transport/pdfs/AQModeling.pdf>, Research Triangle Park, North Carolina.

U.S. EPA, (2011b), Air Quality Modeling Technical Support Document: Final EGU NESHAP (EPA-454/R-11-009), Research Triangle Park, North Carolina.

U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM<sub>2.5</sub> and Regional Haze, available at [http://www.epa.gov/scram001/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](http://www.epa.gov/scram001/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

U.S. EPA. 1991. Guideline for Regulatory Application of the Urban Airshed Model. EPA-450/4-91-013. Found at [http://www.epa.gov/ttn/scram/guidance\\_sip.htm](http://www.epa.gov/ttn/scram/guidance_sip.htm)

UNC, 2010, Operational Guidance for the Community Multiscale Air Quality (CMAQ) Modeling System Version 4.7.1., available at [http://www.cmascenter.org/help/model\\_docs/cmaq/4.7.1/CMAQ\\_4.7.1\\_OGD\\_28june10.pdf](http://www.cmascenter.org/help/model_docs/cmaq/4.7.1/CMAQ_4.7.1_OGD_28june10.pdf).

Vijayaraghavan, K., Karamchadania, P., and Seigneur, C., 2006, Plume-in-grid modeling of summer air pollution in Central California, Atmospheric Environment, 40, 5097-5109.

Vizuette, W., Jeffries, H.E., Tesche, T.W., Olaguer, E., Couzo, E., 2011. Issues with Ozone Attainment Methodology for Houston, TX. Journal of the Air and Waste Management Association 61 (3), 238-253.

Wilczak, J. M., Djalalova, I., McKeen, S., Bianco, L., Bao, J., Grell, G, Peckham, S., Mathur, R., McQueen, J., and Lee, P., 2009, Analysis of regional meteorology and surface ozone during the TexAQS II field program and an evaluation of the NMM-CMAQ and WRF-Chem air quality models, Journal of Geophysical Research, 114, D00F14, doi:10.1029/2008JD011675.

Ying, Q., Lu, J., Allen, P., Livingstone, P., Kaduwela, A., and Kleeman, M., 2008a, Modeling air quality during the California Regional PM<sub>10</sub>/PM<sub>2.5</sub> Air Quality Study (CRPAQS) using the UCD/CIT source-oriented air quality model – Part I. Base case model results, *Atmospheric Environment*, 42, 8954-8966.

Ying, Q., Lu, J., Kaduwela, A., and Kleeman, M., 2008b, Modeling air quality during the California Regional PM<sub>10</sub>/PM<sub>2.5</sub> Air Quality Study (CRPAQS) using the UCD/CIT Source Oriented Air Quality Model - Part II. Regional source apportionment of primary airborne particulate matter, *Atmospheric Environment*, 42(39), 8967-8978.

Young, D. E., Kim, H., Parworth, C., Zhou, S., Zhang, X., Cappa, C. D., Seco, R., Kim, S., and Zhang, Q.: Influences of emission sources and meteorology on aerosol chemistry in a polluted urban environment: results from DISCOVER-AQ California, *Atmos. Chem. Phys.*, 16, 5427-5451, doi:10.5194/acp-16-5427-2016, 2016.

Zhang, H., and Ying, Q., 2011, Secondary organic aerosol formation and source apportionment in Southeast Texas, *Atmospheric Environment*, 45, 3217-3227.

Zhang, Y., Liu, P., Liu, X., Pun, B., Seigneur, C., Jacobson, M.Z., and Wang, W., 2010, Fine scale modeling of wintertime aerosol mass, number, and size distributions in Central California, *Journal of Geophysical Research*, 115, D15207, doi:10.1029/2009JD012950.

Zhang, Y., Liu, P., Queen, A., Misenis, C., Pun, B., Seigneur, C., and Wu, S.Y., 2006, A Comprehensive performance evaluation of MM5-CMAQ for the summer 1999 Southern Oxidants Study Episode, Part-II. Gas and aerosol predictions, *Atmospheric Environment*, 40, 4839-4855.

Zhang, Y., Pun, B., Wu, S.Y., Vijayaraghavan, K., and Seigneur, C., 2004, Application and Evaluation of Two Air Quality Models for Particulate Matter for a Southeastern U.S. Episode, *Journal of Air & Waste Management Association*, 54, 1478-1493.

## Appendix B-4

## Modeling Attainment Demonstration

### **Document Title:**

Modeling Attainment Demonstration – Photochemical Modeling for the 8-Hour Ozone State Implementation Plan in the Sacramento Federal Non-attainment Area (SFNA)

### **Document Description:**

This document summarizes the findings of the model attainment demonstration for the 0.075 ppm (or 75 ppb) 8-hour ozone standard in the Sacramento Federal 8-hour ozone Non-attainment Area (SFNA), which forms the scientific basis for the SFNA 2016 8-hour ozone SIP.

# **MODELING ATTAINMENT DEMONSTRATION**

## **Photochemical Modeling for the 8-Hour Ozone State Implementation Plan in the Sacramento Federal Non-attainment Area (SFNA)**

**Prepared by**  
California Air Resources Board  
Sacramento Metropolitan Air Quality Management District

**Prepared for**  
United States Environmental Protection Agency Region IX

October 12, 2016

**TABLE OF CONTENTS**

1. INTRODUCTION..... 8

2. APPROACH ..... 8

    2.1. METHODOLOGY ..... 8

    2.2. MODELING PERIOD..... 9

    2.3. BASELINE DESIGN VALUES ..... 10

    2.4. BASE, REFERENCE, AND FUTURE YEARS ..... 13

    2.5. RELATIVE RESPONSE FACTORS ..... 15

    2.6. FUTURE YEAR DESIGN VALUE CALCULATION ..... 16

3. METEOROLOGICAL MODELING..... 16

    3.1. WRF MODEL SETUP ..... 17

    3.2. WRF MODEL RESULTS AND EVALUATION ..... 20

        3.2.1 PHENOMENOLOGICAL EVALUATION ..... 26

4. EMISSIONS ..... 30

    4.1 EMISSIONS SUMMARIES ..... 30

5. OZONE MODELING ..... 31

    5.1. CMAQ MODEL SETUP ..... 31

    5.2. CMAQ MODEL EVALUATION ..... 34

        5.2.1 DIAGNOSITC EVALUATION ..... 41

    5.3. RELATIVE RESPONSE FACTORS AND FUTURE YEAR DESIGN VALUES... 45

    5.4. UNMONITORED AREA ANALYSIS ..... 47

    5.5. "BANDED" RELATIVE RESPONSE FACTORS AND FUTURE YEAR DESIGN  
    VALUES ..... 49

6. OZONE ISOPLETHS ..... 52



**LIST OF FIGURES**

Figure 1 Spatial distribution of the 8-hour ozone average DVs in the Sacramento Federal 8-hour Ozone Non-attainment Area (SFNA) for the year 2012. The circle markers and the adjacent numbers denote the location of the monitoring sites in SFNA and the corresponding value of the 2012 8-hr ozone weighted average DVs in ppb listed in Table 2. The solid grey and magenta lines denote the county and regional SFNA boundaries. The dashed black lines show the approximate regional boundaries of the Western, Central and Eastern sub-regions of SFNA. .... 13

Figure 2. WRF modeling domains (D01 36km; D02 12km; and D03 4km). .... 18

Figure 3. Meteorological monitoring sites in the model results evaluation: red markers represent sites in the valley; green markers represent sites in the mountain region. The thick black line denotes the spatial extent and regional boundary of the Sacramento Federal 8-hour ozone Non-attainment Area (SFNA) ..... 20

Figure 4. Distribution of daily mean bias (left) and mean error (right) for May-October 5 2012 (mt: Mountain; vly:Valley). Results are shown for wind speed (top), temperature (middle), and RH (bottom). .... 23

Figure 5. Spatial distribution of mean bias (left) and mean error (right) for May-October 5 2012 (mt: Mountain; vly:Valley). Results are shown for wind speed (top), temperature (middle), and RH (bottom). .... 24

Figure 6. Comparison of modeled and observed hourly wind speed (top row), 2-meter temperature (middle row), and relative humidity (bottom row). Results for Valley are shown in left column, and Mountain in right column (mt: Mountain; vly:Valley). .... 25

Figure 7 Surface wind field at 04:00 PST July 09, 2012. .... 27

Figure 8 Surface wind field at 14:00 PST July 10, 2012. .... 28

Figure 9 Surface wind field at 16:00 PST July 11, 2012. .... 29

Figure 10. Monthly average biogenic ROG emissions for 2012. .... 31

Figure 11. The CMAQ modeling domains used in this SIP modeling. The outer box of the top panel is the California statewide 12 km modeling domain, while the inner box shows the 4km modeling domain covering Central California. The shaded and gray line contours denote the gradients in topography (km). The insert on the bottom shows the zoomed-in view of the spatial extent (magenta lines), approximate regional boundaries

of the Western, Central and Eastern sub-regions (dashed black lines) and the location of ozone monitoring sites (circle markers) in the SFNA..... 33

Figure 12. Comparison of various statistical metrics from the model attainment demonstration modeling to the range of statistics from the 69 peer-reviewed studies summarized in Simon et al. (2012). (MDA denotes Maximum Daily Average) ..... 40

Figure 13. Illustrates a typical ozone isopleth plot, where each line represents ozone mixing ratio, in 10 ppb increments, as a function of initial NO<sub>x</sub> and VOC (or ROG) mixing ratio (adapted from Seinfeld and Pandis, 1998, Figure 5.15). General chemical regimes for ozone formation are shown as NO<sub>x</sub>-disbenefit (red circle), transitional (blue circle), and NO<sub>x</sub>-limited (green circle). ..... 41

Figure 14. Site-specific average weekday and weekend maximum daily average 8-hour ozone for each year from 2000 to 2014 for the Western (top), Central (middle), and Eastern (bottom) sub-regions of SFNA. The colored circle markers denote observed values while the magenta triangle, light gray diamond and dark gray square markers denote the simulated baseline 2012, future 2022 and future 2026 values respectively. Points falling below the 1:1 dashed line represent a NO<sub>x</sub>-disbenefit regime, those on the 1:1 dashed line represent a transitional regime, and those above the 1:1 dashed line represent a NO<sub>x</sub>-limited regime. .... 44

Figure 15. Spatial distribution of the future 2026 DVs based on the unmonitored area analysis in the SFNA. Color scale is in ppb of ozone. .... 49

Figure 16. The 8-hr ozone isopleth based on 2026 emission levels at the Folsom Natoma Street monitoring site located in Central SFNA..... 53

**LIST OF TABLES**

Table 1. Illustrates the data from each year that are utilized in the Design Value calculation for a specific year (DV Year), and the yearly weighting of data for the average Design Value calculation (or DV<sub>R</sub>)..... 10

Table 2. Year-specific 8-hour ozone design values for 2012, 2013, and 2014, and the average baseline design value (represented as the average of the three year-specific design values) for the monitoring sites located in the SFNA. .... 11

Table 3. Description of CMAQ model simulations. .... 14

Table 4. WRF vertical layer structure. .... 19

Table 5. WRF Physics Options. .... 19

Table 6. Meteorological site location and parameter measured. .... 21

Table 7. Hourly surface wind speed, temperature and relative humidity statistics by region for May through October 5, 2012. IOA denotes index of agreement ..... 23

Table 8. SFNA Summer Planning Emissions for 2012, 2022 and 2026 (tons/day). ..... 30

Table 9. CMAQ configuration and settings..... 34

Table 10. Daily maximum 8-hour ozone performance statistics by modeling sub-regions and entire SFNA for the 2012 ozone season (May-October 5<sup>th</sup> 2012). .... 36

Table 11. Daily maximum 1-hour ozone performance statistics by modeling sub-regions and entire SFNA for the 2012 ozone season (May-October 5<sup>th</sup> 2012). .... 37

Table 12. Hourly ozone performance statistics by modeling sub-regions and entire SFNA for the 2012 ozone season (May-October 5<sup>th</sup> 2012). Note that only statistics for the grid cell in which the monitor is located were calculated for hourly ozone..... 38

Table 13. Summary of key parameters related to the calculation of future year 2022 and 2026 8-hour ozone design values (DV). Note that final future year design values are truncated, and fractional values are shown for reference only. .... 46

Table 14. Summary of future year (2022 and 2026) design values projected using a banded RRF approach. Note that final future year design values are truncated, and fractional values are shown for reference only. .... 51

## ACRONYMS

ARB – Air Resources Board  
BCs – Boundary Conditions  
CMAQ Model – Community Multi-scale Air Quality Model  
DV – Design Value  
GEOS-5 – Goddard Earth Observing System Model, Version 5  
GMAO – Global Modeling and Assimilation Office  
ICs – Initial Conditions  
MCAB Mountain Counties Air Basin  
MOZART – Model for Ozone and Related chemical Tracers  
MDA8 – Maximum Daily Average 8-hour Ozone  
NASA – National Aeronautics and Space Administration  
NARR - North American Regional Reanalysis  
NCAR – National Center for Atmospheric Research  
NOAA - National Oceanic and Atmospheric Administration  
NO<sub>x</sub> – Oxides of nitrogen  
OFP - Ozone Forming Potential  
ROG – Reactive Organic Gases  
RH – Relative Humidity  
RRF – Relative Response Factor  
SAPRC – Statewide Air Pollution Research Center  
SIP – State Implementation Plan  
SJV – San Joaquin Valley  
SVAB – Sacramento Valley Air Basin  
SFNA – Sacramento Federal Non-attainment Area  
U.S. EPA – United States Environmental Protection Agency  
VOCs – Volatile Organic Compounds  
WRF Model – Weather and Research Forecast Model

## 1. INTRODUCTION

The purpose of this document is to summarize the findings of the model attainment demonstration for the 0.075 ppm (or 75 ppb) 8-hour ozone standard in the Sacramento Federal 8-hour ozone Non-attainment Area (SFNA), which forms the scientific basis for the SFNA 2016 8-hour ozone SIP. The 75 ppb standard was promulgated by the U.S. EPA in 2008 and became effective in 2010. Currently, the SFNA is designated as a severe ozone non-attainment area for this standard and is mandated to demonstrate attainment of the standard by 2026.

Findings from the model attainment demonstration are summarized in terms of three sub-regions within the SFNA: 1) Western SFNA (Yolo, Solano and southwest portion of Sacramento counties), 2) Central SFNA (Most of Sacramento and western portion of Placer counties), and 3) Eastern SFNA (Placer and El Dorado counties). These three sub-regions are characterized by distinct features in terms of geography, meteorology, and air quality as described in Section 2 of the Photochemical Modeling Protocol Appendix. The general approach utilized in the attainment demonstration is described in Section 2, while the remaining sections discuss the meteorological modeling (Section 3), the emissions inventory (Section 4), and the photochemical modeling and results (Sections 5 and 6). A more detailed description of the modeling and development of the model-ready emissions inventory is presented in the Photochemical Modeling Protocol Appendix.

## 2. APPROACH

This section describes the Air Resources Board's (ARB's) procedures, based on U.S. EPA guidance<sup>1</sup>, for projecting ozone Design Values (DVs) to the future using model output and a Relative Response Factor (RRF) approach in order to show future year attainment of the 0.075 ppm 8-hour ozone standard.

### 2.1. METHODOLOGY

The U.S. EPA modeling guidance<sup>1</sup> outlines the approach for utilizing models to predict future attainment of the 0.075 ppm 8-hour ozone standard. Consistent with the previous modeling guidance<sup>2</sup>, which was utilized in the most recent 8-hour ozone SIPs in

---

<sup>1</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM<sub>2.5</sub> and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

<sup>2</sup> U.S. EPA, 2007. Guidance on the Use of Models and Other Analyses for Demonstrating Attainment of Air Quality Goals for Ozone, PM<sub>2.5</sub>, and Regional Haze. EPA-454/B07-002, 2007, available at <https://www.epa.gov/ttn/scram/guidance/guide/final-03-pm-rh-guidance.pdf>

California's Central Valley, the 2009 Sacramento SIP<sup>1</sup> and the 2007 San Joaquin Valley (SJV) SIP<sup>2</sup> for the 0.08 ppm 8-hour ozone standard, the current guidance recommends utilizing modeling in a relative sense. A brief summary of how models are applied in the attainment demonstration, as prescribed by U.S. EPA modeling guidance (U.S. EPA, 2014<sup>3</sup>), is provided below. A more detailed description of the methodology is provided below and in subsequent sections is provided in the Photochemical Modeling Protocol Appendix.

## 2.2. MODELING PERIOD

Based on analysis of the conduciveness of recent years' meteorological conditions leading to elevated ozone, as well as the availability of the most detailed emissions inventory, the year 2012 was selected for both baseline modeling and design value calculation in the model attainment test. These baseline design value mixing ratios serve as the anchor point for projecting future year design values.

The severe non-attainment designation for the SFNA requires that attainment of the 2008 8-hour ozone standard be demonstrated by 2026. Therefore, 2026 was the future year modeled in this attainment demonstration. An additional future year 2022 was also modeled to assess progress toward the stipulated attainment deadline (2026).

The revised U.S. EPA modeling guidance<sup>3</sup> requires that the 8-hour ozone model attainment demonstration utilize the top ten modeled days when projecting design values to the future. Recent ozone SIP modeling applications in California's Central Valley<sup>4,5</sup>, which encompassed both the SFNA and SJV, have generally simulated the entire ozone season (May – September) as the peak ozone mixing ratios tend to occur between June and September. However, in 2012, the Sacramento region experienced a period of elevated ozone from September 30 through October 4 (see ARB's Air

---

<sup>1</sup> 2009 Sacramento Regional 8-Hour Ozone Attainment and Reasonable Further Progress Plan, available at [http://www.airquality.org/ProgramCoordination/Documents/4\)%202013%20SIP%20Revision%20Report%201997%20Std.pdf](http://www.airquality.org/ProgramCoordination/Documents/4)%202013%20SIP%20Revision%20Report%201997%20Std.pdf)

<sup>2</sup> 2007 Plan for the 1997 8-Hour Ozone Standard available at [http://www.valleyair.org/Air\\_Quality\\_Plans/AQ\\_Final\\_Adopted\\_Ozone2007.htm](http://www.valleyair.org/Air_Quality_Plans/AQ_Final_Adopted_Ozone2007.htm)

<sup>3</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM2.5 and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

<sup>4</sup> 2016 Plan for the 2008 8-Hour Ozone Standard available at [http://www.valleyair.org/Air\\_Quality\\_Plans/Ozone-Plan-2016.htm](http://www.valleyair.org/Air_Quality_Plans/Ozone-Plan-2016.htm)

<sup>5</sup> 2013 Plan for the Revoked 1-Hour Ozone Standard available at [http://www.valleyair.org/Air\\_Quality\\_Plans/Ozone-OneHourPlan-2013.htm](http://www.valleyair.org/Air_Quality_Plans/Ozone-OneHourPlan-2013.htm)

Quality and Meteorological Information System<sup>1</sup> database). Consequently, the modeling period utilized in the SFNA SIP attainment demonstration was extended to include this period, and an ozone season from May – October 5<sup>th</sup> was modeled for 2012, 2022 and 2026 to ensure that all of the top ozone days were included in the SFNA simulations.

### 2.3. BASELINE DESIGN VALUES

Specifying the baseline design value is a key consideration in the model attainment test, since this value is projected forward and used to test for future attainment at each site. The starting point for the attainment demonstration is with the observational based design value (DV), which represents the three-year average of the annual 4<sup>th</sup> highest 8-hour ozone mixing ratio observed at a specific monitor for the year in consideration. For example, a DV for 2012 would represent the average of the 4<sup>th</sup> highest 8-hour ozone mixing ratio from 2010, 2011, and 2012.

The U.S. EPA recommends using an average of three DVs that straddle the baseline year in order to better account for the year-to-year variability inherent in meteorology. Since 2012 was chosen as the base year for projecting DVs to the future, site-specific DVs were calculated for the three three-year periods ending in 2012, 2013, and 2014 and then these three DVs were averaged. This average DV is called a weighted DV (in the context of this SIP, the weighted DV will also be referred to as the reference year DV or DV<sub>R</sub>). Table 1 illustrates the observational data from each year that goes into the calculation of average DV at a particular monitoring site.

Table 1. Illustrates the data from each year that are utilized in the Design Value calculation for a specific year (DV Year), and the yearly weighting of data for the average Design Value calculation (or DV<sub>R</sub>).

DV Year	Years Averaged for the Design Value (4 <sup>th</sup> highest observed 8-hr O <sub>3</sub> )				
2012	2010	2011	2012		
2013		2011	2012	2013	
2014			2012	2013	2014
Yearly Weightings for the Average Design Value Calculation					
2012-2014 Average	$DV_R = \frac{8hrO3_{2010} + (2)8hrO3_{2011} + (3)8hrO3_{2012} + (2)8hrO3_{2013} + 8hrO3_{2014}}{9}$				

<sup>1</sup>ARB's AQMIS database is available at [www.arb.ca.gov/airqualitytoday/](http://www.arb.ca.gov/airqualitytoday/)



Table 2. Year-specific 8-hour ozone design values for 2012, 2013, and 2014, and the average baseline design value (represented as the average of the three year-specific design values) for the monitoring sites located in the SFNA.

Sub-region	Site (County, Air Basin)	8-hr Ozone Design Value (ppb)			
		2012	2013	2014	2012-2014 Average
Eastern SFNA	Placerville-Gold Nugget Way (El Dorado, MCAB <sup>1</sup> )	81	82	84	82.3
	Cool-Hwy193 (El Dorado, MCAB)	83	81	80	81.3
	Auburn - Atwood Rd (Placer, SVAB <sup>1</sup> )	80	79	78	79.0
	Colfax-City Hall (Placer, MCAB)	75	73	73	73.7
	Echo Summit (El Dorado, MCAB)	69	69	69	69.0
Central SFNA	Folsom-Natoma Street (Sacramento, SVAB)	95	90	85	90.0
	Sloughhouse (Sacramento, SVAB)	88	84	80	84.0
	Roseville-N Sunrise Ave (Placer, SVAB)	85	81	81	82.3
	Sacramento-Del Paso Manor (Sacramento, SVAB)	78	77	77	77.3
	North Highlands-Blackfoot Way (Sacramento, SVAB)	77	76	75	76.0
	Sacramento - 1309 T Street (Sacramento, SVAB)	71	70	69	70.0
	Sacramento-Goldenland Court (Sacramento, SVAB)	69	70	71	70.0
Western SFNA	Elk Grove - Bruceville Road (Sacramento, SVAB)	74	71	70	71.7
	Woodland-Gibson Road (Yolo, SVAB)	69	69	68	68.7
	Vacaville-Ulatis Drive (Solano, SVAB)	69	67	66	67.3
	Davis-UCD Campus (Yolo, SVAB)	70	66	64	66.7

<sup>1</sup> SVAB and MCAB denote the Sacramento Valley Air Basin and Mountain Counties Air Basin respectively.

Table 2 lists the design values for the sites within the three major sub-regions of the SFNA that are used in this model attainment demonstration. Note that the DVs are listed in descending order for sites within each sub-region. The Folsom – Natoma Street monitor (highlighted in black bold text), and located in Sacramento county within the Central sub-region, is the SFNA’s design site (i.e. site with the highest average DV in the SFNA) with an average DV of 90 ppb. The Placerville monitoring site, located in El Dorado county, is the design site for the eastern SFNA sub-region with an average DV of 82.3 ppb. All the monitoring sites in the western SFNA have average DVs that are below the 75 ppb standard and are already in attainment of the 2008 standard.

Figure 1 shows the spatial distribution of the baseline DVs in the SFNA. The central and eastern portions of the SFNA tend to have higher baseline DVs, and that exceed the 75 ppb standard at many sites. In contrast, baseline DVs are considerably lower, and below the 75 ppb standard, at sites located in the upwind western SFNA and at sites far downwind near the eastern edge of the SFNA. The spatial heterogeneity seen in the baseline DVs is consistent with the general characteristics of Sacramento region’s ozone plume production and evolution, which has been described as a Lagrangian air parcel that produces peak ozone levels a few kilometers downwind of the urban city center (LaFranchi et. al., 2011<sup>1</sup> and the references therein). Due to prevailing northeast wind flow patterns in this region (U.S. EPA, 2012<sup>2</sup>), the ozone plume is diluted as it migrates farther away from the urban core and downwind into the Sierra foothills (located to the east/northeast). The transport of ozone precursor emissions from the urban Sacramento area dominates ozone production in the downwind Sierra foothills, where ozone levels are heavily dependent upon the proximity to the upwind urban source. Further details on the regional topography, flow patterns and conceptual model for ozone formation in the SFNA region can be found in the modeling protocol appendix.

---

<sup>1</sup> LaFranchi, B. W., Goldstein, A. H., and Cohen, R. C.: Observations of the temperature dependent response of ozone to NO<sub>x</sub> reductions in the Sacramento, CA urban plume, *Atmos. Chem. Phys.*, 11, 6945-6960, doi:10.5194/acp-11-6945-2011, 2011

<sup>2</sup> U.S. EPA, (2012) 2008 Ground-Level Ozone Standards - Final Designations  
[https://www3.epa.gov/region9/air/ozone/pdf/R9\\_CA\\_Sacramento\\_FINAL.pdf](https://www3.epa.gov/region9/air/ozone/pdf/R9_CA_Sacramento_FINAL.pdf)




Figure 1. Spatial distribution of the 8-hour ozone baseline DVs in the Sacramento Federal 8-hour Ozone Non-attainment Area (SFNA), where the baseline DV is the average of the 2012, 2013, and 2014 DVs. Circles denote the location of each monitoring site while the baseline DV for each site is shown next to the site name in parenthesis (see Table 2). Solid grey and magenta lines denote the county and regional SFNA boundaries, while dashed black lines show the approximate regional boundaries of the Western, Central and Eastern sub-regions of the SFNA.

#### 2.4. BASE, REFERENCE, AND FUTURE YEARS

The model attainment demonstration consists of the following three primary model simulations, which all utilized the same model inputs, including meteorology, chemical boundary conditions, and biogenic emissions. The only difference between the simulations was in the year represented by the anthropogenic emissions (2012, 2022 or 2026) and certain day-specific emissions.

##### 1. Base Year (or Base Case) Simulation

The base year simulation for 2012 was used to assess model performance and includes as much day-specific detail as possible in the emissions inventory such as hourly adjustments to the motor vehicle and biogenic inventories based on observed local meteorological conditions, known wildfire and agricultural burning

events, and exceptional events like the Chevron refinery fire in the Bay Area, which occurred over 6 days from August 19-24, 2012.

**2. Reference (or Baseline) Year Simulation**

The reference year simulation was identical to the base year simulation, except that certain emissions events which are either random and/or cannot be projected to the future were removed from the emissions inventory. For the 2012 reference year modeling there are two categories/emissions sources that were excluded: 1) wildfires, which are difficult to predict in the future and can influence the model response to anthropogenic emissions reductions in regions with large fires, and 2) the Chevron refinery fire mentioned above.

**3. Future Year Simulation**

The future year simulation is identical to the reference year simulation, except that projected future year (2022 and 2026) anthropogenic emission levels were used rather than the 2012 emission levels. All other model inputs (e.g., meteorology, chemical boundary conditions, biogenic emissions, and calendar for day-of-week specifications in the inventory) are the same as those used in the reference year simulation.

To summarize (Table 3), the base year 2012 simulation was used for evaluating model performance, while the reference (or baseline) 2012 and future year 2022 and 2026 simulations were used to project the baseline DVs to the future as described in the Photochemical Modeling Protocol Appendix and in subsequent sections of this document.

Table 3. Description of CMAQ model simulations.

Simulation	Anthropogenic Emissions	Biogenic Emissions	Meteorology	Chemical Boundary Conditions
Base year (2012)	2012 w/ wildfires and Chevron refinery fire	2012 MEGAN	2012 WRF	2012 MOZART
Reference year (2012)	2012 w/o wildfires and w/o Chevron refinery fire	2012 MEGAN	2012 WRF	2012 MOZART
Future year (2022)	2022 w/o wildfires and w/o Chevron refinery fire	2012 MEGAN	2012 WRF	2012 MOZART
Future year (2026)	2026 w/o wildfires and w/o Chevron refinery fire	2012 MEGAN	2012 WRF	2012 MOZART

## 2.5. RELATIVE RESPONSE FACTORS

As part of the model attainment demonstration, the fractional changes in ozone mixing ratios between the model reference year and model future year were calculated at each of the monitors. These ratios, called “relative response factors” (RRFs), were calculated based on the ratio of future year modeled maximum daily average 8-hour (MDA8) ozone to modeled reference year MDA8 ozone (Equation 1).

$$\text{RRF} = \frac{\text{average MDA8 ozone}_{\text{future}}}{\text{average MDA8 ozone}_{\text{reference}}} \quad (1)$$

The MDA8 values, used in calculating the RRF, were based on the maximum simulated ozone within a 3x3 array of cells with the grid cells containing the monitor located at the center of the array<sup>1</sup>. The future and reference year ozone values used in the RRF calculations were paired in space and time (i.e., using the future year MDA8 ozone for the same modeled day and at the same grid cell where the MDA8 ozone for the reference year is located within the 3x3 array of cells). The modeled days utilized in the RRF calculation were selected based on the following U.S. EPA recommended criteria<sup>1</sup>.

- Begin with days that have simulated baseline MDA8 > 60 ppb and calculate RRFs based on the top 10 high ozone days.
- If there are fewer than 10 days with MDA8 > 60 ppb then all days > 60 ppb are used in the RRF calculation, as long as there are at least 5 days used in the calculation.
- If there are fewer than 5 days > 60 ppb, an RRF is not calculated at that monitor.
- Restrict the simulated days used in the RRF calculation by only including days with reference MDA8 within +/- 20% of the observed value at the monitor. This ensures that only modeled days which are consistent with the observed ozone levels are used in the RRF calculation.

RRFs were calculated for all monitors within the SFNA following the procedure described above, except for the Folsom monitor. The Folsom monitor is located adjacent to Folsom Lake, such that the northeast corner grid cell of the 3x3 array of grid cells centered at the monitor overlays a portion of Folsom Lake. High ozone mixing ratios are frequently observed over lake surfaces due to a shallow convective boundary layer. Recent studies have shown that simulated ozone over lake surfaces tend to

---

<sup>1</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM2.5 and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

exhibit a higher positive bias than over the surrounding land<sup>1</sup>, which may be due to a simulated boundary layer that is too low over lake surfaces. However, these high biases do not appear to propagate strongly over the inland areas that are located in the vicinity of a lake. Because of this high bias in ozone over lake surfaces, data from the grid cell over Folsom Lake will not be used when calculating the daily maximum 8-hour ozone in the 3x3 array of grid cells centered over the Folsom monitor (i.e., the daily maximum will be calculated from 8 grid cells rather than the standard 9 grid cells).

## 2.6. FUTURE YEAR DESIGN VALUE CALCULATION

Future year design values for each site were calculated by multiplying the corresponding baseline design value (Table 2) by the site-specific RRF (Equation 2).

$$DV_F = DV_R \times RRF \quad (2)$$

where,

$DV_F$  = the future year design value,

$DV_R$  = the reference year design value (from Table 2), and

RRF = the site specific RRF from Equation 1

Future year design values from the model attainment demonstration are discussed in Section 5.3.

## 3. METEOROLOGICAL MODELING

California's proximity to the ocean, complex terrain, and diverse climate represent a unique challenge for developing meteorological fields that adequately represent the synoptic and mesoscale features of the regional meteorology. In summertime, the majority of the storm tracks are far away to the north of the state and a semi-permanent Pacific high typically sits off the California coast. Interactions between this eastern Pacific subtropical high pressure system and the thermal low pressure further inland over the Central Valley or South Coast lead to conditions conducive to pollution buildup

---

<sup>1</sup> Cleary, P. A., Fuhrman, N., Schulz, L., Schafer, J., Fillingham, J., Bootsma, H., McQueen, J., Tang, Y., Langel, T., McKeen, S., Williams, E. J., and Brown, S. S.: Ozone distributions over southern Lake Michigan: comparisons between ferry-based observations, shoreline-based DOAS observations and model forecasts, *Atmos. Chem. Phys.*, 15, 5109-5122, doi:10.5194/acp-15-5109-2015, 2015.

(Fosberg and Schroeder, 1966<sup>1</sup>; Bao et al., 2008<sup>2</sup>). In the past, the ARB has utilized both prognostic and diagnostic meteorological models, as well as hybrid approaches in an effort to develop meteorological fields for use in air quality modeling that most accurately represent the meteorological processes that are important to air quality (e.g., Jackson et al., 2006<sup>3</sup>). In this work, the state-of-the-science Weather and Research Forecasting (WRF) prognostic model (Skamarock et al., 2005<sup>4</sup>) version 3.6 was utilized to develop the meteorological fields used in the subsequent photochemical model simulations.

### 3.1. WRF MODEL SETUP

The WRF meteorological modeling domain consisted of three nested Lambert projection grids of 36-km (D01), 12-km (D02), and 4-km (D03) horizontal grid spacing (Figure 2). WRF was run simultaneously for the three nested domains with two-way feedback between the parent and the nested grids. The D01 and D02 grids were used to resolve the larger scale synoptic weather systems, while the D03 grid resolved the finer details of the atmospheric conditions and was used to drive the air quality model simulations. All three domains utilized 30 vertical sigma layers (defined in Table 4), with the major physics options for each domain listed in Table 5.

Initial and boundary conditions (IC/BCs) for the WRF modeling were based on the 32-km horizontal resolution North American Regional Reanalysis (NARR) data that are archived at the National Center for Atmospheric Research (NCAR). Boundary conditions to WRF were updated at 6-hour intervals for the 36-km grid (D01). In addition, surface and upper air observations obtained from NCAR were used to further refine the analysis data that were used to generate the IC/BCs. Analysis nudging was employed in the outer 36-km grid (D01) to ensure that the simulated meteorological fields were constrained and did not deviate from the observed meteorology. No

---

<sup>1</sup> Fosberg, M.A., Schroeder, M.J., Marine air penetration in Central California, *Journal of Applied Meteorology*, 5, 573-589, 1966.

<sup>2</sup> Bao, J.W., Michelson, S.A., Persson, P.O.G., Djalalova, I.V., Wilczak, J.M., Observed and WRF-simulated low-level winds in a high-ozone episode during the Central California ozone study, *Journal of Applied Meteorology and Climatology*, 47, 2372-2394, 2008.

<sup>3</sup> Jackson, B.S., Chau, D., Gurer, K., Kaduwela, A.: Comparison of ozone simulations using MM5 and CALMET/MM5 hybrid meteorological fields for the July/August 2000 CCOS episode, *Atmos. Environ.*, 40, 2812-2822, 2006.

<sup>4</sup> Skamarock, W. C., J. B. Klemp, J. Dudhia, D. O. Gill, D. M. Barker, W. Wang, and J. G. Powers, 2005: A description of the Advanced Research WRF Version 2. NCAR Tech Notes-468+STR

nudging was used on the two inner domains to allow model physics to work fully without externally imposed forcing (Rogers et al., 2013<sup>1</sup>).




Figure 2. WRF modeling domains (D01 36km; D02 12km; and D03 4km).

<sup>1</sup> Rogers, R.E., Deng, A., Stauffer, D. Gaudet, B.J., Jia, Y., Soong, S.-T., Tanrikulu, S., Application of the Weather Research and Forecasting model for air quality modeling in the San Francisco Bay area, *Journal of Applied Meteorology and Climatology*, 52, 1953-1973, 2013.



Table 4. WRF vertical layer structure.

Layer Number	Height (m)	Layer Thickness (m)	Layer Number	Height (m)	Layer Thickness (m)
30	16082	1192	14	1859	334
29	14890	1134	13	1525	279
28	13756	1081	12	1246	233
27	12675	1032	11	1013	194
26	11643	996	10	819	162
25	10647	970	9	657	135
24	9677	959	8	522	113
23	8719	961	7	409	94
22	7757	978	6	315	79
21	6779	993	5	236	66
20	5786	967	4	170	55
19	4819	815	3	115	46
18	4004	685	2	69	38
17	3319	575	1	31	31
16	2744	482	0	0	0
15	2262	403			

Note: Shaded layers denote the subset of vertical layers used in the CMAQ photochemical model simulations.

Table 5. WRF Physics Options.

Physics Option	Domain		
	D01 (36 km)	D02 (12 km)	D03 (4 km)
Microphysics	WSM 6-class graupel scheme	WSM 6-class graupel scheme	WSM 6-class graupel scheme
Longwave radiation	RRTM	RRTM	RRTM
Shortwave radiation	Dudhia scheme	Dudhia scheme	Dudhia scheme
Surface layer	Revised MM5 Monin-Obukhov	Revised MM5 Monin-Obukhov	Revised MM5 Monin-Obukhov
Land surface	Pleim-Xiu LSM	Pleim-Xiu LSM	Pleim-Xiu LSM
Planetary Boundary Layer	YSU	YSU	YSU
Cumulus Parameterization	Kain-Fritsch scheme	Kain-Fritsch scheme	None

### 3.2. WRF MODEL RESULTS AND EVALUATION

Simulated surface wind speed, temperature, and relative humidity from the 4 km domain were validated against hourly observations at 31 surface stations (Figure 3).

Considering the geographical and meteorological differences, the area covered by these sites was divided into two regions: the lower elevation (Valley) and higher elevation mountain (Mountain) areas. Among the 31 surface sites used in this analysis, 17 of them are located in the valley zone with the remaining 14 sites located in the mountain region.

The observational data for the surface stations were obtained from the ARB archived meteorological database available at <http://www.arb.ca.gov/aqmis2/aqmis2.php>. Table 6 lists the monitoring stations and the meteorological parameters that are measured at each station, including wind speed and direction (wind), temperature (T) and relative humidity (RH). Figure 3 shows the location of each of these sites with the red and green circle markers denoting the sites in the valley and mountain sub-regions while the black lines denote the regional boundary of the (SFNA).




Figure 3. Meteorological monitoring sites in the model results evaluation: red markers represent sites in the valley; green markers represent sites in the mountain region. The thick black line denotes the spatial extent and regional boundary of the Sacramento Federal 8-hour ozone Non-attainment Area (SFNA)

Table 6. Meteorological site location and parameter measured.

Site	Site ID	Site Name	Region	Parameter Measured
1	2958	Yuba City-Almond Street	Valley	Wind, T
2	3290	Lincoln (RAWS)	Valley	Wind, T, RH
3	2956	Roseville-N Sunrise Blvd	Valley	Wind, T, RH
4	3397	Brooks	Valley	Wind, T, RH
5	5833	Esparto	Valley	T, RH
6	3187	Folsom-Natoma Street	Valley	Wind, T, RH
7	5776	Fair Oaks #2	Valley	T, RH
8	2731	Sacramento-Del Paso Manor	Valley	Wind, T, RH
9	5799	Bryte	Valley	R, RH
10	3011	Sacramento-T Street	Valley	Wind, T, RH
11	5710	Davis #2	Valley	T, RH
12	2143	Davis-UCD Campus	Valley	Wind, T
13	5784	Winters	Valley	T, RH
14	3209	Sloughhouse	Valley	Wind
15	5767	Dixon	Valley	T, RH
16	2977	Elk Grove-Bruceville Road	Valley	Wind, T, RH
17	5785	Twitchell Island	Valley	T, RH
18	5880	Duncan #2	Mountain	Wind, T, RH
19	3564	Foresthill #2	Mountain	Wind, T, RH
20	3288	Hell Hole	Mountain	Wind, T, RH
21	2948	South Lake Tahoe-Sandy Way	Mountain	Wind, T
22	3289	Bald Mountain Location	Mountain	Wind
23	3196	Cool-Highway 193	Mountain	Wind, T
24	5832	Auburn #3	Mountain	T, RH
25	3454	Meyers	Mountain	Wind, T, RH
26	3291	Pilot Hill Station	Mountain	Wind, T, RH
27	3487	Echo Summit	Mountain	Wind, T
28	5714	Camino #2	Mountain	T, RH
29	3292	Owens Camp	Mountain	Wind, T, RH
30	3017	Placerville-Gold Nugget Way	Mountain	Wind, T
31	3293	Ben Bolt	Mountain	Wind, T, RH

Several quantitative performance metrics were used to compare hourly surface observations and modeled estimates: mean bias (MB), mean error (ME) and index of agreement (IOA) based on the recommendations from Simon et al. (2012)<sup>1</sup>. A summary of these statistics by performance region is shown in Table 7. The distribution of daily mean bias and mean error are shown in Figure 4. The spatial distributions of the mean bias and mean error of modeled surface wind, temperature and relative humidity are shown in Figure 5, while observed vs. modeled scatter plots are shown in Figure 6. Wind Speed biases are positive in each of the two regions. The average bias for the valley sites is 0.69 m/s. The model generally over-predicted the wind speed for the mountain sites, with an average positive bias of 1.28 m/s. This is also evident in the wind speed scatter plot (top right panel of Figure 6). Temperature bias is relatively small in the valley with a bias of -0.02 °K, and higher in the mountain areas (-1.22 °K). Temperature generally shows good agreement between the observations and simulation with IOA above 0.90. Relative humidity biases range from 1.03% to 7.96%. These results are comparable to other recent WRF modeling efforts in California investigating ozone formation in Central California (e.g., Hu et al., 2012<sup>2</sup>) and modeling analysis for the CalNex and CARES field studies (e.g., Fast et al., 2014<sup>3</sup>; Baker et al., 2013<sup>4</sup>; Kelly et al., 2014<sup>5</sup>; Angevine et al., 2012<sup>6</sup>). Detailed hourly time-series of surface

---

<sup>1</sup> Simon, H., Baker, K. R., and Phillips, S.: Compilation and interpretation of photochemical model performance statistics published between 2006 and 2012, *Atmospheric Environment*, 61, 124-139, 2012

<sup>2</sup> Hu, J., Howard, C. J., Mitloehner, F., Green, P. G., and Kleeman, M. J.: Mobile Source and Livestock Feed Contributions to Regional Ozone Formation in Central California, *Environmental Science & Technology*, 46, 2781-2789, 2012.

<sup>3</sup>Fast, J. D., Gustafson Jr, W. I., Berg, L. K., Shaw, W. J., Pekour, M., Shrivastava, M., Barnard, J. C., Ferrare, R. A., Hostetler, C. A., Hair, J. A., Erickson, M., Jobson, B. T., Flowers, B., Dubey, M. K., Springston, S., Pierce, R. B., Dolislager, L., Pederson, J., and Zaveri, R. A.: Transport and mixing patterns over Central California during the carbonaceous aerosol and radiative effects study (CARES), *Atmos. Chem. Phys.*, 12, 1759-1783, 2012, doi:10.5194/acp-12-1759-2012.

<sup>4</sup>Baker, K. R., Misenis, C., Obland, M. D., Ferrare, R. A., Scarino, A. J., and Kelly, J. T.: Evaluation of surface and upper air fine scale WRF meteorological modeling of the May and June 2010 CalNex period in California, *Atmos. Environ.*, 80, 299-309, 2013.

<sup>5</sup> Kelly, J. T., Baker, K. R., Nowak, J. B., Murphy, J. G., Milos, Z. M., VandenBoer, T. C., Ellis, R. A., Neuman, J. A., Weber, R. J., Roberts, J. M., Veres, P. R., de Gouw, J. A., Beaver, M. R., Newman, S., and Misenis, C.: Fine-scale simulation of ammonium and nitrate over the South Coast Air Basin and San Joaquin Valley of California during CalNex-2010, *J. Geophysical Research*, 119, 3600-3614, doi:10.1002/2013JD021290.

<sup>6</sup> Angevine, W. M., Eddington, L., Durkee, K., Fairall, C., Bianco, L., Brioude, J.: Meteorological model evaluation for CalNex 2010, *Monthly Weather Review*, 140, 3885-3906, 2012.

temperature, relative humidity, wind speed, and wind direction for each sub-region can be found in the supplementary material.

Table 7. Hourly surface wind speed, temperature and relative humidity statistics by region for May through October 5, 2012. IOA denotes index of agreement

Region	Observed Mean	Modeled Mean	Mean Bias	Mean Error	IOA
<b>Wind Speed (m/s)</b>					
Valley	2.11	2.80	0.69	1.21	0.67
Mountain	1.75	3.03	1.28	1.63	0.44
<b>Temperature (K)</b>					
Valley	295.48	295.46	-0.02	2.42	0.94
Mountain	292.42	291.19	-1.22	2.83	0.94
<b>Relative Humidity (%)</b>					
Valley	48.63	49.66	1.03	11.32	0.85
Mountain	40.56	48.52	7.96	15.97	0.72




Figure 4. Distribution of daily mean bias (left) and mean error (right) from May-October 5, 2012 for Mountain (mt) and Valley (vly) sites. Results are shown for wind speed (top), temperature (middle), and RH (bottom).




Figure 5. Spatial distribution of mean bias (left) and mean error (right) from May-October 5, 2012 for Mountain (mt) and Valley (vly) sites. Results are shown for wind speed (top), temperature (middle), and RH (bottom).




Figure 6. Comparison of modeled and observed hourly wind speed (top row), 2-meter temperature (middle row), and relative humidity (bottom row). Results for Valley (vly) are shown in the left column, and Mountain (mt) in the right column.

### 3.2.1 PHENOMENOLOGICAL EVALUATION

Conducting a detailed phenomenological evaluation for all modeled days can be resource intensive given that the entire ozone season was modeled. However, some insight and confidence that the model is able to reproduce the meteorological conditions leading to elevated ozone can be gained by investigating the meteorological conditions during a period of peak ozone within the Sacramento non-attainment area in more detail. Meteorological conditions that produced the highest ozone levels in the area occurred on or around July 10, 2012. The July 10<sup>th</sup> episode represents a typical ozone episode in the Sacramento area consistent with the conceptual model for ozone described in the Modeling Protocol Appendix. Surface weather analysis during the episode showed that the Sacramento area was caught between a high pressure center off the California coast and a large high pressure system over an area spanning from the Rockies to the Midwest. The surface wind distributions (Figures 7, 8, 9) indicate the model was able to capture many of the important features of the meteorological fields in this area. In the early morning of July 9 (Figure 7), the bifurcation of the delta breeze, one branch up to the Sacramento valley and one down to the San Joaquin Valley, is not as strong as during the afternoon of July 11 (Figure 9). The downslope flows on the west slope of the Sierra and east side of the Coastal Ranges created some convergence zones along the foothills. Figure 8 shows a lower valley convergence formed along the Solano-Yolo border in the afternoon of July 10 with upslope flows fully developed in the mountain areas. This is a wind pattern which occurs relatively infrequently in the area (Hayes et al., 1984<sup>1</sup>). The upslope flows are stronger than those in the afternoon of July 11, but less orderly. Overall, the modeled winds are in general agreement with the observations on both the valley floor and mountain areas during this episode. Although a phenomenological evaluation of a single episode does not necessarily mean the model performs equally well on all days, the fact that the model can adequately reproduce wind flows consistent with the ozone conceptual model, combined with reasonable performance statistics over the ozone season (Table 7), provides added confidence in the meteorological fields.

---

<sup>1</sup> Hayes, T.P., J.J. Kinney, and N.J. Wheeler 1984: California surface wind climatology. California Air Resources Board, Sacramento, CA, 180pp.






Figure 7. Surface wind field at 04:00 PST July 09, 2012.




Figure 8. Surface wind field at 14:00 PST July 10, 2012.




Figure 9. Surface wind field at 16:00 PST July 11, 2012.

#### 4. EMISSIONS

The emissions inventory used in this modeling was based on the most recent inventory submitted to the U.S. EPA, with base year 2012

(<http://www.arb.ca.gov/planning/sip/2012iv/2012iv.htm>). For a detailed description of the emissions inventory, updates to the inventory, and how it was processed from the planning totals to a gridded inventory for modeling, see the Modeling Emissions Inventory Appendix.

#### 4.1 EMISSIONS SUMMARIES

Table 8 summarizes the 2012, 2022 and 2026 SFNA anthropogenic emissions used in this work. Overall, anthropogenic NO<sub>x</sub> was projected to decrease ~45% by 2022 (from 104 tpd to 56.8 tpd) and ~55% by 2026 (from 104 tpd to 47.3 tpd) when compared to 2012 emissions levels. In contrast, anthropogenic ROG was projected to decrease ~23 % by 2022 (from 109.8 tpd to 84.7 tpd) and ~26 % by 2026 (from 109.8 tpd to 81.7 tpd).

Table 8. SFNA Summer Planning Emissions for 2012, 2022 and 2026 (tons/day).

Source Category	NO <sub>x</sub>					ROG						
	2012		2022		2026		2012		2022		2026	
	[tpd]	[tpd]	% diff <sup>#</sup>	[tpd]	% diff <sup>#</sup>	[tpd]	[tpd]	% diff <sup>#</sup>	[tpd]	% diff <sup>#</sup>	[tpd]	% diff <sup>#</sup>
Stationary	9.2	7.6	-17	7.6	-17	20.6	21.9	6	22.1	7		
Area	2.7	2.1	-22	2.1	-22	28.5	29.5	4	30.4	7		
On-Road Mobile	62	24.5	-60	17.7	-71	35	15.6	-55	13.3	-62		
Other Mobile	30.1	22.6	-25	19.9	-34	25.7	17.7	-32	15.9	-39		
Total	104	56.8	-45	47.3	-55	109.8	84.7	-23	81.7	-26		

<sup>#</sup> % diff denotes percent difference with respect to 2012 emission levels.

Monthly biogenic ROG totals for 2012 within the SFNA are shown in Figure 10 (note that the same biogenic emissions were used in 2012, 2022 and 2026 modeling).

Throughout the summer, biogenic ROG emissions ranged from ~450 tpd in May to over 900 tpd in July and August, with the difference in emissions primarily due to differences in temperature, solar radiation, and leaf area from month-to-month.




Figure 10. Monthly average biogenic ROG emissions for 2012.

## 5. OZONE MODELING

### 5.1. CMAQ MODEL SETUP

Figure 11 shows the CMAQ modeling domains used in this work. The larger domain covering all of California has a horizontal grid resolution of 12 km with 107x97 lateral grid cells for each vertical layer and extends from the Pacific Ocean in the west to Eastern Nevada in the east, and runs from the U.S.-Mexico border in the south to the California-Oregon border in the north. The smaller nested domain (dashed black line) covering the Central valley region including the San Joaquin Valley, Sacramento Valley, and Mountain Counties air basins has a finer scale 4 km grid resolution and includes 192x192 lateral grid cells. The 12 km and 4 km domains are based on a Lambert Conformal Conic projection with reference longitude at -120.5°W, reference latitude at 37°N, and two standard parallels at 30°N and 60°N, which is consistent with WRF domain settings. The 30 vertical layers from WRF were mapped onto 18 vertical layers for CMAQ extending from the surface to 100 mb such that majority of the vertical layers fall within the planetary boundary layer. This vertical layer structure is based on the

WRF sigma-pressure coordinates and the exact layer structure used can be found in Table 4.

The photochemical modeling for this attainment demonstration utilized CMAQ version 5.0.2, released by the U.S. EPA (<https://www.cmascenter.org/cmaq/>) in May 2014. The SAPRC07 mechanism was selected as the photochemical mechanism for the CMAQ simulations. Further details of the CMAQ configuration used in this work are summarized in Table 9 and in the Photochemical Modeling Protocol Appendix. The same configuration has been used for all simulations including the base, reference, and future years. CMAQ was compiled using the Intel FORTRAN compiler version 12.

The entire ozone season (May – October 5<sup>th</sup> 2012) was simulated through individual monthly simulations conducted in parallel. For each month, the CMAQ simulations included a seven day spin-up period (i.e., the last seven days of the previous month) for the outer 12 km domain, where initial conditions for the first day were set to the default initial conditions included with the CMAQ release. The 4 km inner domain simulations utilized a three day spin-up period, with initial conditions derived from output from the corresponding day of the 12 km domain simulation.

Chemical boundary conditions (BCs) for the outer 12 km domain were extracted from the global chemical transport Model for Ozone and Related chemical Tracers, version 4 (MOZART-4; Emmons et al., 2010<sup>1</sup>). The MOZART-4 data for 2012 was obtained from the National Center for Atmospheric Research (NCAR; <http://www.acom.ucar.edu/wrf-chem/mozart.shtml>) for the simulations driven by meteorological fields from the NASA GMAO GEOS-5 model. The same MOZART derived BCs for the 12 km outer domain, were used for all simulations (e.g., Base, Reference, Future, and any sensitivity simulation). The inner 4 km domain simulations utilized BCs that were based on the output from the corresponding day of the 12 km domain simulation.

---

<sup>1</sup> Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43-67, doi:10.5194/gmd-3-43-2010, 2010.




Figure 11. The CMAQ modeling domains used in this SIP modeling. The outer box of the top panel is the California statewide 12 km modeling domain, while the inner box shows the 4 km modeling domain covering Central California. The shaded and gray line contours denote the gradients in topography (km). The insert on the bottom shows the zoomed-in view of the spatial extent (magenta lines), approximate regional boundaries of the Western, Central and Eastern sub-regions (dashed black lines) and the location of ozone monitoring sites (circle markers) in the SFNA.

Table 9. CMAQ configuration and settings.

<b>Process</b>	<b>Scheme</b>
Horizontal advection	Yamo (Yamartino scheme for mass-conserving advection)
Vertical advection	WRF-based scheme for mass-conserving advection
Horizontal diffusion	Multi-scale
Vertical diffusion	ACM2 (Asymmetric Convective Model version 2)
Gas-phase chemical mechanism	SAPRC-07 gas-phase mechanism with version "C" toluene updates
Chemical solver	EBI (Euler Backward Iterative solver)
Aerosol module	Aero6 (the sixth-generation CMAQ aerosol mechanism with extensions for sea salt emissions and thermodynamics; includes a new formulation for secondary organic aerosol yields)
Cloud module	ACM_AE6 (ACM cloud processor that uses the ACM methodology to compute convective mixing with heterogeneous chemistry for AERO6)
Photolysis rate	phot_inline (calculate photolysis rates in-line using simulated aerosols and ozone concentrations)

## 5.2. CMAQ MODEL EVALUATION

Observed ozone data from the Air Quality and Meteorological Information System (AQMIS) database ([www.arb.ca.gov/airqualitytoday/](http://www.arb.ca.gov/airqualitytoday/)) was used to evaluate the accuracy of the 4 km CMAQ modeling for all ozone monitors listed in Table 2 and Figure 11. The U.S. EPA modeling guidance<sup>1</sup> recommends using model output from the grid cell in which the monitor is located in the operational evaluation of the model predictions. However, the future year design value calculations (discussed in Sections 2.5 and 2.6) are based on simulated values > 60 ppb near the monitor (i.e., the maximum simulated ozone within a 3x3 array of grid cells with the grid cell containing the monitor located at the center of the array). Hence, model performance was evaluated at each monitor by comparing observations against the simulated values using only data above the 60 ppb threshold at the monitored grid cell as well as the peak grid cell within the 3x3 grid array

<sup>1</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM<sub>2.5</sub> and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)



centered on the monitor (i.e., the 3x3 maximum). Model performance was further summarized separately for the three sub-regions in Figure 11.

As recommended by U.S. EPA<sup>1</sup>, a number of statistical metrics have been used to evaluate the model performance for ozone. These metrics include mean bias (MB), mean error (ME), mean fractional bias (MFB), mean fractional error (MFE), normalized mean bias (NMB), normalized mean error (NME), root mean square error (RMSE), and correlation coefficient ( $R^2$ ). In addition, the following plots were used in evaluating the modeling: time-series comparing predictions and observations, scatter plots for comparing the magnitude of simulated and observed mixing ratios, box plots to summarize the time series data across different regions and averaging times, as well as frequency distributions.

The model performance evaluation is presented for the entire SFNA region and also disaggregated for the three sub-regions. Performance statistics for data above 60 ppb are reported separately for different ozone metrics including 8-hour daily maximum ozone, 1-hour daily maximum ozone, and hourly ozone (all hours of the day) for the monitored grid cell as well as the 3x3 maximum.

Performance statistics for Maximum Daily Average 8-hour ozone (MDA8) are shown in Table 10. Overall, when simulated data extracted at the grid cell is used for comparison with observations, the model shows a slight negative bias in MDA8 ozone greater than 60 ppb in the Central SFNA (-2.4 ppb) and Eastern SFNA (-1.3 ppb), while a very small positive bias (0.4) is seen in the Western SFNA. However, when the 3x3 maximum is used, the model shows a slight positive bias in MDA8 in the Western (1.9 ppb) and Eastern (0.1) SFNA, with a slight negative bias in Central SFNA (-0.1 ppb). Mean error shows a consistent trend with the error increasing slightly by 0.2 ppb (from 7.2 ppb to 7.4 ppb) for the entire SFNA when the 3x3 maximum is considered. Similar statistics for daily maximum 1-hour ozone and hourly ozone can be found in Table 11 and Table 12, respectively.

---

<sup>1</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM2.5 and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

Table 10. Daily maximum 8-hour ozone performance statistics by modeling sub-regions and entire SFNA for the 2012 ozone season (May-October 5<sup>th</sup> 2012).

<b>Daily Maximum 8-hour ozone &gt; 60 ppb with simulated data extracted at grid cell where the monitor is located</b>				
<b>Parameter</b>	<b>Western SFNA</b>	<b>Central SFNA</b>	<b>Eastern SFNA</b>	<b>Entire SFNA</b>
Number of data points	64	244	227	535
Mean obs (ppb)	67.2	72.1	69.4	70.4
Standard Deviation obs (ppb)	6.6	9.5	6.9	8.3
Mean Bias (ppb)	0.4	-2.4	-1.3	-1.6
Mean Error (ppb)	5.1	7.9	7.1	7.2
RMSE (ppb)	6.6	10	9.1	9.3
Normalized Mean Bias (%)	0.6	-3.3	-1.9	-2.3
Normal Mean Error (%)	7.6	10.9	10.3	10.3
R-squared	0.15	0.14	0.03	0.1
Index of Agreement	0.61	0.62	0.49	0.58

<b>Daily Maximum 8-hour ozone &gt; 60 ppb with simulated data extracted from the 3x3 grid cell array maximum centered at the monitor</b>				
<b>Parameter</b>	<b>Western SFNA</b>	<b>Central SFNA</b>	<b>Eastern SFNA</b>	<b>Entire SFNA</b>
Number of data points	69	275	259	603
Mean obs (ppb)	67.3	71.7	69.2	70.1
Standard Deviation obs (ppb)	6.4	9.3	6.9	8.2
Mean Bias (ppb)	1.9	-0.1	0.1	0.2
Mean Error (ppb)	6.1	7.9	7.3	7.4
RMSE (ppb)	7.4	10.2	9.3	9.5
Normalized Mean Bias (%)	2.7	-0.2	0.2	0.3
Normal Mean Error (%)	9	11	10.5	10.6
R-squared	0.11	0.14	0.04	0.11
Index of Agreement	0.58	0.63	0.51	0.6

Table 11. Daily maximum 1-hour ozone performance statistics by modeling sub-regions and entire SFNA for the 2012 ozone season (May-October 5<sup>th</sup> 2012).

<b>Daily Maximum 1-hour ozone &gt; 60 ppb with simulated data extracted at grid cell where the monitor is located</b>				
<b>Parameter</b>	<b>Western SFNA</b>	<b>Central SFNA</b>	<b>Eastern SFNA</b>	<b>Entire SFNA</b>
Number of data points	189	455	361	1005
Mean obs (ppb)	71	77	73.4	74.6
Standard Deviation obs (ppb)	8.3	12.4	9.5	11
Mean Bias (ppb)	0.7	-2.8	-0.8	-1.4
Mean Error (ppb)	6.8	9.5	8.3	8.6
RMSE (ppb)	8.9	12.5	10.8	11.3
Normalized Mean Bias (%)	1	-3.6	-1.1	-1.9
Normal Mean Error (%)	9.6	12.4	11.4	11.5
R-squared	0.16	0.22	0.15	0.2
Index of Agreement	0.64	0.68	0.63	0.67

<b>Daily Maximum 1-hour ozone &gt; 60 ppb with simulated data extracted from the 3x3 grid cell array maximum centered at the monitor</b>				
<b>Parameter</b>	<b>Western SFNA</b>	<b>Central SFNA</b>	<b>Eastern SFNA</b>	<b>Entire SFNA</b>
Number of data points	207	505	395	1107
Mean obs (ppb)	70.7	76.6	73	74.2
Standard Deviation obs (ppb)	8.2	12.2	9.5	10.9
Mean Bias (ppb)	3.2	0.6	1.5	1.4
Mean Error (ppb)	7.3	9.8	8.6	8.9
RMSE (ppb)	9.5	13.2	11.1	11.8
Normalized Mean Bias (%)	4.5	0.8	2.1	1.9
Normal Mean Error (%)	10.3	12.8	11.8	12
R-squared	0.2	0.22	0.18	0.21
Index of Agreement	0.67	0.69	0.65	0.68

Table 12. Hourly ozone performance statistics by modeling sub-regions and entire SFNA for the 2012 ozone season (May-October 5<sup>th</sup> 2012). Note that only statistics for the grid cell in which the monitor is located were calculated for hourly ozone.

Hourly ozone > 60 ppb with simulated data extracted at grid cell where the monitor is located				
Parameter	Western SFNA	Central SFNA	Eastern SFNA	Entire SFNA
Number of data points	608	2044	1571	4223
Mean obs (ppb)	69.7	73.9	70.4	72
Standard Deviation obs (ppb)	7.6	10.8	8.2	9.7
Mean Bias (ppb)	0.4	-2.2	-0.6	-1.2
Mean Error (ppb)	6.7	8.7	7.8	8.1
RMSE (ppb)	8.8	11.4	10.4	10.7
Normalized Mean Bias (%)	0.5	-3	-0.8	-1.7
Normal Mean Error (%)	9.6	11.8	11	11.2
R-squared	0.08	0.17	0.06	0.13
Index of Agreement	0.57	0.64	0.55	0.61

Model performance statistics within the range of values shown in Tables 10, 11 and 12 are consistent with previous studies in the SFNA, SJV and studies elsewhere in the U.S. Hu et al. (2012)<sup>1</sup>, simulated an ozone episode in central California (July 27 – August 2, 2000) using a different chemical mechanisms and found that modeled bias ranged from -2.7 to -10.8 ppb for daily maximum 8-hour ozone (compared to -1.6 and 0.2 ppb for the entire SFNA in this work) and -3.6 to -12.7 ppb for daily maximum 1-hour ozone in Central California (compared to -1.4 and 1.4 ppb in this work). Similarly, Shearer et al. (2012)<sup>2</sup> compared model performance in Central California during two episodes in 2000 (July 24 – 26 and July 31 – August 2) for two different chemical mechanisms and found that normalized bias for daily maximum 8-hour ozone ranged from -7% to -14% with hourly peak ozone showing a slightly larger range from -7% to -18%. These values are greater than the statistics found in this work, which were calculated as -2.3% (or 0.3% with 3x3 maximum values) for daily maximum 8-hour

<sup>1</sup> Hu, J., Howard, C. J., Mitloehner, F., Green, P. G., and Kleeman, M. J.: Mobile Source and Livestock Feed Contributions to Regional Ozone Formation in Central California, *Environmental Science & Technology*, 46, 2781-2789, 2012.

<sup>2</sup> Shearer, S. M., Harley, R. A., Jin, L., and Brown, N. J.: Comparison of SAPRC99 and SAPRC07 mechanisms in photochemical modeling for central California, *Atmos. Environ.*, 46, 205-216, 2012.

ozone and -3% (or 1.9% with 3x3 maximum values) for daily maximum 1-hour ozone. Jin et al. (2010)<sup>1</sup> conducted a longer term simulation over Central California (summer 2000) and found a RMSE for daily maximum 8-hour ozone of 14 ppb, which is greater than the 9.3 ppb (or 9.5 ppb with 3x3 maximum values) found in this work. Jin et al. (2010) also showed an overall negative bias of -2 ppb, which is consistent with the -1.6 ppb (0.2 ppb with 3x3 maximum values) found in this work.

Simon et al. (2012)<sup>2</sup> conducted a review of photochemical model performance statistics published between 2006 and 2012 for North America (from 69 peer-reviewed articles). In Figure 12, the statistical evaluation of this model attainment demonstration is compared to the model performance summary presented in Simon et al. (2012) by overlaying the various summary statistics from this attainment demonstration onto the Simon et al. (2012) model performance summary. Note that the box-whisker plot (colored in gray) shown in Figure 12 is reproduced using data from Figure 4 of Simon et al. (2012). The blue and red colored horizontal line markers in each of the panels in Figure 12 denote the model performance statistics from the current modeling work, calculated using the simulated monitor grid cell and the 3x3 maximum, respectively. Figure 11 clearly shows that the model performance statistical metrics for hourly, daily maximum 8-hour and daily maximum 1-hour ozone from this work are consistent with previous modeling studies reported in the scientific literature. In particular, the Simon et al. (2012) study found that mean bias for daily maximum 8-hour ozone ranged from approximately -7 ppb to 13 ppb, while mean error ranged from around 4 ppb to 22 ppb, and RMSE ranged from approximately 8 ppb to 23 ppb; all of which are similar in magnitude to the statistics presented in Table 10. Time series, scatter plots, box plots of mean bias (grouped into 10 ppb bins based on observed values), frequency distribution along with the spatial distribution of mean bias and error plots of the hourly, 1-hour daily maximum and 8-hour daily maximum ozone data used to generate Tables 10, 11 and 12 can be found in the supplementary material.

---

<sup>1</sup> Jin, L., Brown, N. J., Harley, R. A., Bao, J.-W., Michelson, S. A., and Wilczak, J. M.: Seasonal versus episodic performance evaluation for an Eulerian photochemical air quality model, *J. Geophys. Res.*, 115, D09302, doi:10.1029/2009JD012680, 2010.

<sup>2</sup> Simon, H., Baker, K. R., and Phillips, S.: Compilation and interpretation of photochemical model performance statistics published between 2006 and 2012, *Atmospheric Environment*, 61, 124-139, 2012.




Figure 12. Comparison of various statistical metrics from the model attainment demonstration modeling to the range of statistics from the 69 peer-reviewed studies summarized in Simon et al. (2012)<sup>1</sup>. (MDA denotes Maximum Daily Average).

<sup>1</sup> Simon, H., Baker, K. R., and Phillips, S.: Compilation and interpretation of photochemical model performance statistics published between 2006 and 2012, *Atmospheric Environment*, 61, 124-139, 2012.

### 5.2.1 DIAGNOSTIC EVALUATION

In addition to the statistical evaluation presented above, since the modeling is utilized in a relative sense, it is also useful to consider whether the model is able to reproduce observable relationships between changes in emissions and ozone. One approach to this would be to conduct a retrospective analysis where additional years are modeled (e.g., 2000 or 2005) and the ability of the modeling system to reproduce the observed change in ozone over time is investigated. Since this approach is extremely time consuming and resource intensive, it is generally not feasible to perform such an analysis under the constraints of a typical SIP modeling application. Another approach to investigating the ozone response to changes in emissions is through the so called “weekend effect”.




Figure 13. Illustrates a typical ozone isopleth plot, where each line represents ozone mixing ratio, in 10 ppb increments, as a function of initial NO<sub>x</sub> and VOC (or ROG) mixing ratio (adapted from Seinfeld and Pandis, 1998<sup>1</sup>, Figure 5.15). General chemical regimes for ozone formation are shown as NO<sub>x</sub>-disbenefit (red circle), transitional (blue circle), and NO<sub>x</sub>-limited (green circle).

The weekend effect is a well-known phenomenon in some major urbanized areas where emissions of NO<sub>x</sub> are substantially lower on weekends than on weekdays, but

<sup>1</sup> Seinfeld J. H. and Pandis S. N. (1998) Atmospheric Chemistry and Physics: From Air Pollution to Climate Change, 1st edition, J. Wiley, New York.

measured levels of ozone are higher on weekends than on weekdays. This is due to the complex and non-linear relationship between  $\text{NO}_x$  and ROG precursors and ozone (e.g., Swamy et al., 2012)<sup>1</sup>. Ozone formation exhibits a nonlinear dependence to  $\text{NO}_x$  and ROG precursors in the atmosphere. In general terms, under ambient conditions of high- $\text{NO}_x$  and low-ROG ( $\text{NO}_x$ -disbenefit region in Figure 13), ozone formation tends to exhibit a disbenefit to reductions in  $\text{NO}_x$  emissions (i.e., ozone increases with decreases in  $\text{NO}_x$ ) and a benefit to reductions in ROG emissions (i.e., ozone decreases with decreases in ROG). In contrast, under ambient conditions of low- $\text{NO}_x$  and high-ROG ( $\text{NO}_x$ -limited region in Figure 13), ozone formation shows a benefit to reductions in  $\text{NO}_x$  emissions, while changes in ROG emissions result in only minor decreases in ozone. These two distinct “ozone chemical regimes” are illustrated in Figure 13 along with a transitional regime that can exhibit characteristics of both the  $\text{NO}_x$ -disbenefit and  $\text{NO}_x$ -limited regimes. Note that Figure 13 is shown for illustrative purposes only, and does not represent the actual ozone sensitivity within the SFNA for a given combination of  $\text{NO}_x$  and ROG (VOC) emissions.

In this context, the prevalence of a weekend effect in a region suggests that the region is in a  $\text{NO}_x$ -disbenefit regime (Heuss et al., 2003)<sup>2</sup>. A lack of a weekend effect (i.e., no pronounced high  $\text{O}_3$  occurrences during weekends) would suggest that the region is in a transition regime and moving between exhibiting a  $\text{NO}_x$ -disbenefit and being  $\text{NO}_x$ -limited. A reversed weekend effect (i.e., lower  $\text{O}_3$  during weekends) would suggest that the region is  $\text{NO}_x$ -limited.

Investigating the “weekend effect” and how it has changed over time is a useful real world metric for evaluating the ozone chemistry regime in the SFNA and how well it is represented in the modeling. The trend in day-of-week dependence of SFNA’s sub-regional observed ozone levels between 2000 and 2014 is shown in Figure 14. The three-panel scatter plot shown in Figure 14 compares the average site-specific weekday (Wednesday and Thursday) and weekend (Sunday) observed summertime (June through September) maximum daily average (MDA) 8-hour ozone by year (2000 to 2014), separated into three sub-regions: Western SFNA (top), Central SFNA (middle), and Eastern SFNA (bottom). Different definitions of weekday and weekend

---

<sup>1</sup> Swamy, Y.V., Venkanna, R., Nikhil, G.N., Chitanya, D.N.S.K., Sinha, P.R., Ramakrishna, M., and Rao, A.G., 2012. Impact of Nitrogen Oxides, Volatile Organic Compounds and Black Carbon on Atmospheric Ozone Levels at a Semi Arid Urban Site in Hyderabad. *Aerosol and Air Quality Research* 12, 662–671.

<sup>2</sup> Heuss, J.M., Kahlbaum, D.F., and Wolff, G.T., 2003. Weekday/weekend ozone differences: What can we learn from them? *Journal of the Air & Waste Management Association* 53(7), 772-788



days were also investigated and did not show appreciable differences from the Wednesday/Thursday and Sunday definitions.

From Figure 14, it can be seen that ozone levels are highest in the eastern and central regions of the SFNA consistent with their location downwind to and within the urban core of the Sacramento Metropolitan Area. The lowest ozone levels are seen in the western SFNA region, which is located upwind of the urban Sacramento emissions source. In addition, in all regions, summertime average weekday and weekend ozone levels have steadily declined between 2000 and 2014.

Along with the declining ozone, there was shift in the relative difference between weekday and weekend ozone from 2000 and 2014. In the early 2000's, the central region of the SFNA exhibited a roughly equal number sites with weekend ozone greater than weekday ozone as sites with weekday ozone greater than weekend ozone, which suggests that the region may have been in a transitional chemistry regime for ozone formation. By the mid-2000's, the majority of sites were showing weekday ozone greater than weekend ozone, which is consistent with a shift into complete NO<sub>x</sub>-limited chemistry. By 2014, however, some of the sites had shifted back towards a more equal distribution between weekday and weekend ozone, likely due to variability in the biogenic emissions and meteorology that can shift the ozone chemistry between NO<sub>x</sub>-limited and NO<sub>x</sub>-disbenefit regimes in the Sacramento area (LaFranchi et al., 2011)<sup>1</sup>.

The Western SFNA region clearly experienced a greater NO<sub>x</sub>-disbenefit in the early 2000's and then moved into a transitional chemical regime in the mid-2000's and transitioned into the NO<sub>x</sub>-limited regime around the 2010/2011 timeframe. There was a shift back towards a more equal distribution between weekday and weekend ozone by 2014, similar to the Central sub-region. However, this shift occurred at very low ozone levels (below 50 ppb) that are well below the 75 ppb 8-hour ozone standard.

In contrast to the central and western regions described above, the eastern portion of SFNA has been in a NO<sub>x</sub>-limited regime since before 2000, which can be seen from the greater weekday ozone when compared to the weekend ozone. This region is in close proximity to large biogenic ROG emission sources and farther away from the anthropogenic NO<sub>x</sub> sources in the urban Sacramento Metropolitan area, which are conditions (i.e. low NO<sub>x</sub> and high ROG) which place the region in a NO<sub>x</sub>-limited regime.

---

<sup>1</sup> LaFranchi, B. W., Goldstein, A. H., and Cohen, R. C.: Observations of the temperature dependent response of ozone to NO<sub>x</sub> reductions in the Sacramento, CA urban plume, *Atmos. Chem. Phys.*, 11, 6945-6960, doi:10.5194/acp-11-6945-2011, 2011




Figure 14. Site-specific average weekday and weekend maximum daily average 8-hour ozone for each year from 2000 to 2014 for the Western (top), Central (middle), and Eastern (bottom) sub-regions of SFNA. The colored circle markers denote observed values while the magenta triangle markers denote the simulated baseline 2012 values. Points falling below the 1:1 dashed line represent a NO<sub>x</sub>-disbenefit regime, those on the 1:1 dashed line represent a transitional regime, and those above the 1:1 dashed line represent a NO<sub>x</sub>-limited regime.

The simulated baseline 2012 weekday/weekend values (magenta triangle markers in Figure 14) from the attainment demonstration modeling show greater weekday ozone compared to weekend ozone for all three sub-regions, with smaller differences seen in the Central and Western SFNA. These predicted values are consistent with observed findings in 2012 that show a shift into a NO<sub>x</sub>-limited chemistry regime for the Central and Western SFNA and prevalence of NO<sub>x</sub>-limited conditions in Eastern SFNA.

These findings are consistent with an independent analysis by UC Berkeley researchers that examined the observed ozone response due to the decline in NO<sub>x</sub> emissions within the Sacramento area between 2001 and 2007<sup>1</sup>. The study showed a significant decline in 1-hour ozone exceedance days corresponding to a 30% decrease in observed NO<sub>x</sub> due to reductions in NO<sub>x</sub> emissions, and suggesting that NO<sub>x</sub> emission reductions have been effective at reducing ozone levels at all points in Sacramento urban plume. This study concluded that the decline in NO<sub>x</sub> emissions levels has successfully transitioned the region to a NO<sub>x</sub>-limited chemistry regime except within the urban core of the Sacramento Metropolitan Area and predicted that the future cumulative NO<sub>x</sub> controls over time will likely transition the entire SFNA (including the urban core) to a NO<sub>x</sub> limited regime.

### **5.3. RELATIVE RESPONSE FACTORS AND FUTURE YEAR DESIGN VALUES**

The RRFs (Section 2.5) and future year design values (Section 2.6) for the representative sites in the western, central and eastern regions of the SFNA were calculated using the procedures outlined in the corresponding sections, respectively, and are summarized in Table 13. Note that the results shown in Table 13 are ordered by each sub-region in descending order of the average reference year 2012 DVs.

The results in Table 13 show that all monitoring sites in the SFNA have a future DV less than 75 ppb based on the 2026 emissions inventory, with the Folsom monitor in Central SFNA having the highest predicted future design of 70 ppb in 2026 (Note that Folsom is also the valley's design site for base year 2012). Therefore, the air quality simulations predict that the entire region will attain the 75 ppb 8-hour O<sub>3</sub> standard by 2026.

---

<sup>1</sup> LaFranchi, B. W., Goldstein, A. H., and Cohen, R. C.: Observations of the temperature dependent response of ozone to NO<sub>x</sub> reductions in the Sacramento, CA urban plume, *Atmos. Chem. Phys.*, 11, 6945-6960, doi:10.5194/acp-11-6945-2011, 2011

Table 13. Summary of key parameters related to the calculation of future year 2022 and 2026 8-hour ozone design values (DV). Note that final future year design values are truncated, and fractional values are shown for reference only.

Sub-region	Site (County, Air Basin)	Base year 2012	Future year 2022		Future year 2026	
		Average DV (ppb)	RRF	Average DV (ppb)	RRF	Average DV (ppb)
Eastern SFNA	Placerville- Gold Nugget Way (El Dorado, MCAB)	82.3	0.8259	68.0	0.7778	64.0
	Cool-Hwy193 (El Dorado, MCAB)	81.3	0.8336	67.8	0.7882	64.1
	Auburn - Atwood Rd (Placer, SVAB)	79.0	0.8180	64.6	0.7669	60.6
	Colfax-City Hall (Placer, MCAB)	73.7	0.8270	60.9	0.7804	57.5
	Echo Summit (El Dorado, MCAB)	69.0	0.9411	64.9	0.9260	63.9
Central SFNA	<b>Folsom-Natoma Street<sup>1</sup> (Sacramento, SVAB)</b>	<b>90.0</b>	<b>0.8358</b>	<b>75.2<sup>1</sup></b>	<b>0.7857</b>	<b>70.7<sup>1</sup></b>
	Sloughhouse (Sacramento, SVAB)	84.0	0.8459	71.1	0.7998	67.2
	Roseville-N Sunrise Ave (Placer, SVAB)	82.3	0.8487	69.8	0.8055	66.3
	Sacramento-Del Paso Manor (Sacramento, SVAB)	77.3	0.8595	66.4	0.8162	63.1
	North Highlands- Blackfoot Way (Sacramento, SVAB)	76.0	0.8578	65.2	0.8149	61.9
	Sacramento - 1309 T Street (Sacramento, SVAB)	70.0	0.8644	60.5	0.8242	57.7
	Sacramento- Goldenland Court (Sacramento, SVAB)	70.0	0.8820	61.7	0.8415	58.9
Western SFNA	Elk Grove - Bruceville Road (Sacramento, SVAB)	71.7	0.8558	61.4	0.8129	58.3
	Woodland-Gibson Road (Yolo, SVAB)	68.7	0.8459	58.1	0.7996	54.9
	Vacaville-Ulatis Drive (Solano, SVAB)	67.3	0.8459	56.9	0.8009	53.9

<sup>1</sup> The RRF and projected future DVs at the Folsom site do not include the grid cell where the Folsom lake is located (See section 2.5 for details). This does not impact the findings of this attainment demonstration as the future year 2022 and 2026 DVs at Folsom site are estimated to be 73.6 ppb and 68.8 ppb, when the Folsom lake grid cell location is included in RRF and future DV calculations.

	Davis-UCD Campus (Yolo, SVAB)	66.7	0.8495	56.7	0.8052	53.7
--	-------------------------------	------	--------	------	--------	------

The projected 2022 and 2026 DVs in SFNA show a large decrease when compared to 2012 levels (e.g., at the Folsom Natoma Street monitoring site, the SFNA’s design site for 2012, the DV declined by ~15 ppb in 2022 and ~20 ppb in 2026 compared to 2012), which is consistent with the peer-reviewed, published study conducted by the UC Berkeley researchers on the observed response of ozone to NO<sub>x</sub> reductions in the Sacramento area<sup>1</sup>. This study concluded that the region’s ozone exceedance days have been decreasing linearly with decreases in NO<sub>x</sub>, which suggests that cumulative NO<sub>x</sub> controls over time have successfully transitioned the SFNA into a NO<sub>x</sub>-limited chemistry regime, where NO<sub>x</sub> emission reductions are becoming increasingly effective at reducing ozone levels in the region.

**5.4. UNMONITORED AREA ANALYSIS**

The unmonitored area analysis is used to ensure that there are no regions outside of the existing monitoring network that would exceed the NAAQS if a monitor was present (U.S. EPA, 2014<sup>2</sup>). U.S. EPA recommends combining spatially interpolated design value fields with modeled ozone gradients and grid-specific RRFs in order to generate gridded future year gradient adjusted design values.

This analysis can be done using the Model Attainment Test Software (MATS) (Abt, 2014<sup>3</sup>). However, this software is not open source and comes as a precompiled software package. To maintain transparency and flexibility in the analysis, in-house R codes (<https://www.r-project.org/>) developed at ARB, were utilized in this analysis.

The unmonitored area analysis was conducted using the 8-hr O<sub>3</sub> weighted DVs from all the available sites that fall within the 4 km inner modeling domain along with the reference year 2012 and future year 2026 4 km CMAQ model output. The steps followed in the unmonitored area analysis are as follows:

<sup>1</sup> LaFranchi, B. W., Goldstein, A. H., and Cohen, R. C.: Observations of the temperature dependent response of ozone to NO<sub>x</sub> reductions in the Sacramento, CA urban plume, Atmos. Chem. Phys., 11, 6945-6960, doi:10.5194/acp-11-6945-2011, 2011

<sup>2</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM2.5 and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

<sup>3</sup> Abt, 2014. Modeled Attainment Test Software: User’s Manual. MATS available at: [http://www.epa.gov/scram001/modelingapps\\_mats.htm](http://www.epa.gov/scram001/modelingapps_mats.htm)

**Step 1:** At each grid cell, the top-10 modeled maximum daily average 8-hour ozone mixing ratios from the reference year simulation were averaged, and a gradient in this top-10 day average between each grid cell and grid cells which contain a monitor was calculated.

**Step 2:** A single set of spatially interpolated 8-hr ozone DV fields was generated based on the observed 5-year weighted base year 8-hr ozone DVs from the available monitors. The interpolation is done using normalized inverse distance squared weightings for all monitors within a grid cell's Voronoi Region (calculated with the R tripack library; <https://cran.r-project.org/web/packages/tripack/README>), and adjusted based on the gradients between the grid cell and the corresponding monitor from Step 1.

**Step 3:** At each grid cell, the RRFs are calculated based on the reference- and future-year modeling following the same approach outlined in Section 8.3, except that the +/- 20% limitation on the simulated and observed maximum daily average 8-hour ozone was not applied because observed data do not exist for grid cells in unmonitored areas.

**Step 4:** The future year gridded 8-hr ozone DVs were calculated by multiplying the gradient-adjusted interpolated 8-hr ozone DVs from Step 2 with the gridded RRFs from Step 3

**Step 5:** The future-year gridded 8-hr ozone DVs (from Step 4) were examined to determine if there are any peak values higher than those at the monitors, which could potentially cause violations of the applicable 8-hr ozone NAAQS.

Figure 15 shows the spatial distribution of gridded DVs in 2026 for the SFNA based on the unmonitored area analysis (described above). The black colored triangle markers denote the monitoring sites, which had valid reference year 2012 DVs and were used in the analysis. The entire region shows gridded DVs that are below 70 ppb, except for a small region near the center of the spatial map in Figure 15, which shows DVs between 71 and 75 ppb. Those grid cells are located over Folsom Lake and the higher DVs are likely an artifact of the lower mixing heights predicted by the model. Therefore, the unmonitored area analysis predicts that all unmonitored regions within the SFNA will attain the 75 ppb 8-hour O<sub>3</sub> standard by 2026.




Figure 15. Spatial distribution of the future 2026 DVs based on the unmonitored area analysis in the SFNA. Color scale is in ppb of ozone.

### 5.5. “BANDED” RELATIVE RESPONSE FACTORS AND FUTURE YEAR DESIGN VALUES

The “Banded-RRF” approach expands upon the standard “Single-RRF” (Section 5.3) approach to account for differences in model response to emissions controls at varying ozone levels. The most recent U.S. EPA modeling guidance (U. S. EPA, 2014<sup>1</sup>) accounts for some of these differences by focusing on the top ten modeled days, but even the top ten days may contain a significant range of ozone mixing ratios. The Banded-RRF approach accounts for these differences more explicitly by grouping the simulated ozone into bands of lower, medium, and higher ozone mixing ratios.

<sup>1</sup> U.S. EPA, 2014, Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM2.5 and Regional Haze, available at [https://www.epa.gov/ttn/scram/guidance/guide/Draft\\_O3-PM-RH\\_Modeling\\_Guidance-2014.pdf](https://www.epa.gov/ttn/scram/guidance/guide/Draft_O3-PM-RH_Modeling_Guidance-2014.pdf)

In this work, the banded RRFs were calculated to project the future year 2022 and 2026 DVs. The data used for this analysis is inherently consistent with the data used in the single RRF calculations (Sections 2.5 and 5.3). The various steps involved in the calculation of banded RRFs are as follows:

1. MDA 8-hour ozone mixing ratios for all days that are above 60 ppb and that fall within +/- 20% of observations are stratified into 5 ppb increments in the 60 -100 ppb range. (All days above 100 ppb are grouped into a single bin)
2. A separate RRF is calculated for each ozone band following a similar approach as for the standard Single-RRF. A linear regression is then fit to the data resulting in an equation relating RRF to ozone band as long as there are at least 3 bands (without missing data). The band RRF calculations were not available for sites that had fewer than 3 bands of valid RRFs. Similar to the Single-RRF; this equation is unique to each monitor/location.
3. The top ten days for each monitor, based on observed 8-hour ozone for each year of the 5 years that is utilized in the design value calculation (see Table 1), are then projected to the future using the appropriate RRF for the corresponding ozone band.
4. The top ten future days for each individual year are then re-sorted, the fourth highest 8-hour ozone is selected, and the future year design value is calculated in a manner consistent with the base/reference year design value calculation.
5. The future Design Values were then compared with the 75 ppb 8-hour O<sub>3</sub> standard to determine the attainment status for each monitor.

More detailed information on the Banded-RRF approach can be found in Kulkarni et al. (2014)<sup>1</sup> and the SJV 2013 1-Hour Ozone SIP<sup>2</sup>.

The banded RRFs and the corresponding future year 2022 and 2026 design values for the representative sites in the eastern, central, and western regions of the SFNA were calculated using the procedure outlined above, and are summarized in Table 14. Note

<sup>1</sup> Kulkarni, S., Kaduwela, A. P., Avise, J. C., DaMassa, J. A., and Chau, D.: An extended approach to calculate the ozone relative response factors used in the attainment demonstration for the National Ambient Air Quality Standards, *J. Air & Waste Management Association*, 64(10), 1204-1213, 2014, doi:10.1080/10962247.2014.936984.

<sup>2</sup> [http://www.valleyair.org/Air\\_Quality\\_Plans/Ozone-OneHourPlan-2013.htm](http://www.valleyair.org/Air_Quality_Plans/Ozone-OneHourPlan-2013.htm)



that the results shown in Table 14 are ordered by each sub-region in the descending order of average reference year 2012 DVs.

Table 14. Summary of future year (2022 and 2026) design values projected using a banded RRF approach. Note that final future year design values are truncated, and fractional values are shown for reference only.

Sub-region	Site (County, Air Basin)	Base year 2012	Future Year 2022	Future Year 2026
		Average DV (ppb)	Average DV (ppb)	Average DV (ppb)
Eastern SFNA	Placerville-Gold Nugget Way (El Dorado, MCAB)	82.3	67.0	62.3
	Cool-Hwy193 (El Dorado, MCAB)	81.3	66.0	62.0
	Auburn - Atwood Rd (Placer, SVAB)	79.0	65.3	61.3
	Colfax-City Hall (Placer, MCAB)	73.7	60.3	57.3
	Echo Summit (El Dorado, MCAB)	69.0	65.7	65.0
Central SFNA	<b>Folsom-Natoma Street (Sacramento, SVAB)</b>	<b>90.0</b>	<b>74.0</b>	<b>69.0</b>
	Sloughhouse (Sacramento, SVAB)	84.0	70.7	67.0
	Roseville-N Sunrise Ave (Placer, SVAB)	82.3	68.7	64.7
	Sacramento-Del Paso Manor (Sacramento, SVAB)	77.3	66.0	63.0
	North Highlands-Blackfoot Way (Sacramento, SVAB)	76.0	65.0	62.0
	Sacramento - 1309 T Street (Sacramento, SVAB)	70.0	61.0	58.0
	Sacramento-Goldenland Court (Sacramento, SVAB)	70.0	61.3	59.3
Western SFNA	Elk Grove - Bruceville Road (Sacramento, SVAB)	71.7	61.7	58.7
	Woodland-Gibson Road (Yolo, SVAB)	68.7	58.3	55.7
	Vacaville-Ulatis Drive (Solano, SVAB)	67.3	57.0	54.0
	Davis-UCD Campus (Yolo, SVAB)	66.7	-	-

The results in Table 14 show that all the monitoring sites in the SFNA have a future DV less than 75 ppb, with the Folsom Natoma Street monitoring site in Central SFNA having the highest predicted future design value with an estimated future design value of 74 and 69 ppb in 2022 and 2026, respectively. These future DV's are ~1 ppb lower than the corresponding single-RRF values (Table 13).

## 6. OZONE ISOPLETHS

Since the entire SFNA is projected to be in attainment for the 2008 75 ppb 8-hour O<sub>3</sub> standard, no additional emission reductions beyond what is being implemented through the current control program will be necessary. However, the U.S. EPA revised the 8-hr O<sub>3</sub> standard to a level of 0.070 ppm (70 ppb) in October 2015<sup>1</sup>, for which the final designations are due in late 2017. Hence, it is important to know the emission targets in the future to assess the level of emissions controls needed to attain the 2015 8-hour O<sub>3</sub> standard of 0.070 ppm (70 ppb). Although 2026 DVs at all monitoring sites are predicted to be in attainment of the 70 ppb ozone standard, it is still useful to examine the future DV sensitivity to precursor emissions to evaluate how this sensitivity may change in the future.

To examine the future ozone sensitivity within the SFNA for different combinations of NO<sub>x</sub> and VOC (ROG) emissions in the region, modeling sensitivity simulations were conducted to generate 8-hr ozone isopleths. These sensitivity simulations are identical to the future year 2026 simulation discussed earlier in Section 2.4 and Table 3, except that domain-wide fractional reductions were applied to future year 2026 anthropogenic NO<sub>x</sub> and ROG emission levels. Each sensitivity simulation was run for the entire ozone season (May – October 5<sup>th</sup> 2012) and included statewide 12 km simulations nested down to 4 km. The inner 4 km domain sensitivity simulations utilized BCs based on output from the corresponding 12 km sensitivity simulation, while the 12 km simulations all utilized the same MOZART derived BCs. The RRF methodology described in Section 2.5 was then applied to the inner 4 km domain output of each fractional ROG and NO<sub>x</sub> sensitivity simulation to calculate the future year DV (for that specific NO<sub>x</sub>-ROG combination) at each monitoring site in the SFNA.

Figure 16 shows the 2026 8-hour ozone isopleths for the Folsom monitoring site (isopleths for other sites are not shown since their projected DVs are below 70 ppb). In Figure 16, the bottom and top axes represent the domain-wide fractional ROG emissions and the corresponding SFNA emission totals (tons per day) in 2026,

---

<sup>1</sup> Federal Register, Vol. 80 , No. 206, October 26, 2015, National Ambient Air Quality Standards for Ozone, Final Rule, Pages 65291-65468

respectively. Similarly, the left and right axes represent the domain-wide fractional NO<sub>x</sub> emissions and the corresponding SFNA emission totals (tons per day) in 2026, respectively. The upper right point on each diagram represents the projected DV for the attainment demonstration modeling (listed in Table 13).

The shape of the ozone isopleth shown in Figure 16 indicates that it falls in the bottom right corner of the Figure 13, where the NO<sub>x</sub>-limited regime is prevalent. It is evident from this diagram that the future O<sub>3</sub> mixing ratios throughout the SFNA are predicted to be in the NO<sub>x</sub>-limited regime and that the sensitivity to ROG emissions controls will be much lower when compared to NO<sub>x</sub> controls.




Figure 16. The 8-hr ozone isopleth based on 2026 emission levels at the Folsom Natoma Street monitoring site located in Central SFNA.

## SUPPLEMENTAL MATERIALS

## SUPPLEMENTAL MATERIALS TABLE OF CONTENTS

MONTHLY METEOROLOGICAL TIME SERIES PLOTS.....	60
OZONE PLOTS.....	76
HOURLY OZONE TIMESERIES PLOTS.....	83
DAILY MAXIMUM 1 – HOUR OZONE TIME SERIES PLOTS .....	100
DAILY MAXIMUM 8 – HOUR OZONE TIME SERIES PLOTS .....	107

## SUPPLEMENTAL MATERIALS LIST OF FIGURES

Figure S. 1 Time series of wind speed, direction, and temperature for Valley in May 2012. ....	61
Figure S. 2 Time series of wind speed, direction, and temperature for Mountain in May 2012. ....	62
Figure S. 3 Time series of wind speed, direction, and temperature for Valley in June 2012. ....	63
Figure S. 4 Time series of wind speed, direction, and temperature for Mountain in June 2012. ....	64
Figure S. 5 Time series of wind speed, direction, and temperature for Valley in July 2012. ....	65
Figure S. 6 Time series of wind speed, direction, and temperature for Mountain in July 2012. ....	66
Figure S. 7 Time series of wind speed, direction, and temperature for Valley in August 2012. ....	67
Figure S. 8 Time series of wind speed, direction, and temperature for Mountain in August 2012. ....	68
Figure S. 9 Time series of wind speed, direction, and temperature for Valley in September through October 5 2012. ....	69
Figure S. 10 Time series of wind speed, direction, and temperature for Mountain in September through October 5 2012. ....	70
Figure S. 11 Time series of relative humidity for Valley and Mountain in May 2012. ....	71
Figure S. 12 Time series of relative humidity for Valley and Mountain in June 2012. ....	72
Figure S. 13 Time series of relative humidity for Valley and Mountain in July 2012. ....	73
Figure S. 14 Time series of relative humidity for Valley and Mountain in August 2012. ....	74
Figure S. 15 Time series of relative humidity for Valley and Mountain in September through October 5 2012. ....	75
Figure S. 16 Observed and modeled ozone frequency distribution for the ozone season (May – October 5 <sup>th</sup> 2012) ....	77
Figure S. 17 Comparison of modeled ozone with observations for the ozone season (May – October 5 <sup>th</sup> 2012) ....	78
Figure S. 18 Spatial distribution of ozone mean bias (left) and mean error (right) for the ozone season (May-October 5 <sup>th</sup> 2012). ....	79

Figure S. 19 Hourly Ozone Site Mean Bias Distribution for the ozone season (May-October 5 <sup>th</sup> 2012) .....	80
Figure S. 20 Daily Maximum 1-hour Ozone Site Mean Bias Distribution for the ozone season (May-October 5 <sup>th</sup> 2012).....	81
Figure S. 21 Daily Maximum Average 8-hour Ozone Site Mean Bias Distribution for the ozone season (May-October 5 <sup>th</sup> 2012).....	82
Figure S. 22 Time-series of hourly ozone at Cool Hwy 193 monitor .....	84
Figure S. 23 Time-series of hourly ozone at Placerville Gold Nugget Way .....	85
Figure S. 24 Time-series of hourly ozone at Auburn Atwood road .....	86
Figure S. 25 Time-series of hourly ozone at Colfax City Hall .....	87
Figure S. 26 Time-series of hourly ozone at Roseville-N. Sunrise Ave .....	88
Figure S. 27 Time-series of hourly ozone at Elk Grove – Bruceville Road .....	89
Figure S. 28 Time-series of hourly ozone at Folsom Natoma Street .....	90
Figure S. 29 Time-series of hourly ozone at North Highlands – Blackfoot way .....	91
Figure S. 30 Time-series of hourly ozone at Sacramento – Del Paso Manor.....	92
Figure S. 31 Time-series of hourly ozone at Sacramento – Goldenland Court .....	93
Figure S. 32 Time-series of hourly ozone at Sacramento – T Street.....	94
Figure S. 33 Time-series of hourly ozone at Sloughhouse .....	95
Figure S. 34 Time-series of hourly ozone at Vacaville Ulatis Drive .....	96
Figure S. 35 Time-series of hourly ozone at Davis – UCD Campus.....	97
Figure S. 36 Time-series of hourly ozone at Woodland – Gibson road .....	98
Figure S. 37 Time-series of hourly ozone at Echo Summit .....	99
Figure S. 38 Time-series of daily maximum 1-hour ozone at Cool – Highway 193 .....	101
Figure S. 39 Time-series of daily maximum 1-hour ozone at Placerville – Gold Nugget way.....	101
Figure S. 40 Time-series of daily maximum 1-hour ozone at Auburn – Antwoo Road	101
Figure S. 41 Time-series of daily maximum 1-hour ozone at Colfax City Hall.....	102
Figure S. 42 Time-series of daily maximum 1-hour ozone at Roseville – N Sunrise Ave .....	102
Figure S. 43 Time-series of daily maximum 1-hour ozone at Elk Grove – Bruceville road .....	102

Figure S. 44 Time-series of daily maximum 1-hour ozone at Folsom – Natoma street 103

Figure S. 45 Time-series of daily maximum 1-hour ozone at North Highlands –  
Blackfoot way ..... 103

Figure S. 46 Time-series of daily maximum 1-hour ozone at Sacramento – Del Paso  
Manor ..... 103

Figure S. 47 Time-series of daily maximum 1-hour ozone at Sacramento – Goldenland  
Court ..... 104

Figure S. 48 Time-series of daily maximum 1-hour ozone at Sacramento – T street.. 104

Figure S. 49 Time-series of daily maximum 1-hour ozone at Sloughhouse ..... 104

Figure S. 50 Time-series of daily maximum 1-hour ozone at Vacaville-Ulatis Drive ... 105

Figure S. 51 Time-series of daily maximum 1-hour ozone at Davis – UCD campus... 105

Figure S. 52 Time-series of daily maximum 1-hour ozone at Woodland – Gibson road  
..... 105

Figure S. 53 Time-series of daily maximum 1-hour ozone at Echo Summit ..... 106

Figure S. 54 Time-series of daily maximum average 8-hour ozone at Cool – Highway  
193 ..... 108

Figure S. 55 Time-series of daily maximum average 8-hour ozone at Placerville – Gold  
Nugget Way ..... 108

Figure S. 56 Time-series of daily maximum average 8-hour ozone at Auburn Antwook  
Road..... 108

Figure S. 57 Time-series of daily maximum average 8-hour ozone at Colfax – City Hall  
..... 109

Figure S. 58 Time-series of daily maximum average 8-hour ozone at Roseville N.  
Sunrise Ave ..... 109

Figure S. 59 Time-series of daily maximum average 8-hour ozone at Elk Grive  
Bruceville Road ..... 109

Figure S. 60 Time-series of daily maximum average 8-hour ozone at Folsom Natoma  
Street..... 110

Figure S. 61 Time-series of daily maximum average 8-hour ozone at North Highlands –  
Blackfoot way ..... 110

Figure S. 62 Time-series of daily maximum average 8-hour ozone at Sacramento – Del  
Paso Manor ..... 110



Figure S. 63 Time-series of daily maximum average 8-hour ozone at Sacramento Goldenland Court ..... 111

Figure S. 64 Time-series of daily maximum average 8-hour ozone at Sacramento – T street ..... 111

Figure S. 65 Time-series of daily maximum average 8-hour ozone at Sloughhouse .. 111

Figure S. 66 Time-series of daily maximum average 8-hour ozone at Vacaville – Ulatis Drive ..... 112

Figure S. 67 Time-series of daily maximum average 8-hour ozone at Davis – UCD campus..... 112

Figure S. 68 Time-series of daily maximum average 8-hour ozone at woodland- Gibson Road..... 112

Figure S. 69 Time-series of daily maximum average 8-hour ozone at Echo Summit .. 113

## MONTHLY METEOROLOGICAL TIME SERIES PLOTS




Figure S. 1 Time series of wind speed, direction, and temperature for Valley in May 2012.




Figure S. 2 Time series of wind speed, direction, and temperature for Mountain in May 2012.




Figure S. 3 Time series of wind speed, direction, and temperature for Valley in June 2012.




Figure S. 4 Time series of wind speed, direction, and temperature for Mountain in June 2012.




Figure S. 5 Time series of wind speed, direction, and temperature for Valley in July 2012.




Figure S. 6 Time series of wind speed, direction, and temperature for Mountain in July 2012.






Figure S. 7 Time series of wind speed, direction, and temperature for Valley in August 2012.




Figure S. 8 Time series of wind speed, direction, and temperature for Mountain in August 2012.




Figure S. 9 Time series of wind speed, direction, and temperature for Valley in September through October 5 2012.




Figure S. 10 Time series of wind speed, direction, and temperature for Mountain in September through October 5 2012.




Figure S. 11 Time series of relative humidity for Valley and Mountain in May 2012.




Figure S. 12 Time series of relative humidity for Valley and Mountain in June 2012.




Figure S. 13 Time series of relative humidity for Valley and Mountain in July 2012.




Figure S. 14 Time series of relative humidity for Valley and Mountain in August 2012.






Figure S. 15 Time series of relative humidity for Valley and Mountain in September through October 5 2012.

## OZONE PLOTS




Figure S. 16 Observed and modeled ozone frequency distribution for the ozone season (May – October 5<sup>th</sup> 2012)




Figure S. 17 Comparison of modeled ozone with observations for the ozone season (May – October 5<sup>th</sup> 2012)




Figure S. 18 Spatial distribution of ozone mean bias (left) and mean error (right) for the ozone season (May-October 5<sup>th</sup> 2012).




Figure S. 19 Hourly Ozone Site Mean Bias Distribution for the ozone season (May-October 5<sup>th</sup> 2012)




Figure S. 20 Daily Maximum 1-hour Ozone Site Mean Bias Distribution for the ozone season (May-October 5<sup>th</sup> 2012)




Figure S. 21 Daily Maximum Average 8-hour Ozone Site Mean Bias Distribution for the ozone season (May-October 5<sup>th</sup> 2012)



## HOURLY OZONE TIMESERIES PLOTS




Figure S. 22 Time-series of hourly ozone at Cool Hwy 193 monitor




Figure S. 23 Time-series of hourly ozone at Placerville Gold Nugget Way




Figure S. 24 Time-series of hourly ozone at Auburn Atwood road




Figure S. 25 Time-series of hourly ozone at Colfax City Hall




Figure S. 26 Time-series of hourly ozone at Roseville – N. Sunrise Ave




Figure S. 27 Time-series of hourly ozone at Elk Grove – Bruceville Road




Figure S. 28 Time-series of hourly ozone at Folsom Natoma Street






Figure S. 29 Time-series of hourly ozone at North Highlands – Blackfoot way




Figure S. 30 Time-series of hourly ozone at Sacramento – Del Paso Manor




Figure S. 31 Time-series of hourly ozone at Sacramento – Goldenland Court




Figure S. 32 Time-series of hourly ozone at Sacramento – T Street




Figure S. 33 Time-series of hourly ozone at Sloughhouse




Figure S. 34 Time-series of hourly ozone at Vacaville Ulatis Drive




Figure S. 35 Time-series of hourly ozone at Davis – UCD Campus




Figure S. 36 Time-series of hourly ozone at Woodland – Gibson road






Figure S. 37 Time-series of hourly ozone at Echo Summit

**DAILY MAXIMUM 1 – HOUR OZONE TIME SERIES PLOTS**




Figure S. 38 Time-series of daily maximum 1-hour ozone at Cool – Highway 193




Figure S. 39 Time-series of daily maximum 1-hour ozone at Placerville – Gold Nugget way




Figure S. 40 Time-series of daily maximum 1-hour ozone at Auburn – Antwoot Road




Figure S. 41 Time-series of daily maximum 1-hour ozone at Colfax City Hall




Figure S. 42 Time-series of daily maximum 1-hour ozone at Roseville – N Sunrise Ave




Figure S. 43 Time-series of daily maximum 1-hour ozone at Elk Grove – Bruceville road




Figure S. 44 Time-series of daily maximum 1-hour ozone at Folsom – Natoma street




Figure S. 45 Time-series of daily maximum 1-hour ozone at North Highlands – Blackfoot way




Figure S. 46 Time-series of daily maximum 1-hour ozone at Sacramento – Del Paso Manor




Figure S. 47 Time-series of daily maximum 1-hour ozone at Sacramento – Goldenland Court




Figure S. 48 Time-series of daily maximum 1-hour ozone at Sacramento – T street




Figure S. 49 Time-series of daily maximum 1-hour ozone at Sloughouse




Figure S. 50 Time-series of daily maximum 1-hour ozone at Vacaville-Ulatis Drive




Figure S. 51 Time-series of daily maximum 1-hour ozone at Davis – UCD campus




Figure S. 52 Time-series of daily maximum 1-hour ozone at Woodland – Gibson road




Figure S. 53 Time-series of daily maximum 1-hour ozone at Echo Summit



**DAILY MAXIMUM 8 – HOUR OZONE TIME SERIES PLOTS**




Figure S. 54 Time-series of daily maximum average 8-hour ozone at Cool – Highway 193




Figure S. 55 Time-series of daily maximum average 8-hour ozone at Placerville – Gold Nugget Way




Figure S. 56 Time-series of daily maximum average 8-hour ozone at Auburn Antwoot Road




Figure S. 57 Time-series of daily maximum average 8-hour ozone at Colfax – City Hall




Figure S. 58 Time-series of daily maximum average 8-hour ozone at Roseville N. Sunrise Ave




Figure S. 59 Time-series of daily maximum average 8-hour ozone at Elk Grove Bruceville Road




Figure S. 60 Time-series of daily maximum average 8-hour ozone at Folsom Natoma Street




Figure S. 61 Time-series of daily maximum average 8-hour ozone at North Highlands – Blackfoot way




Figure S. 62 Time-series of daily maximum average 8-hour ozone at Sacramento – Del Paso Manor




Figure S. 63 Time-series of daily maximum average 8-hour ozone at Sacramento Goldenland Court




Figure S. 64 Time-series of daily maximum average 8-hour ozone at Sacramento – T street




Figure S. 65 Time-series of daily maximum average 8-hour ozone at Sloughouse




Figure S. 66 Time-series of daily maximum average 8-hour ozone at Vacaville – Ulatis Drive




Figure S. 67 Time-series of daily maximum average 8-hour ozone at Davis – UCD campus




Figure S. 68 Time-series of daily maximum average 8-hour ozone at woodland- Gibson Road




Figure S. 69 Time-series of daily maximum average 8-hour ozone at Echo Summit

## **Appendix B-5**

## **Modeling Emissions Inventory**

### **Document Title:**

Modeling Emission Inventory for the 8-Hour Ozone State Implementation Plan in the Sacramento Non-Attainment Area

### **Document Description:**

This document describes how the base and future year gridded photochemical modeling emissions inventory are prepared.



# **Modeling Emission Inventory for the 8-Hour Ozone State Implementation Plan in the Sacramento Non-Attainment Area**

## **Prepared by**

California Air Resources Board

El Dorado County Air Quality Management District

Feather River Air Quality Management District

Placer County Air Pollution Control District

Sacramento Air Quality Management District

Yolo-Solano Air Quality Management District

## **Prepared for**

United States Environmental Protection Agency Region IX

January 27, 2017

## Contents

1. Development of Ozone Emissions Inventories .....	7
1.1. Inventory Coordination .....	7
1.2. Background .....	8
1.3. Inventory Years .....	9
1.3.1. Base Case Modeling Inventory (2012).....	9
1.3.2. Reference Year (or Baseline) Modeling Inventory (2012).....	10
1.3.3. Future Year Modeling Inventory (2022/2026) .....	10
1.3.4. 2012 Base Case Modeling Inventory .....	11
1.3.5. 2012 Reference Year (Baseline) Modeling Inventory .....	11
1.3.6. 2022/2026 Future Year Modeling Inventories.....	11
1.4. Spatial Extent of Emission Inventories .....	11
2. Estimation of Base Year Modeling Inventory .....	13
2.1. Terminology .....	13
2.2. Temporal Distribution of Emissions.....	14
2.2.1. Monthly Variation .....	15
2.2.2. Weekly Variation.....	15
2.2.3. Daily Variation .....	16
2.3. Spatial Allocation.....	18
2.3.1. Spatial Allocation of Area Sources .....	22
2.3.2. Spatial Allocation of Point Sources.....	22

---

2.3.3.	Spatial Allocation of Wildfires, Prescribed Burns and Wildland Fire Use ...	22
2.3.4.	Spatial Allocation of Ocean going vessels (OGV).....	22
2.3.5.	Spatial Allocation of On-road Motor Vehicles .....	23
2.3.6.	Spatial Allocation of Biogenic Emissions .....	23
2.4.	Speciation Profiles .....	23
3.	Methodology for Developing Base Case and Baseline Emissions Inventories.....	26
3.1.	Surface Temperature and Relative Humidity Fields .....	27
3.2.	Insolation Effects.....	28
3.3.	Estimation of Gridded Area and Point sources.....	28
3.4.	Estimation of On-road Motor Vehicle Emissions .....	29
3.4.1.	General Methodology .....	30
3.4.2.	ITN Activity Data .....	33
3.4.3.	Spatial Adjustment.....	34
3.4.4.	Temporal Adjustment (Day-of-Week adjustments to EMFAC daily totals).	35
3.4.5.	Temporal Adjustment (Hour-of-Day re-distribution of hourly travel network volumes) .....	36
3.4.6.	Summary of On-road Emissions Processing Steps .....	37
3.5.	Estimation of Gridded Biogenic Emissions.....	39
3.6.	Estimation of Other Day-Specific Sources .....	40
3.6.1.	Wildfires and Prescribed Burns .....	40
3.6.2.	Paved Road Dust .....	42

---

3.6.3. Unpaved Road Dust .....	43
3.6.4. Agricultural Burning .....	44
3.6.6. Closed Facilities .....	48
4. Quality Assurance of Modeling Inventories .....	49
4.1. Area and Point Sources .....	49
4.1.1. Area and Point Sources Temporal Profiles .....	51
4.2. On-road Emissions.....	52
4.3. Day-specific Sources .....	53
4.3.1. Wildfires and Prescribed Burns .....	53
4.3.2. Paved Road Dust .....	54
4.3.3. Unpaved Road Dust .....	54
4.3.4. Agricultural Burning .....	54
4.3.5. Refinery Fire .....	55
4.4. Additional QA .....	55
4.5. Model ready files QA.....	58
Bibliography .....	59
Appendix A: Day of week redistribution factors by vehicle type and county .....	63
Appendix B: Hour of Day Profiles by vehicle type and county .....	68
Appendix C: Scaling procedures after DTIM processing .....	95
Appendix D: Additional temporal profiles.....	97

## List of Figures

Figure 1 Spatial coverage and parameter summary of modeling domains.....	12
Figure 2 Block diagram for on-road processing.....	32
Figure 3 Example of a spatial plot by source category .....	50
Figure 4 Screen capture of a SMOKE-generated QA report .....	51
Figure 5 Screenshot of comparison of inventories report.....	56
Figure 6 Daily variation of NOx emissions for mobile sources for San Luis Obispo .....	57

## List of Tables

Table 1 Modeling domain parameters.....	13
Table 2 Inventory terms for emission source types .....	14
Table 3 Day of week variation factors .....	16
Table 4 Daily variation factors .....	17
Table 5 Spatial Surrogates.....	20
Table 6 Vintage of travel demand models for link based and traffic analysis zone .....	33
Table 7 DTIM Emission Categories.....	34
Table 8 Vehicle classification and type of adjustment .....	35
Table 9 Day of week adjustment by vehicle class and county.....	63
Table 10 Hour of Day Profiles by vehicle type and county .....	68

Table 11 Day of week temporal profiles from the Agricultural Emissions Temporal and Spatial Allocation Tool (AgTool) ..... 97

Table 12 Daily temporal profiles from the Agricultural Emissions Temporal and Spatial Allocation Tool (AgTool) ..... 99

## **1. Development of Ozone Emissions Inventories**

Emission inputs for air quality modeling (commonly and interchangeably referred to as 'modeling inventories' or 'gridded inventories') have been developed by ARB and district staff. These inventories support the different SIPs across California to meet various federal PM<sub>2.5</sub> standards. ARB maintains an electronic database of emissions and other useful information to generate aggregate emission estimates at the county, air basin and district level. This database is called the California Emission Inventory Development and Reporting System (CEIDARS). CEIDARS provides a foundation for the development of a more refined (hourly, grid-cell specific) set of emission inputs that are required by air quality models. The CEIDARS base year inventory is a primary input to the state's emission forecasting system, known as the California Emission Projection Analysis Model (CEPAM). CEPAM produces the projected emissions that are then gridded and serve as the emission input for the particulate matter models.

The following sections of this document describe how base and future year emissions inventory estimates are prepared.

### **1.1. Inventory Coordination**

The Air Resources Board convened the SIP Inventory Working Group (SIPIWG) to provide an opportunity and means for interested parties (ARB, districts, etc.) to discuss issues pertaining to the development and review of base year, future year, planning and gridded inventories to be used in SIP modeling. The group has met every four to six weeks since March 2013. Group participants included district staff from Bay Area, Butte, Eastern Kern, El Dorado, Feather River, Imperial, Northern Sierra, Placer, Sacramento, San Diego, San Joaquin, San Luis Obispo, South Coast, Ventura and Yolo-Solano.

Additionally, ARB established the SIPIWG Spatial Surrogate Sub-committee, which focused on improving input data to spatially disaggregate emissions at a more refined level needed for air quality modeling. Local air districts that participated included San Joaquin Valley APCD, South Coast AQMD, Ventura County APCD and Sacramento Metropolitan AQMD.

In addition to the two coordination groups described above, a great deal of work preceded this modeling effort through the Central California Air Quality Studies (CCAQS). CCAQS consisted of two studies: 1) the Central California Ozone Study (CCOS); and 2) the California Regional PM<sub>10</sub>/PM<sub>2.5</sub> Air Quality Study (CRPAQS).

## **1.2. Background**

California's emission inventory is an estimate of the amounts and types of pollutants emitted from thousands of industrial facilities, millions of motor vehicles and a myriad of emission sources such as consumer products and fireplaces. The development and maintenance of the emission inventory involves several agencies. This multi-agency effort includes: ARB, 35 local air pollution control and air quality management districts (Districts), regional transportation planning agencies (RTPAs), and the California Department of Transportation (Caltrans). The ARB is responsible for the compilation of the final statewide emission inventory, and for maintaining this information in CEIDARS. In addition to the statewide emission inventory, emissions from northern Mexico (Jackson, 2012) are also incorporated in the final emission inventory used for modeling. The final emission inventory reflects the best information available at the time.

The basic principle for estimating county-wide regulatory emissions is to multiply an estimated, per-unit emission factor by an estimate of typical usage or activity. For example, on-road motor vehicle emission factors are estimated for a specific vehicle type and applied to all applicable vehicles. The estimates are based on dynamometer tests of a small sample for a vehicle type. The activity for any given vehicle type is based on an estimate of typical driving patterns, number of vehicle starts, and typical miles driven. Assumptions are also made regarding typical usage; it is assumed that all vehicles of a certain vehicle type are driven under similar conditions in each region of the state.

Developing emission estimates for stationary sources involves the use of per unit emission factors and activity levels. Under ideal conditions, facility-specific emission factors are determined from emission tests for a particular process at a facility. A continuous emission monitoring system (CEMS) can also be used to determine a gas or



particulate matter concentration or emission rate (U.S. EPA, 2016). More commonly, a generic emission factor is developed by averaging the results of emission tests from similar processes at several different facilities. This generic factor is then used to estimate emissions from similar types of processes when a facility-specific emission factor is not available. Activity levels from stationary sources are measured in terms such as the amount of product produced, solvent used, or fuel used.

The district reported or ARB estimated emissions totals are stored in the CEIDARS database for any given pollutant. Both criteria and toxic air pollutant emission inventories are stored in this complex database. These are typically annual average emissions for each county, air basin, and district. Modeling inventories for reactive organic gases (ROG) are estimated from total organic gases (TOG). Similarly, the modeling inventories for total particulate matter 10 $\mu$  in diameter and smaller (PM<sub>10</sub>) and total particulate matter 2.5 $\mu$  in diameter and smaller (PM<sub>2.5</sub>) are estimated from total particulate matter (PM). Details about chemical and size resolved speciation of emissions for modeling can be found in Section 2.4. Additional information on ARB emission inventories can be found at: <http://www.arb.ca.gov/ei/ei.htm>.

### 1.3. Inventory Years

The emission inventory scenarios used for air quality modeling must be consistent with U.S. EPA's Modeling guidance (U.S. EPA, 2014). Since changes in the emissions inventory can affect the calculation of the relative response factors (RRFs), the terms used in the preparation of the emission inventory scenarios must be clearly defined. In this document the following inventory definitions will be used:

**1.3.1. Base Case Modeling Inventory (2012):** Base case modeling is intended to evaluate model performance and demonstrate confidence in the modeling system used for the modeled attainment test. The base case modeling inventory is not used as part of the modeled attainment test itself. Model performance is assessed relative to how well model-simulated concentrations match actual measured concentrations. The modeling inputs are developed to represent (as best as possible) actual, day-specific conditions. Therefore, the

base case modeling inventory for 2012 includes day-specific emissions for certain sectors. This includes, for instance, actual district-reported point source emissions information for 2012, as well as available day-specific activities and emission adjustments. The year 2012 was selected to coincide with the year selected for baseline design values (described below). The U.S. EPA modeling guidance states that once the model has been shown to perform adequately, the use of day-specific emissions is no longer needed. In preparation for SIP development, both ARB and the local air districts began a comprehensive review and update of the emission inventory several years ago resulting in a comprehensive emissions inventory for 2012.

**1.3.2. Reference Year (or Baseline) Modeling Inventory (2012):** The baseline or reference year inventory is intended to be a representation of emission patterns occurring through the baseline design value period and the emission patterns expected in the future year. U.S. EPA modeling guidance describes the reference year modeling inventory as “a common starting point” that represents average or “typical” conditions that are consistent with the baseline design value period. U.S. EPA guidance also states “using a ‘typical’ or average reference year inventory provides an appropriate platform for comparisons between baseline and future years.” The 2012 reference year inventory represents typical average conditions and emission patterns through the 2012 design value period. The baseline inventory includes temperature, relative humidity and solar insolation effects, for 2012.

**1.3.3. Future Year Modeling Inventory (2022/2026):** Future year modeling inventories, along with the reference year modeling inventory, are used in the model-derived RRF calculation. These inventories maintain the “typical”, average patterns of the 2012 reference year modeling inventory. The 2022 or 2026 inventory will include temperature, relative humidity, and solar insolation effects from reference year (2012) meteorology. Future year point and area source emissions are projected from the 2012 baseline emissions used in the

2012 reference year modeling inventory. Additionally, future year on-road emission inventories are used, as projected by EMFAC.

In summary and based on the definitions above, the following modeling emission inventories were developed:

**1.3.4. 2012 Base Case Modeling Inventory:** This day-specific inventory is used for the model performance evaluation.

**1.3.5. 2012 Reference Year (Baseline) Modeling Inventory:** This 2012 reference year inventory is used to determine site-specific RRFs in the modeled attainment test. The 2012 reference year modeling inventory represents typical, average conditions and emission patterns over the baseline design value period, and includes 2012 meteorological effects.

**1.3.6. 2022/2026 Future Year Modeling Inventories:** These typical, average-day inventories are used to determine site-specific RRFs in the modeled attainment test. Consistent with the 2012 reference year modeling inventory, the 2022 or 2026 inventory is projected from the 2012 baseline inventory and includes 2012 meteorological effects.

#### **1.4. Spatial Extent of Emission Inventories**

The emissions model-ready files that are prepared for use as an input for the air quality model conform to the definition and extent of the grids shown in Figure 1.




Figure 1 Spatial coverage and parameter summary of modeling domains

The domain uses a Lambert projection and assumes a spherical Earth. The emissions inventory grid uses a Lambert Conical Projection with two parallels. The parallels are at 30° and 60° N latitude, with a central meridian at 120.5° W longitude. The coordinate system origin is offset to 37° N latitude. The emissions inventory uses a grid with a spatial resolution of 4 km x 4 km. The state modeling domain (ST4K) extends entirely over California and 100 nautical miles west over the Pacific Ocean. A smaller subdomain (CCOS) is used for Sacramento area. It has the same grid definitions and resolution as the main domain, but has a smaller area offset to cover central and

northern California. The specifications of the emissions inventory domain and CCOS subdomain are summarized in Table 1.

Table 1 Modeling domain parameters

Parameter	Statewide domain (ST4K)	Subdomain (CCOS)
Map Projection	Lambert Conformal Conic	Lambert Conformal Conic
<b>Datum</b>	None (Clarke 1866 spheroid)	None (Clarke 1866 spheroid)
1st Standard Parallel	30.0° N	30.0° N
2nd Standard Parallel	60.0° N	60.0° N
Central Meridian	-120.5° W	-120.5° W
Latitude of projection origin	37.0° N	37.0° N
<b>COORDINATE SYSTEM</b>		
Units	Meters	Meters
Semi-major axis	6370 km	6370 km
Semi-minor axis	6370 km	6370 km
<b>DEFINITION OF GRID</b>		
Grid size	4km x 4km	4km x 4km
Number of cells	321 x 291 cells	192 x 192 cells
Lambert origin	(-684,000 m, -564,000 m)	(-384,000 m, -300,000 m)
Geographic center	-120.5° Lat and 37.0° Lon	-120.5° Lat and 37.0° Lon

## 2. Estimation of Base Year Modeling Inventory

As mentioned in Section 1.3, base case modeling is intended to demonstrate confidence in the modeling system used for the modeled attainment test. The following sections describe the temporal and spatial distribution of emissions and how the different sectors of the modeling inventories are prepared.

### 2.1. Terminology

The terms “point sources” and “area sources” are often confused. Traditionally, these terms have had different meanings to the developers of emissions inventories and the developers of modeling inventories. Table 2 summarizes the difference in the terms. Both sets of terms are used in this document. In modeling terminology, “point sources”

traditionally refer to elevated emission sources that exit from a stack and have an associated plume rise. While the current inventory includes emissions from stacks, all emission sources reported by the SJVAPCD associated with a facility are treated as potential elevated sources. The emissions processor calculates plume rise if appropriate; non-elevated sources are treated as ground-level sources. Examples of non-elevated emissions sources include gas dispensing facilities and storage piles. “Area sources” refers collectively to area-wide sources, stationary-aggregated sources, and other mobile sources (including aircraft, trains, ships, and all off-road vehicles and equipment). That is, “area sources” are low-level sources from a modeling perspective.

Table 2 Inventory terms for emission source types

Modeling Term	Emission Inventory Term	Examples
Point	Stationary – Point Facilities	Stacks at Individual Facilities
Area	Off-Road Mobile	Construction Equipment, Farm Equipment, Trains, Recreational Boats
Area	Area-wide	Residential Fuel Combustion, Livestock Waste, Consumer Products, Architectural Coatings
Area	Stationary - Aggregated	Industrial Fuel Use
On-Road Motor Vehicles	On-Road Mobile	Cars and Trucks
Biogenic	Biogenic	Trees

The following sections describe in more detail the temporal, spatial and chemical disaggregation of the emissions inventory for point sources and area sources.

## 2.2. Temporal Distribution of Emissions

Emission inventories that are temporally and spatially resolved are needed for modeling purposes, for the base case and baseline modeling inventories as well as future year inventories. The temporal distribution of on-road emissions and biogenic emissions are discussed in Sections 3.4 and 3.5, respectively. How emissions are temporally

distributed for the remaining sources (point, area and off-road mobile sources) is discussed below.

Emissions are adjusted temporally to represent variations by month, day of week and hour of day. Temporal data are stored in ARB's emission inventory database. Each local air district assigns temporal data for all processes at each facility in their district to represent when emissions at each process occur. For example, emissions from degreasing may operate differently than a boiler. ARB or district staff also assigns temporal data for each area source category by county/air basin/district.

**2.2.1. Monthly Variation:** Emissions are adjusted temporally to represent variations by month. Some emission sources operate the same throughout a year. For example, a process heater at a refinery or a line haul locomotive likely operates the same month to month. Other emission categories, such as a tomato processing plant or use of recreational boats, vary significantly by season. ARB's emission inventory database stores the relative monthly fractional activity for each process, the sum of which is 100. Using an example of emission sources that typically operate the same over each season, emissions from refinery heaters and line haul locomotives would have a monthly fraction (throughput) of 8.33 for each month (calculated as  $100/12 = 8.33$ ). This is considered a flat monthly profile. To apply monthly variations to create a gridded inventory, the annual average day's emissions (yearly emissions divided by 365) is multiplied by the typical monthly throughput. For example, a typical monthly throughput in July for recreational boats of 15 results in about 1.8 times higher ( $15 / 8.33 = 1.8$ ) emissions than a day with flat monthly profile.

**2.2.2. Weekly Variation:** Emissions are adjusted temporally to represent variations by day of week. Some operations are the same over a week, such as a utility boiler or a landfill. Many businesses operate only 5 days per week. Other emissions sources are similar on weekdays, but may operate differently on weekend days, such as architectural coatings or off-road motorcycles. To

accommodate variations in days of the week, each process or emission category is assigned a days per week code or DPWK. Table 3 below shows the current DPWK codes and Table 11 in Appendix D shows additional DPWK codes used for agricultural-related emissions.

Table 3 Day of week variation factors

Code	WEEKLY CYCLE CODE DESCRIPTION	M	T	W	TH	F	S	S
1	One day per week	1	1	1	1	1	0	0
2	Two days per week	1	1	1	1	1	0	0
3	Three days per week	1	1	1	1	1	0	0
4	Four days per week	1	1	1	1	1	0	0
5	Five days per week - Uniform activity on week days; non on Saturday and Sunday	1	1	1	1	1	0	0
6	Six days per week - Uniform activity on week days; non on Saturday and Sunday	1	1	1	1	1	1	0
7	Seven days per week - Uniform activity every day Of the week	1	1	1	1	1	1	1
20	Uniform activity on Saturday and Sunday; No activity the remainder of the week	0	0	0	0	0	1	1
21	Uniform activity on Saturday and Sunday; No activity the remainder of the week	5	5	5	5	5	10	10
22	Uniform activity on week days; Reduced activity on weekends	10	10	10	10	10	7	4
23	Uniform activity on week days; Reduced activity on weekends (For onroad motor vehicles)	10	10	10	10	10	8	8
24	Uniform activity on week days; half as much activity on Saturday. Little activity on Sunday	10	10	10	10	10	5	1
25	Uniform activity on week days; one third as much on Saturday; little on Sunday	10	10	10	10	10	3	1
26	Uniform activity on week days; little activity on Saturday; no activity on Sunday	10	10	10	10	10	3	0
27	Uniform activity on week days; half as much activity on weekends	10	10	10	10	10	5	5
28	Uniform activity on week days; Five times as much activity on weekends	2	2	2	2	2	10	10
29	Uniform activity on Monday through Thursday; increased activity on Friday, Saturday, Sunday	8	8	8	8	10	10	10

**2.2.3. Daily Variation:** Emissions are adjusted temporally to represent variations by hour of day. Many emission sources occur 24 hours per day, such as livestock waste or a sewage treatment plant. Many businesses operate 8 hours per day. Other emissions sources vary significantly over a day, such as residential space heating or pesticide application. Each process or emission category is assigned an hours per day code or HPDY. Table 4 below shows the daily variation factors or current HPDY codes. Table 12 in Appendix D shows additional DPWK codes used for agricultural-related emissions.



Table 4 Daily variation factors

Code	CODE DESCRIPTION	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	11 HOUR PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
2	22 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
3	33 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
4	44 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
5	55 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
6	66 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
7	77 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
8	88 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
9	99 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
10	1010 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
11	1111 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
12	1212 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
13	1313 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
14	1414 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
15	1515 HOURS PER DAY	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
16	1616 HOURS PER DAY - UNIFORM ACTIVITY FROM 8 A.M. TO MIDNIGHT (2 WORKING SHIFTS)	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
17	1717 HOURS PER DAY	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
18	1818 HOURS PER DAY	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
19	1919 HOURS PER DAY	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
20	2020 HOURS PER DAY	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
21	2121 HOURS PER DAY	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
22	2222 HOURS PER DAY	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
23	2323 HOURS PER DAY	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
24	2424 HOURS PER DAY - UNIFORM ACTIVITY DURING THE DAY	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
31	31 MAJOR ACTIVITY 5-9 P.M. - AVERAGE DURING DAY. MINIMAL IN EARLY A.M. (GAS STATIONS)	3	1	1	1	1	1	1	1	1	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
33	33 MAX ACTIVITY 7-9 A.M. & 7-11 P.M. - AVERAGE DURING DAY. LOW AT NIGHT (RESIDENTIAL FUEL COMBUSTION)	2	2	2	2	2	2	2	2	2	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
34	34 ACTIVITY 1 TO 9 A.M.; NO ACTIVITY REMAINDER OF DAY (i.e. ORCHARD HEATERS)	0	8	8	8	8	10	10	10	10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
35	35 MAX ACTIVITY 7 A.M. TO 1 A.M.; REMAINDER IS LOW (i.e. COMMERCIAL AIRCRAFT)	10	1	1	1	1	1	1	1	1	8	10	10	10	10	10	10	10	10	10	10	10	10	10	10
37	37 ACTIVITY DURING DAYTIME HOURS; LESS CHANGE IN EARLY MORNING AND LATE EVENING	0	0	0	0	0	0	0	0	0	1	3	6	9	10	10	10	10	10	10	10	10	10	10	10
38	38 ACTIVITY DURING MEAL TIME HOURS (i.e. RESIDENTIAL COOKING)	0	0	0	0	0	0	0	0	0	2	6	6	2	2	4	4	2	1	1	3	10	8	7	6
50	50 PEAK ACTIVITY AT 7 A.M. & 4 P.M.; AVERAGE DURING DAY (ON-ROAD MOTOR VEHICLES)	1	1	1	1	1	1	1	1	1	10	6	5	5	5	5	6	10	8	6	4	1	1	1	1
51	51 ACTIVITY FROM 6 A.M. TO 12 P.M. (PETROLEUM DRY CLEANING)	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
52	52 MAJOR ACTIVITY FROM 6 A.M.-12 P.M.; LESS FROM 12-7 P.M. (PESTICIDES)	0	0	0	0	0	0	0	0	0	1	6	10	10	10	6	3	3	3	4	0	0	0	0	0
53	53 ACTIVITY FROM 7 A.M. TO 12 P.M. (AGRICULTURAL AIRCRAFT)	0	0	0	0	0	0	0	0	0	2	2	2	2	2	1	0	0	0	0	0	0	0	0	0
54	54 UNIFORM ACTIVITY FROM 7 A.M. TO 9 P.M. (DAYTIME BIOGENICS)	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
55	55 UNIFORM ACTIVITY FROM 9 P.M. TO 7 A.M. (NIGHTTIME BIOGENICS)	1	1	1	1	1	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
56	56 MAX ACTIVITY 8 A.M. TO 5 P.M. - MINIMAL AT NIGHT & EARLY MORNING (CAN&COIL/METAL PARTS COATINGS)	0	0	0	0	0	0	0	0	0	1	2	3	10	10	10	10	10	10	10	10	10	10	10	10
57	57 MAX ACTIVITY 7 A.M. TO 2 P.M. - MINIMAL AT NIGHT & EARLY MORNING (CONSTRUCTION EQUIPMENT ON HOT)	0	0	0	0	0	0	0	0	0	1	6	10	10	10	10	8	4	2	1	1	1	1	1	1
58	58 MAX ACTIVITY 7 A.M. TO NOON; REDUCED ACTIVITY FROM NOON TO 6 P.M. (AUTO REFINISHING)	0	0	0	0	0	0	0	0	0	0	10	10	10	10	8	8	8	8	8	8	8	8	8	8
59	59 MAXIMUM ACTIVITY FROM 7:00 AM TO 3:00 PM; REDUCED ACTIVITY FROM 3:00 TO 6:00 PM (CONSTRUCTION)	0	0	0	0	0	0	0	0	0	2	10	10	10	10	10	10	10	10	10	10	10	10	10	10
60	60 MAXIMUM ACTIVITY FROM NOON TO 7:00 PM; REDUCED ACTIVITY EVENING AND MORNING HOURS (RECREATIONAL)	0	0	0	0	0	0	0	0	0	2	4	6	7	9	10	10	10	10	10	10	10	10	10	10
81	81 MAX ACTIVITY 9 AM TO 3 PM; HALF THE ACTIVITY REMAINING HOURS (WASTE FROM DAIRY CATTLE)	7	6	5	4	4	4	5	7	8	9	10	10	7	3	3	4	4	5	6	7	8	9	10	10
82	82 ACTIVITY FROM 10 AM TO 9 PM RISING TO PEAK AT 3; NO ACTIVITY REMAINDER OF DAY (WASTE FROM POULTRY)	0	0	0	0	0	0	0	0	0	0	3	7	7	10	10	7	3	3	3	3	3	3	3	3
83	83 ACTIVITY FROM 9 AM TO 12 AM RISING TO PEAK AT 3; MINIMUM ACTIVITY REMAINDER OF DAY (WASTE FROM SWINE)	0	0	0	0	0	0	0	0	0	0	1	2	4	6	8	9	10	8	4	3	2	1	1	1
84	84 MAJOR ACTIVITY FROM 11 AM TO 8 PM; REDUCED OTHER HOURS (EVAP-COASTAL COUNTIES)	7	7	6	6	6	6	7	8	8	9	10	10	10	10	9	8	7	7	7	7	7	7	7	7
85	85 MAJOR ACTIVITY FROM 11 AM TO 8 PM; REDUCED OTHER HOURS (EVAP-NON-COASTAL COUNTIES)	5	5	5	5	4	4	5	5	6	7	8	9	9	10	10	10	10	9	8	7	6	6	6	6

### **2.3. Spatial Allocation**

Once the base case, baseline or future year inventories are developed, the next step of modeling inventory development is to spatially allocate the emissions. Air quality modeling attempts to replicate the physical and chemical processes that occur in an inventory domain. Therefore, it is important that the physical location of emissions be specified as accurately as possible. Ideally, the actual location of all emissions would be known exactly. In reality, however, some categories of emissions would be virtually impossible to determine – for example, the actual amount and location of consumer products (e.g. deodorant) used every day. To the extent possible, the spatial allocation of emissions in a modeling inventory approximates as closely as possible the actual location of emissions.

Spatial allocation is typically accomplished by using spatial surrogates. These spatial surrogates are processed into spatial allocation factors in order to geographically distribute county-wide area source emissions to individual grid cells. Spatial surrogates are developed based on demographic, land cover and other data that exhibit patterns which vary geographically. The spatial surrogates have been updated over the years mainly by Sonoma Technology, Inc. (STI) (Funk, et al., 2001) who created a 2000 base year and various future years. Later, STI updated the underlying spatial data and developed new surrogates (Reid, et al., 2006) completing the project in 2008. ARB and districts have continued to update and improve many of the spatial surrogates and added new ones.

Three basic types of surrogate data were used to develop the original spatial allocation factors: land use and land cover; facility location; and demographic and socioeconomic data. Land use and land cover data are associated with specific land uses, such as agricultural harvesting or recreational boats. Facility locations are used for sources such as gas stations and dry cleaners. Demographic and socioeconomic data, such as population and housing, are associated with residential, industrial, and commercial activity (e.g. residential fuel combustion). To develop spatial allocation factors of high quality and resolution, local socioeconomic and demographic data were used where available for developing base case, baseline and future year inventories. These data

were available from local Metropolitan Planning Organizations (MPO) or Regional Transportation Planning Agencies (RTPA), where they are used as inputs for travel demand models. In rural regions for which local data were not available, data from Caltrans' Statewide Transportation Model were used.

Since 2008, ARB and district staffs have continued to search for more recent or improved sources of data, since the underlying data used by STI were pre-recession. ARB and district staffs have updated many of the spatial surrogates and added many new ones.

- Updates to land use categories were made using the National Land Cover Database 2011 (Homer, et al., 2015).
- Many surrogates were updated using the locations from Dun & Bradstreet's Market Insight Database (Dun and Bradstreet, 2015). The types of sources were defined by SIC (Standard Industrial Classification). Fourteen new surrogates were developed for industrial-related sources using SIC and whether manufacturing occurred at the facility.
- U.S. Census American Community Survey (FactFinder, 2011) data by census block were used to update residential fuel use.
- Sierra Research developed nine new surrogates related to agricultural activities (Anderson, et al., 2012), some of which incorporated crop-specific factors.
- Seven new surrogates were developed using vessel traffic data, or Automatic Identification System (AIS) data, collected by the U.S. Coast Guard.
- A new surrogate was created to represent the location of construction equipment. The distribution is a combination of two sets of data: 90% change in "imperviousness" between 2006 and 2011 from NLCD 2011 and 10% road network. Impervious surfaces are mainly artificial structures such as pavements (roads, sidewalks, driveways and parking lots) that are covered by materials impenetrable to a satellite such as asphalt, concrete, brick, stone and rooftops.
- A new surrogate was compiled to distribute emissions from transport refrigeration units (TRU) from three sources: 65% distribution centers, 34% road network and 1% grocery stores / food processing facilities. Information on

distribution centers were retrieved from ARBER, the ARB Equipment Registration software for the Transport Refrigeration Unit (TRU) ATCM and the Drayage Truck Regulation.

In all, a total of 99 unique surrogates are available for use. A summary of the spatial surrogates for which spatial allocation factors were developed is shown below in Table 5.

Table 5 Spatial Surrogates

Surrogate Name	Surrogate Definition
AEROSPACE	Spatial distribution of businesses involved in aerospace
Airports	Spatial locations of all airports
All_PavedRds	Spatial distribution of road network (all paved roads)
AutobodyShops	Locations of autobody repair and refinishing shops
CAFO	Spatial distribution of concentrated animal feeding operations
CANCOIL	Spatial distribution of businesses involved in can and coil operations
Cemeteries	Spatial locations of cemeteries
Comm_Airports	Spatial locations of commercial airports
COMPOST	Spatial distribution of composting
CONSTRUCTION_EQUIP	Spatial distribution of where construction equipment is used
Devplnd_HiDensity	Spatial distribution of developed land - low density, medium density and high density
Devplnd_LoDensity	Spatial distribution of developed land - open space (lowest density)
DREDGE	Locations of dredging
Drycleaners	Locations of dry cleaning facilities
DryLakeBeds	Locations of dry lake beds
Elev5000ft	Topological contours – areas above 5000 feet
Employ_Roads	Spatial distribution of total employment and road density (all paved roads)
FABRIC	Spatial distribution of businesses involved in fabric manufacturing
FERRIES	Locations of ferry ports and routes
FISHING_COMM	Locations of commercial fishing
Forestland	Spatial distribution of forest land
Fugitive_Dust	Spatial distribution of barren land
GAS_DISTRIBUTION	Location of gas pipelines
GAS_SEEP	Location of natural-occurring gas seeps
GasStations	Locations of gasoline service stations
GASWELL	Locations of gas wells
GolfCourses	Spatial locations of golf courses
HE_Sqft	Computed surrogate based on housing and employment (est. ft <sup>2</sup> / person)
Hospitals	Spatial locations of hospitals
Housing	Spatial distribution of total housing
Housing_Autobody	Spatial distribution of housing and autobody refinishing shops
Housing_Com_Emp	Spatial distribution of total housing and commercial employment
Housing_Restaurants	Spatial distribution of total housing and restaurants/bakeries
Surrogate Name	Surrogate Definition
INDUSTRIAL	Spatial distribution of industrial businesses where manufacturing occurs (SIC<4000)
Industrial_Emp	Spatial distribution of industrial employment
InlandShippingLanes	Spatial distribution of major shipping lanes within bays and inland areas
Irr_Cropland	Spatial location of agricultural cropland
Lakes_Coastline	Locations of lakes, reservoirs, and coastline
LAKES_RIVERS_RECBOAT	Locations of lakes, rivers and reservoirs where recreational boats are used
LANDFILLS	Locations of landfills
LANDPREP	Spatial distribution of dust from land preparation operations (e.g. tilling)
LINEHAUL	Spatial distribution of Class I rail network
LiveStock	Spatial distribution of cattle ranches, feedlots, dairies, and poultry farms
MARINE	Spatial distribution of businesses involved in marine

Surrogate Name	Surrogate Definition
METALFURN	Spatial distribution of businesses involved in metal furniture
METALPARTS	Spatial distribution of businesses involved in metal parts and products
Metrolink_Lines	Spatial distribution of metrolink network
MILITARY_AIRCRAFT	Locations of landing strips on military bases
MILITARY_SHIPS	Locations of military ship activity
MILITARY_TACTICAL	Military bases where tactical equipment are used
MilitaryBases	Locations of military bases
NON_PASTURE_AG	Spatial distribution of farmland
NonIrr_Pastureland	Spatial location of pasture land
NonRes_Chg	Computed surrogate based on spatial distribution of non-residential areas
OCEAN_RECBOAT	Locations of recreational boat activity that can occur on the ocean and SF Bay
OIL_SEEP	Location of naturally-occurring oil seeps
OILWELL	Locations of oil wells (both onshore and offshore)
OTHERCOAT	Spatial distribution of businesses with SIC<4000 not included in another category
PAPER	Spatial distribution of businesses involved in paper
PASTURE	Spatial distribution of grazing land
PEST_ME_BR	Spatial distribution of methyl bromide pesticides
PEST_NO_ME_BR	Spatial distribution of non-methyl bromide pesticides
PLASTIC	Spatial distribution of businesses involved in plastic
Pop_ComEmp_Hos	Spatial distribution of hospitals, population and commercial employment
Population	Spatial distribution of population
Ports	Locations of shipping ports
POTWs	Coordinate locations of POTWs
PrimaryRoads	Spatial distribution of road network (primary roads)
PRINT	Spatial distribution of print businesses
Raillines	Spatial distribution of railroad network
RailYards	Locations of rail yards
Rds_HE	Calculated surrogate based on road densities and housing/employment (est. ft <sup>2</sup> / person)
RefineriesTankFarms	Coordinate locations of refineries and tank farms
Res_NonRes_Chg	Computed surrogate based on spatial distribution of residential and non-residential areas
ResGasHeating	Spatial distribution of homes using gas supplied by a utility as primary source of heating
Residential_Chg	Computed surrogate based on spatial distribution of residential areas
ResLPGHeat	Spatial distribution of homes using gas (bottled, tank or LP) as primary source of heating
ResNonResChg_IndEmp	Spatial distribution of industrial employment and residential/non-residential change
ResOilHeat	Spatial distribution of homes using fuel oil or kerosene as primary source of heating
Restaurants	Locations of restaurants
ResWoodHeating	Spatial distribution of homes using wood as primary source of heating
Surrogate Name	Surrogate Definition
SandandGravelMines	Locations of sand/gravel excavation and mining
Schools	Spatial locations of schools
SecondaryPavedRds	Spatial distribution of road network (secondary roads)
SEMICONDUCT	Spatial distribution of businesses involved in semiconductors
Ser_ComEmp_Sch_GolfC_Cem	Spatial distribution of service and commercial employment, schools, cemeteries, of courses
Service_Com_Emp	Spatial distribution of service and commercial employment
Shiplanes	Spatial distribution of major shipping lanes
SILAGE	Spatial distribution of silage operations
SingleHousingUnits	Spatial distribution of single dwelling units
TRU	Spatial distribution of transport refrigeration units
TUG_TOW	Spatial distribution of tug and tow boats
UnpavedRds	Spatial distribution of road network (unpaved roads)
Wineries	Locations of wineries
WOOD	Spatial distribution of businesses using wood
WOODFURN	Spatial distribution of businesses involved in wood furniture

The following sections describe in more detail the type of spatial disaggregation used for each sector of the emissions inventory.

**2.3.1. Spatial Allocation of Area Sources:** Each area source category is assigned a spatial surrogate that is used to allocate emissions to a grid cell in ARB's 4km statewide modeling domain. Examples of surrogates include population, land use, and other data with known geographic distributions for allocating emissions to grid cells, as described above.

**2.3.2. Spatial Allocation of Point Sources:** Each point source is allocated to grid cells using the latitude and longitude reported for each stack. If there are no stack latitude and longitude, the facility coordinates are used. There are two types of point sources: elevated and non-elevated sources. Vertical distribution of elevated sources is allocated using the plume rise algorithm in the emissions processor, SMOKE (see Section 3.3), while non-elevated are allocated to the first layer. Most stationary point sources with existing stacks are regarded as elevated sources. Those without physical stacks that provide only latitude/longitude, such as airports or landfills, are considered non-elevated.

**2.3.3. Spatial Allocation of Wildfires, Prescribed Burns and Wildland Fire Use:** Emissions from these sources are event and location-based. A fire event can last a few hours or span multiple days. Each fire is spatially allocated to grid cells using the extent of each fire event, while the temporal distribution also reflects the actual duration of the fire. The spatial information to allocate the fire emissions comes from a statewide interagency fire perimeters geodatabase maintained by the Fire and Resource Assessment Program (FRAP) of the California Department of Forestry and Fire Protection (CALFIRE). More details on the methodology and estimation of the wildfire emissions can be found in Section 3.6.1.

**2.3.4. Spatial Allocation of Ocean going vessels (OGV):** Ship emissions are allocated to the grids corresponding to the vessel traffic lanes in ARB's OGV

model (ARB-PTSD, 2011) These traffic lanes were estimated from three different sources:

- a. National Waterway Network
- b. The Ship Traffic, Energy and Environment Model
- c. Automated instrumentation system (AIS) telemetry data collected in 2007

**2.3.5. Spatial Allocation of On-road Motor Vehicles:** The spatial allocation of on-road motor vehicles is based on DTIM as described in Section 3.4.

**2.3.6. Spatial Allocation of Biogenic Emissions:** As described in Section 3.5, gridded biogenic emissions are derived using the Model of Emissions of Gases and Aerosols from Nature (MEGAN). MEGAN utilizes gridded emission factor and plant functional type data, adjusted by local meteorological conditions and satellite derived leaf area data, to estimate hourly biogenic emissions within each grid cell of the modeling domain. More details about MEGAN can be found at <http://lar.wsu.edu/megan/>.

## 2.4. Speciation Profiles

ARB's emission inventory lists the amount of pollutants discharged into the atmosphere by source in a certain geographical area during a given time period. It currently contains estimates for CO, NH<sub>3</sub>, NO<sub>x</sub>, SO<sub>x</sub>, total organic gases (TOG) and particulate matter (PM). CO and NH<sub>3</sub> are single species; NO<sub>x</sub> emissions are composed of NO, NO<sub>2</sub> and HONO; and SO<sub>x</sub> emissions are composed of SO<sub>2</sub> and SO<sub>3</sub>. Emissions of TOG and PM for many sources can actually contain over hundreds of different chemical species, and speciation is the process of disaggregating these inventory pollutants into individual chemical species components or groups of species. ARB maintains and updates such species profiles for organic gases (OG) and PM for a variety of source categories.

Photochemical models simulate the physical and chemical processes in the lower atmosphere, and include all emissions of the important classes of chemicals involved in

photochemistry. Organic gases emitted to the atmosphere are referred to as Total Organic Gas or TOG. TOG includes all organic compounds that can become airborne (through evaporation, sublimation, as aerosols, etc.), excluding carbon monoxide, carbon dioxide, carbonic acid, metallic carbides or carbonates and ammonium carbonate. TOG emissions reported in the ARB's emission inventory are the basis for deriving the Reactive Organic Gas (ROG) emission components, which are also reported in the inventory. ROG is defined as TOG minus ARB's exempt compounds (e.g., methane, ethane, various chlorinated fluorocarbons, acetone, perchloroethylene, volatile methyl siloxanes, etc.). ROG is nearly identical to U.S. EPA's Volatile Organic Compounds (VOC), which is based on EPA's exempt list. For all practical purposes, use of the term ROG and VOC are interchangeable. Also, various regulatory uses of the term VOC, such as that for consumer products exclude specific, additional compounds from particular control requirements.

The OG speciation profiles are applied to estimate the amounts of various organic compounds that make up TOG emissions. A speciation profile contains a list of organic compounds and the weight fraction that each compound comprises of the TOG emissions from a particular source type. In addition to the chemical name for each chemical constituent, the file also shows the chemical code (a 5-digit ARB internal identifier). The speciation profiles are applied to TOG to develop both the photochemical model inputs and the emission inventory for ROG. It should be noted that districts are allowed to report their own reactive fraction of TOG that is used to calculate ROG rather than use the information from the assigned organic gas speciation profiles. These district-reported fractions are not used in developing modeling inventories because the information needed to calculate the amount of each organic compound is not available.

The PM emissions are size fractionated by using PM size profiles, which contain the total weight fraction for PM<sub>2.5</sub> and PM<sub>10</sub> out of total PM. The fine and coarse PM chemical compositions are characterized by applying the PM chemical speciation profiles for each source type, which contain the weight fractions of each chemical



species for PM<sub>2.5</sub>, PM<sub>10</sub> and total PM. PM chemical speciation profiles may also vary for different PM size fractions even for the same emission source. PM size profiles and speciation profiles are typically generated based on source testing data. In most previous source testing studies aimed at determining PM chemical composition, filter-based sampling techniques were used to collect PM samples for chemical analyses.

The organic gas profiles and PM profiles used in the emission inventory are available for download from the ARB's web site at: <http://www.arb.ca.gov/ei/speciate/speciate.htm>

Each process or product category is keyed to one of the OG profiles and one of the PM profiles. Also available for download from ARB's web site is a cross-reference file that indicates which OG profile and PM profile are assigned to each category in the inventory. The inventory source categories are represented by an 8-digit source classification code (SCC) for point sources, or a 14-digit emission inventory code (EIC) for area and mobile sources. Some of the organic gas profiles and PM profiles related to motor vehicles, ocean going vessels, and fuel evaporative sources vary by the inventory year of interest, due to changes in fuel composition, vehicle fleet composition and diesel particulate filter (DPF) requirements over time. Details can be found in ARB's documentation of heavy-duty diesel vehicle exhaust PM speciation profiles (ARB, 2011).

Research studies are conducted regularly to improve ARB's speciation profiles. These profiles support ozone and PM modeling studies but are also designed to be used for aerosol and regional toxics modeling. The profiles are also used to support other health or welfare related modeling studies where the compounds of interest cannot always be anticipated. Therefore, speciation profiles need to be as complete and accurate as possible. ARB has an ongoing effort to update speciation profiles as data become available, such as the testing of emission sources or surveys of product formulations. New speciation data generally undergo technical and peer review, and updating of the profiles is coordinated with users of the data. The recent addition to ARB's speciation profiles include:

- (1) Organic gas profile
  - Consumer products
  - Architectural coating
  - Gasoline fuel and headspace vapor
  - Gasoline vehicle hot soak and diurnal evaporation
  - Gasoline vehicle start and running exhaust
  - Silage
  - Aircraft exhaust
  - Compressed Natural Gas (CNG) bus running exhaust
  
- (2) PM profile
  - Gasoline vehicle exhaust
  - On-road diesel exhaust
  - Off-road diesel exhaust
  - Ocean going vessel exhaust
  - Aircraft exhaust
  - Concrete batching
  - Commercial cooking
  - Residential fuel combustion-natural gas
  - Coating/painting
  - Cotton ginning
  - Stationary combustion

### **3. Methodology for Developing Base Case and Baseline Emissions Inventories**

As mentioned in Section 0, the base case and baseline inventories include temperature, humidity and solar insolation effects for some emission categories; development of these data is described in Sections 3.1 and 3.2. The remaining sections of Chapter 3 detail how the base case and baseline inventories were created for different sectors of the inventory, such as for point, area, on-road motor vehicles, biogenic and other day-specific sources.

### 3.1. Surface Temperature and Relative Humidity Fields

The calculation of gridded emissions for some categories of the emissions inventory is dependent on meteorological variables. More specifically, biogenic emissions are sensitive to air temperatures and solar radiation while emissions from on-road mobile sources are sensitive to air temperature and relative humidity. As a result, estimates of air temperature (T), relative humidity (RH), and solar radiation are needed for each grid cell in the modeling domain in order to take into account the effects of these meteorological variables.

Gridded temperature and humidity fields are readily available from prognostic meteorological models such as the Weather Research and Forecasting (WRF) model (<http://www.wrf-model.org/index.php>), which is used to prepare meteorological inputs for the air quality model. However, prognostic meteorological models can at times have difficulty capturing diurnal temperature extremes (Valade, 2009; Caldwell, 2009; Fovell, 2008). Since temperature and the corresponding relative humidity extremes can have an appreciable influence on some emissions categories, such as on-road mobile and biogenic sources, measurement based fields for these parameters are used in processing emissions. The CALMET (<http://www.src.com/>) diagnostic meteorological model is utilized to generate both the gridded temperature and relative humidity fields used in processing emissions. The solar radiation fields needed for biogenic emission inventory calculations were taken from the WRF prognostic model, which is also used to generate meteorology for the air quality model. The principal steps involved in generating a gridded, surface-level temperature field using CALMET include the following:

1. Compute the relative weights of each surface observation station to each grid cell (the weight is inversely proportional to the distance between the surface observation station and grid cell center).
2. Adjust all surface temperatures to sea level. In this step, a lapse rate of  $-0.0049$  °C/m is used (this lapse rate is based on private communication with Gary Moore of Earth Tech, Inc., Concord, MA). This lapse rate ( $=2.7$  F/1000 feet) is based on observational data.

3. Use the weights to compute a spatially-averaged sea-level temperature for each grid cell.
4. Correct all sea-level temperatures back to 10 m height above ground level (i.e. the standard height of surface temperature measurements) using the lapse rate of  $-0.0049\text{ }^{\circ}\text{C/m}$  again.
5. The current version of CALMET does not generate estimates of relative humidity. As a result, a post-processing program was used to produce gridded, hourly relative humidity estimates from observed relative humidity data. The major steps needed to generate gridded, surface-level relative humidity are described as follows:
  - a. Calculate actual vapor pressure from observed relative humidity and temperature at all meteorological stations. The (Mc. Rae, 1980) method is used to calculate the saturated vapor pressure from temperature;
  - b. Compute the relative weights of each surface observation station to each grid in question, exactly as done by CALMET to compute the temperature field;
  - c. Use the weights from step 2 to compute a spatially-averaged estimate of actual vapor pressure in each grid cell;
  - d. For each grid cell, calculate relative humidity from values for actual vapor pressure and temperature for the same grid cell.

### **3.2. Insolation Effects**

Insolation data was used in the estimation of the gridded emissions inventory and provided by the WRF meteorological fields as mentioned in Section 3.5.

### **3.3. Estimation of Gridded Area and Point sources**

Emissions inventories that are temporally, chemically, and spatially resolved are needed as inputs for the photochemical air quality model. Point sources and area sources (area-wide, off-road mobile and aggregated stationary) are processed into emissions inventories for photochemical modeling using the SMOKE (Sparse Matrix Operator

Kernel Emissions) modeling system (<https://www.cmascenter.org/smoke/>).

Improvements to SMOKE were recently implemented under ARB contract for version 4.0 of SMOKE (Baek, 2015).

Inputs for SMOKE are annual emissions totals from CEPAM and information for allocating to temporal, chemical, and spatial resolutions. Temporal inputs for SMOKE are screened for missing or invalid temporal codes as discussed in Section 4.1. Temporal allocation of emissions using SMOKE involves the disaggregation of annual emissions totals into monthly, day of week, and hour of day emissions totals. The temporal codes from Table 3 and Table 4 are reformatted into an input-ready format as explained in the SMOKE user's manual. Chemical speciation profiles, as described in Section 2.4, and emissions source cross-reference files used as inputs for SMOKE are developed by ARB staff. SMOKE uses the files for the chemical speciation of NO<sub>x</sub>, SO<sub>x</sub>, TOG and PM to species needed by photochemical air quality models.

Emissions for area sources are allocated to grid cells as defined by the modeling grid domain defined in Section 1.4. Emissions are spatially disaggregated by the use of spatial surrogates as described in Section 2.3. These spatial surrogates are converted to a SMOKE-ready format as described in the SMOKE user's manual. Emissions for point sources are allocated to grid cells by SMOKE using the latitude and longitude coordinates reported for each stack.

### **3.4. Estimation of On-road Motor Vehicle Emissions**

The EMFAC emissions model is used by ARB to assess emissions from on-road vehicles including cars, trucks, and buses in California, and to support air quality planning efforts to meet the Federal Highway Administration's transportation planning requirements. EMFAC is designed to produce county-level, average-day estimates. As a result, these estimates must be disaggregated spatially and temporally into gridded, hourly estimates for air quality modeling.

The general methodology used to disaggregate EMFAC emission estimates is a two-step approach. The first step uses the Direct Travel Impact Model (DTIM4) (Systems Applications Inc., 2001) to produce gridded, hourly emission estimates. The second

step distributes EMFAC emissions according to the spatiotemporal output from DTIM. This methodology has been peer reviewed by the Institute of Transportation Studies at the University of California, Irvine, under CCOS contract 11-4CCOS.

The spatiotemporal allocation of emissions from DTIM does not vary dramatically with small changes in meteorological data (T/RH), resulting in a negligible monthly variation of the spatial surrogate. However, differences in DTIM's winter versus summer spatiotemporal allocation are slightly appreciable. Therefore, spatial surrogates are created for a winter and a summer day.

The most recent version of EMFAC, EMFAC2014, has three separate modules that are relevant for the preparation of the on-road emissions gridded inventory: one that estimates emissions, one that estimates emission rates, and one that estimates activity data. The emissions module is run for every county and every day of the modeled year using day-specific temperature and relative humidity. On a less granular level, the emissions rates module is run for every county for a summer day and a winter day. Lastly, the activity module is run once to estimate vehicle miles traveled (VMT), number of vehicle trips, fuel consumption, and the number of vehicles in use.

**3.4.1. General Methodology:** Mobile source emissions are sensitive to ambient temperature and humidity. Both EMFAC and DTIM account for meteorological effects using day-specific inputs. For EMFAC, hourly gridded temperature and humidity fields are averaged by county using a gridded VMT weighted average (i.e. weighted proportional to the VMT per grid cell in a county). DTIM accepts gridded, hourly data directly (CALMET formatted data). See Section 3.1 for more information.

EMFAC provides vehicle-class-specific emissions estimates for: exhaust, evaporative, tire wear, and brake wear emissions. EMFAC also produces estimates of: VMT, number of vehicle trips, fuel consumption, and the number of vehicles in use. More information on EMFAC can be found at (ARB-MSEI, 2015) . The vehicle activity is the most important input for spatiotemporal

distribution of emissions. DTIM uses hourly vehicle miles traveled on each highway link and each of the vehicle trips in the modeling domain. The detailed vehicle activity data is obtained from ARB's Integrated Transportation Network (dtiv3) database.

The overall processing of on-road emissions to create the gridded emissions inventory can be seen in Figure . Activity data from the ITN (see Section 3.4.2) is developed for the thirteen EMFAC 2007 vehicle types, but activity is split for gas and diesel, resulting in a total of 26 vehicle types as shown in the block diagram. The forecasted on-road modeling inventories are developed using the same methodology as the baseline year, where future year emissions are based on running EMFAC 2014 in Emissions Mode for the associated future year.




Figure 2 Block diagram for on-road processing



**3.4.2. ITN Activity Data:** The ITN is a database which is populated with link-based and Traffic Analysis Zone (TAZ)-based travel activity from travel demand models provided by different metropolitan planning organizations (MPOs), California Department of Transportation (Caltrans) and other California regional transportation planning agencies. The vintage and types of data used in the current version of the ITN are shown in Table 6. Different types of quality control parameters like vehicle mix, hourly distributions and post-mile coverage are obtained from default EMFAC and Caltrans databases. After these various pieces of data are imported to the database, the data can be examined for quality assurance. These input data sets are later moved into consolidated and geographically referenced master tables of link and TAZ activity data. Finally, these master tables are processed to produce hourly tables and hourly activity data input files for DTIM.

Table 6 Vintage of travel demand models for link based and traffic analysis zone

<b>Metropolitan Planning Organizations</b>	<b>TDM Version Base year</b>	<b>Data types received</b>	<b>Data received on</b>
AMBAG	2010	Links, Trips	06/15/2015
BCAG	2010	Links, Trips	05/13/2015
FCOG	2008	Links†	06/11/2015
CALTRANS	2010	Links, Trips	12/09/2014
KCOG	2010	Links†	06/11/2015
KCAG	2010	Links†	06/11/2015
MTC	2010	Links, Trips	03/23/2015
MCTC	2010	Links†	06/11/2015
MCAG	2010	Links, Trips	06/11/2015
SACOG	2010	Links, Trips	05/08/2014
SANDAG	2008	Links, Trips	12/09/2014
SBCAG	2010	Links, Trips	04/06/2015
SCAG	2008	Links, Trips	01/23/2014
SJCOG	2010	Links, Trips	06/11/2015
SLOCOG	2010	Links, Trips	12/19/2014
StanCOG	2010	Links, Trips	06/11/2015
SCRTPA	2010	Links, Trips	07/13/2015
TCAG	2010	Links†	06/11/2015
TMPO	2010	Links, Trips	04/02/2015

† Trips data from Caltrans Statewide Travel Demand model were used

**3.4.3. Spatial Adjustment:** The spatial allocation of county-wide EMFAC emissions is accomplished using gridded, hourly emission estimates from DTIM normalized by county. DTIM uses emission rates from EMFAC along with activity data, digitized roadway segments (links) and traffic analysis zone centroids to calculate gridded, hourly emissions for travel and trip ends. DTIM considers fewer vehicle categories than EMFAC outputs; therefore a mapping between EMFAC and DTIM vehicle categories is necessary. Categories of emissions after running DTIM are presented in Table 7. The categories are represented by the listed source classification codes (SCC) developed by ARB and depend on vehicle type, technology, and whether the vehicle is catalyst, non-catalyst, or diesel. Light- and medium-duty vehicles are separated from heavy-duty vehicles to allow for separate reporting and control strategy applications.

Table 7 DTIM Emission Categories

SCC for light and medium duty gas vehicles	SCC for heavy-duty gas vehicles	SCC for light-duty and medium-duty diesel vehicles	SCC for heavy-duty diesel vehicles	Description
202	302			Catalyst Start Exhaust
203	303			Catalyst Running Exhaust
204	304			Non-catalyst Start
205	305			Non-catalyst Running
206	306			Hot Soak
207	307			Diurnal Evaporatives
		808	408, 508	Diesel Exhaust
209	309			Running Evaporatives
210	310			Resting Evaporatives
211	311			Multi-Day Resting
212	312			Multi-Day Diurnal
213	313	813	413, 513, 613,	PM Tire Wear
214	314	814	414, 514, 614,	PM Brake Wear
215	315			Catalyst Buses
216	316			Non-catalyst Buses
		817	617, 717	Diesel Bus
218	318			Catalyst Idle
219	319			Non-catalyst Idle
		820	420, 520, 620,	Diesel Idle
221	321			PM Road Dust

DTIM and EMFAC2014 are both run using the 13 vehicle types shown in Table 8. In order to obtain better resolved spatiotemporal surrogates, the DTIM runs are split by light-duty (LDA, LDT1, LDT2, MDV, LHDT1, LHDT2, Urban Bus, MH, MCY) and heavy-duty (T6/T7 HHDT, SBUS, Other BUS) vehicle classes, and also by fuel type (gas, diesel). Each DTIM run outputs emissions for categories from 1-13; therefore, the mapping from Table 8 is used to preserve the spatial surrogates for each of the four DTIM runs. These codes depend on vehicle type, technology, and whether the vehicle is catalyst, non-catalyst, or diesel.

Table 8 Vehicle classification and type of adjustment

DTIM Category	Vehicle type	Type of adjustment
1	LDA	LD
2	LDT1	LD
3	LDT2	LD
4	MDV	LD
5	LHDT1	LM
6	LHDT2	LM
7	T6	LM
8	T7 HHDT	HHDT
9	Other Bus	LM
10	School Bus	Unadjusted on weekdays, zeroed on weekends
11	Urban Bus	LD
12	Motorhomes	LD
13	Motorcycles	LD

**3.4.4. Temporal Adjustment (Day-of-Week adjustments to EMFAC daily totals):** EMFAC2014 produces average day-of-week (DOW) estimates that represent Tuesday, Wednesday, and Thursday. In order to more accurately represent daily emissions, DOW adjustments are made to all emissions estimated on a Friday, Saturday, Sunday or Monday. The DOW adjustment factors were developed using CalVAD data. The California Vehicle Activity Database (CalVAD) developed by UC Irvine for ARB, is a system that fuses available data sources to produce a “best estimate” of vehicle activity by class. The CalVAD data set includes actual daily measurements of VMT on

the road network for 43 of the 58 counties in California. However, there are seven counties that can't be used because the total vehicle miles traveled are less than the sum of the heavy heavy-duty truck vehicle miles traveled and trucks excluding heavy heavy-duty vehicle miles traveled. Furthermore, two more counties that have high vehicle miles traveled on Sunday are also excluded. Therefore, only 34 of these counties had useful data. In order to fill the missing 24 counties' data to cover all of California, a county which is nearby and similar in geography is selected for each of the missing counties. The CalVAD fractions were developed for three categories of vehicles: passenger cars (LD), light- and medium-duty trucks (LM), and heavy-heavy duty trucks (HHDT). Table 8 also shows the corresponding assignment to each vehicle type. Furthermore, the CalVAD fractions are scaled so that a typical workday (Tuesday, Wednesday, or Thursday) gets a scaling factor of 1.0. All other days of the week receive a scaling factor where their VMT is related back to the typical work day. This means there are a total of five weekday scaling factors. Lastly, the CalVAD data were used to create a typical holiday, because the traffic patterns for holidays are quite different than a typical week day. Thus, in the end, there are six daily fractions for each of the three vehicle classes, for all 58 counties. The DOW factors and vehicle type can be found in Appendix A: Day of week redistribution factors by vehicle type and county.

**3.4.5. Temporal Adjustment (Hour-of-Day re-distribution of hourly travel network volumes):** The travel networks provided by local transportation agencies and used with DTIM represent an hourly distribution for an average day. As for EMFAC, it is assumed that these average day-of-week hourly distributions represent hourly mid-week activities (i.e. for Tuesday, Wednesday, and Thursday). As such, they lack the temporal variations that are known to occur on other days of the week. To rectify this, the CalVAD data were used to develop hour-of-day profiles for Friday through Monday and a typical holiday. In a similar manner as the DOW factors, these hour-of-

day profiles are used to re-allocate the hourly travel network distributions used in DTIM to Friday through Monday and a typical holiday. The hour-of-day profiles can be found in Appendix B: Hour of Day Profiles by vehicle type and county.

**3.4.6. Summary of On-road Emissions Processing Steps:** Eight general steps are used to spatially and temporally allocate EMFAC emissions by hour and grid cell:

1. Activity Data
  - a. EMFAC is run in default mode for a single day to generate hourly activity data for each vehicle type and county: VMT, vehicle population, and number of vehicle trips. This is a single day's run, as EMFAC2014 yields the same hourly activity data for every day of the year.
  - b. The activity data are used to generate various input files for ITN and DTIM, the general goal being to determine how much each activity belongs to each vehicle type through the day.
  
2. Road Network
  - a. Pull a full copy of the California road network from the ITN database, using MPO inputs.
  - b. Convert the ITN results to a form readable by DTIM.
  - c. Apply travel network volumes by county hourly DOW fractions.
  
3. Meteorological Input Data
  - a. Gridded, hourly temperature (T) and relative humidity (RH) are modeled using CALMET. Section 3.1 describes the development of these meteorological (met) data in more detail.
  - b. Daily met files are prepared in formats readable by both EMFAC2014 and DTIM4.

4. EMFAC Emission Rates
  - a. EMFAC is run in emissions rates mode (using monthly-average T and RH) to generate a look-up table of on-road mobile source emission rates by speed, temperature, and relative humidity for each county. These results are created on a monthly-average basis to save processing time.
  - b. The emissions rates are pulled from the EMFAC database and reformatted in the DTIM-ready IRS file format.
  
5. EMFAC Emissions
  - a. EMFAC is run in emissions mode (using day-specific T and RH) to provide county-wide on-road mobile source emission estimates by day and hour for EMFAC categories.
  - b. These results are saved for later use.
  
6. DTIM
  - a. DTIM is run for one week (five representative days since Tuesday, Wednesday and Thursday are treated as a single day) and one holiday in the summer and in the winter.
  - b. Convert the DTIM output results into MEDS format for further processing.

More details on the DTIM and scaling processing can be found in the Appendix C.

7. Scale EMFAC Emissions Using DTIM
  - a. For each day of EMFAC emissions, the closest day-of-week matching DTIM file is chosen for scaling.
  - b. The daily, county-wide EMFAC emissions are distributed spatially and temporally using the DTIM MEDS files as surrogates, as shown by the equation:

$$E_{P,ij,hr,cat} = \frac{EF_{P,cat} \times DTIM_{P,ij,hr,cat}}{DTIM_{P,daily,cat,cnty}}$$

where:

E = grid cell emissions  
EF = EMFAC emissions  
DTIM = DTIM emissions  
p = pollutant  
i,j = grid cell  
hr = hourly emissions  
cat = emission category  
daily = daily emissions  
cnty = county

- c. Finally, the Caltrans day-of-week factors are applied to the gridded, hourly emissions to better match traffic patterns.

#### 8. Final Formatting

- a. The final step of on-road emissions processing is to convert the gridded, hourly emissions data to a NetCDF file usable by the CMAQ photochemical model.

### 3.5. Estimation of Gridded Biogenic Emissions

Biogenic emissions were estimated using the Model of Emissions of Gases and Aerosols from Nature (MEGAN) version 2.04 (Guenther, et al., 2006). MEGAN estimates biogenic emissions as a function of normalized emission rates (i.e. emission rates at standard conditions), which are adjusted to reflect variations in temperature, light, leaf area index (LAI), and leaf age (estimated from changes in LAI). The default MEGAN input databases for emission factors (EFs), plant functional types (PFTs), and LAI are not used in the application of MEGAN in California. Instead, California-specific emission factor and PFT databases were translated from those used in the Biogenic Emission Inventory GIS (BEIGIS) system (Scott & Benjamin, 2003) to improve emission estimates and to maintain consistency with previous California biogenic emission inventories. LAI data were derived from the MODIS 8-day LAI satellite product. Hourly surface temperatures were from observations gridded with the CALMET meteorological model and insolation data was provided by the WRF meteorological fields, as discussed in Section 3.1. Emissions of isoprene, monoterpenes, and methylbutenol were

estimated from California-specific gridded emission factor data, while emissions of sesquiterpenes, methanol, and other volatile organic compounds were estimated from California-specific PFT data and PFT-derived emission rates.

MEGAN emissions estimates for California were evaluated during the California Airborne BVOC Emission Research in Natural Ecosystems Transects (CABERNET) field campaign in 2011 (Karl, et al., 2013), (Misztal, et al., 2014) and were shown to agree to within +/-20% of the measured fluxes (Misztal, et al., 2015), which is well within the stated model uncertainty of 50%.

### **3.6. Estimation of Other Day-Specific Sources**

Day-specific data were used for preparing base case inventories when data were available. ARB and district staffs were able to gather hourly/daily emission information for 1) wildfires and prescribed burns 2) paved and unpaved road dust 3) agricultural burns in six districts and 4) a refinery fire. Additionally, emissions in future years were removed for facilities that have closed after 2012.

For the reference and future year inventories, which are used to calculate Relative Response Factors (RRFs), day-specific emissions for wildfires, prescribed burns, wildland fires use (WFU) and the Chevron fire are left out of the inventory. All other day-specific data are included in both reference and future year modeling inventories.

**3.6.1. Wildfires and Prescribed Burns:** Day-specific, base case estimates of emissions from wildfires and prescribed fires were developed in a two-part process. The first part consisted of estimating micro-scale, fire-specific emissions (i.e. at the fire polygon scale, which can be at a smaller spatial scale than the grid cells used in air quality modeling). The second part consisted of several steps of post-processing fire polygon emission estimates into gridded, hourly emission estimates that were formatted for use in air quality modeling.



Fire event-specific emissions were estimated using a combination of geospatial databases and a federal wildland fire emission model, first described in (Clinton, et al., 2006). A series of pre-processing steps were performed using a Geographic Information System (GIS) to develop fuel loading and fuel moisture inputs to the First Order Fire Effects (FOFEM) fire emission model (Lutes, et al., 2012). Polygons from a statewide interagency fire perimeters geodatabase (fire12\_1.gdb, downloaded June 4, 2013) maintained by the Fire and Resource Assessment Program (FRAP) of the California Department of Forestry and Fire Protection (CALFIRE) provided georeferenced information on the location, size (area), spatial shape, and timing of wildfires and prescribed burns. (Under interagency Memorandums of Understanding, federal, state, and local agencies report California wildfire and prescribed burning activity data to FRAP.) Using GIS software, fire polygons were overlaid upon a vegetation fuels raster dataset called the Fuel Characteristic Classification System (FCCS) (Ottmar, et al., 2007). The FCCS maps vegetation fuels at a 30 meter spatial resolution, and is maintained and distributed by LANDFIRE.GOV, a state and federal consortium of wildland fire and natural resource management agencies. With spatial overlay of fire polygons upon the FCCS raster, fuel model codes were retrieved and component areas within each fire footprint tabulated. For each fuel code, loadings (tons/acre) for fuel categories were retrieved from a FOFEM look-up table. Fuel categories included dead woody fuel size classes, overstory live tree crown, understory trees, shrubs, herbaceous vegetation, litter and duff. Fuel moisture values for each fire were estimated by overlaying fire polygons on year- and month-specific 1 km spatial resolution fuel moisture raster files generated from the national Wildland Fire Assessment System (WFAS.net) and retrieving moisture values from fire polygon centroids. Fire event-specific fuel loads and fuel moisture values were compiled and formatted to a batch input file and run through FOFEM.

A series of post-processing steps were performed on the FOFEM batch output to include emission estimates (pounds/acre) for three supplemental

pollutant species ( $\text{NH}_3$ , TNMHC and  $\text{N}_2\text{O}$ ) in addition to the seven species native to FOFEM ( $\text{CO}$ ,  $\text{CO}_2$ ,  $\text{PM}_{2.5}$ ,  $\text{PM}_{10}$ ,  $\text{CH}_4$ ,  $\text{NO}_x$ ,  $\text{SO}_2$ ), and to calculate total emissions (tons) by pollutant species for each fire. Emission estimates for  $\text{NH}_3$ , TNMHC and  $\text{N}_2\text{O}$  were based on mass ratios to emitted  $\text{CO}$  and  $\text{CO}_2$  (Gong, et al., 2003)

Fire polygon emissions were apportioned to CMAQ model grid cells using area fractions, developed using GIS software, by intersecting fire polygons to the grid domain.

Another set of post-processing steps were applied to allocate fire polygon emissions by date and hour of the day. Fire polygon emissions were allocated evenly between fire start and end dates, taken from the fire perimeters geodatabase. Daily emissions were then allocated to hour of day and to the model grid cells and distributed vertically using a method developed by the Western Regional Air Partnership (WRAP), which specifies a pre-defined diurnal temporal profile, plume bottom and plume top for each fire. (WRAP, 2005)

**3.6.2. Paved Road Dust:** Statewide emissions from paved road dust were adjusted for each day of the baseline year. The adjustment reduced emissions by 25% from paved road dust on days when precipitation occurred. Paved road dust emissions are calculated using the AP-42 method described in (U.S. EPA, 2011).

This methodology includes equations that adjust emissions based on average precipitation in a month; these precipitation-adjusted emissions were placed in the CEIDARS and CEPAM databases. Since daily precipitation totals are readily available, ARB and district staff agreed that paved road dust emissions should be estimated for each day rather than by month as described in the AP-42 methodology. The emissions from CEIDARS were

replaced with day-specific data. A description of the steps used to calculate day-specific emissions is as follows:

Daily uncontrolled emissions for each county/air basin are estimated from the AP-42 methodology [Equation (1) on page 13.2.1-4]. No monthly precipitation adjustments are incorporated into the equation to estimate emissions.

To adjust for precipitation, daily precipitation data for 2013 were provided by an in-house database maintained by ARB staff that stores collected meteorology data from outside sources. The specific data sources for these data include: Remote Automated Weather Stations (RAWS), Atmospheric Infrared Sounder (AIRS), California Irrigation Management Information System (CIMIS) networks, SFBMET (a meteorological database maintained by the Bay Area Air Quality Management District), and Federal Aviation Administration (FAA). FAA data provide precipitation data collected from airports in California.

If the precipitation is greater than or equal to 0.01 inches (measured anywhere in a county or county/air basin piece on a particular day), then the uncontrolled emissions are reduced by 25% for that day only. This reduction of emissions follows the recommendation in AP-42 as referenced above.

Replace the annual average emissions with day-specific emissions for every day in the corresponding emission inventory dataset.

**3.6.3. Unpaved Road Dust:** Statewide emissions from unpaved road dust were adjusted for rainfall suppression for each day of the year. The adjustment reduced county-wide emissions by 100% (total suppression) from unpaved road dust on days when precipitation greater than 0.01" occurred in a county/air basin. Dust emissions from unpaved roads were calculated using an emission factor derived from tests conducted by the University of California, Davis, and the Desert Research Institute (DRI). Unpaved road

vehicle miles traveled (VMT) were based on county-specific road mileage estimates.

Emissions were assumed to be suppressed for each day with rainfall of 0.01 inch or greater using equation (2) from the method described in (U.S. EPA, 2011). The equation adjusts emissions based on annual precipitation; these precipitation-adjusted emissions were placed in the CEIDARS database. Similar to paved road dust, ARB and district staff agreed that unpaved road dust emissions should be estimated for each day. The emissions from CEIDARS were replaced with day-specific data for the appropriate years. Following is a description of the steps that were taken to calculate day-specific emissions.

- a) Start with the daily uncontrolled emissions for each county/air basin as estimated from ARB's methodology. In other words, no precipitation adjustments have been incorporated in the emission estimates.
- b) Use the same daily precipitation data as for paved road dust (see above)
- c) If the precipitation is greater than or equal to 0.01 inches measured anywhere in a county or county/air basin portion on a particular day, then the emissions are removed for that day only.
- d) Replace the annual average emissions with day-specific emissions for every day.

**3.6.4. Agricultural Burning:** Agricultural burning day-specific emission estimations were incorporated into the inventory for the following areas:

**San Joaquin Valley**

The San Joaquin Valley Air Pollution Control District estimated emissions for each day of 2012 when agricultural burning occurred. Emissions were estimated for the burning of prunings, field crops, weed abatement and other solid fuels. Information needed to estimate emissions came from the district's

Smoke Management System, which stores information on burn permits issued by the district. In order to obtain a daily burn authorization, the person requesting the burn provides information to the district, including the acres and type of material to be burned, the specific location of the burn and the date of the burn. Acres are converted to tons of fuel burned using a fuel loading factor based on the specific crop to be burned. Emissions are calculated by multiplying the tons of fuel burned by a crop-specific emission factor. More information can be found in (ARB-Miscellaneous Methodologies, 2013).

To determine the location of the burn, district staff created spatial allocation factors for each 4 kilometer grid cell used in modeling. These factors were developed for “burn zones” in the San Joaquin Valley based on the agricultural land coverage. Daily emissions in each “agricultural burn zone” were then distributed across the zone/grid cell combinations using the spatial allocation factors. Emissions were summarized by grid cell and day.

Burning was assumed to occur over three hours from 10:00 a.m. to 1:00 p.m., except for two categories. Orchard removals were assumed to burn over eight hours from 10:00 a.m. to 6:00 p.m. Vineyard removals were assumed to burn over five hours from 10:00 a.m. to 3:00 p.m.

### **Sacramento**

Sacramento Metropolitan Air Quality Management District provided information needed to calculate emissions in Sacramento County from agricultural burning for each day of 2012 when agricultural burning occurred. Using the same methodology as San Joaquin Valley, emissions were estimated for the burning of prunings, field crops, weed abatement and other solid fuels. Information needed to estimate emissions came from burn permits issued by the district. In order to obtain a burn permit, the person requesting the burn provides information to the district, including the acres to be burned,

the specific location of the burn and the date of the burn. Acres are converted to tons of fuel burned using a fuel loading factor based on the specific crop to be burned. Emissions are calculated by multiplying the tons of fuel burned by a crop-specific emission factor. The location of the burn was converted to latitude/longitude based on the address or description of location provided by the burn permit holder, then ultimately to grid cell. Burning was assumed to occur over eight hours from 10:00 a.m. to 6:00 p.m.

### **Yolo-Solano**

Yolo-Solano Air Quality Management District provided information needed to calculate emissions from agricultural burning for each day of 2012 when agricultural burning occurred. Data were provided for their region: all of Yolo County and the Sacramento Valley portion of Solano County. Using the same methodology as San Joaquin Valley, emissions were estimated for the burning of prunings, field crops, weed abatement and range improvement. The location of the burn was converted to latitude/longitude based on the address or description of location provided by the burn permit holder, then ultimately to grid cell. Burning was assumed to occur over five hours from 11:00 a.m. to 4:00 p.m.

### **Feather River**

Feather River Air Quality Management District provided information needed to calculate emissions from agricultural and prescribed burning for each day of 2012 when agricultural burning occurred. Data were provided for Sutter and Yuba Counties. Using the same methodology as San Joaquin Valley, emissions were estimated for the burning of prunings, field crops, weed abatement and other solid waste. The location of each burn was converted to latitude/longitude based on the address or description of location provided by the burn permit holder, then ultimately to grid cell. Orchard prunings were

assumed to occur from 9:00 a.m. to 4:00 p.m. The burning of field crops, rice, weeds and ditch banks were assumed to occur from 10:00 a.m. to 5:00 p.m. from March 1 through August 31 and from 10:00 a.m. to 4:00 p.m. from September 1 through February 29. Prescribed burns over 10 acres were assumed to occur from 9:00 a.m. to 12:00 a.m. while prescribed burns less than 10 acres were assumed to occur from 9:00 a.m. to 6:00 p.m.

### **Ventura**

Ventura County Air Pollution Control District provided emissions in Ventura County from agricultural burning for each day of 2012 when agricultural burning occurred. Using the same methodology as San Joaquin Valley, emissions were estimated for the burning of prunings, field crops, weed abatement, range improvement and prescribed burns not included in the wildfires / prescribed burns discussed in the San Joaquin Valley portion of Section 3.6.4. Information needed to estimate emissions came from burn permits issued by the district. In order to obtain a burn permit, the person requesting the burn provides information to the district, including the acres to be burned, the specific location of the burn and the date of the burn. Acres are converted to tons of fuel burned using a fuel loading factor based on the specific crop to be burned. Emissions are calculated by multiplying the tons of fuel burned by a crop-specific emission factor. The location of the burn was converted to latitude/longitude based on the address or description of location provided by the burn permit holder, then ultimately to grid cell. Burning was assumed to occur over three hours from 9:00 a.m. to 12:00 p.m.

### **Imperial**

Imperial County Air Pollution Control District provided information needed to calculate emissions from agricultural and prescribed burning for each day of 2012 when agricultural burning occurred. Using the same methodology as

San Joaquin Valley, emissions were estimated for the burning of field crops and weed abatement. The location of each burn was converted to latitude/longitude based on the nearest crossroads provided by the burn permit holder, then ultimately to grid cell. Burning was assumed to occur over four hours from 11:00 a.m. to 3:00 p.m.

**3.6.5. Refinery Fire:** On August 6, 2012, the Chevron U.S.A Inc. refinery in Richmond experienced a catastrophic pipe rupture. The flammable, high temperature gas oil flowing through the pipe ignited shortly after the release and burned for approximately 5 hours. Flaring also occurred for four days from August 6 through August 10. Bay Area Air Quality Management District (BAAQMD) staff estimated NO<sub>x</sub> and SO<sub>x</sub> emissions from both the fire and flaring; TOG emissions from flaring were also estimated. The emissions were spread evenly across the hours they occurred.

Additionally, stack data were estimated by the BAAQMD. Based on physical observation of the plume height, the first two hours of the fire were estimated to have the highest gas flow rate used in the calculation of plume rise. The gas flow rate was reduced for the latter three hours of the fire.

**3.6.6. Closed Facilities:** Emissions in future years were removed for facilities that have closed beyond the baseline year. In other words, the emissions were removed from future year inventories for a facility that was included in the 2012 inventory but stopped operating after 2012. Local air district staffs provided the lists of facilities.



## **4. Quality Assurance of Modeling Inventories**

As mentioned in Section 1.3, base case modeling is intended to demonstrate confidence in the modeling system. Quality assurance of the data is fundamental in order to detect any possible outliers and potential problems with emission estimates. The most important quality assurance checks of the modeling emissions inventory are summarized in the following sections.

### **4.1. Area and Point Sources**

Before utilizing SMOKE to process the annual emissions totals into temporally, chemically, and spatially-resolved emissions inventories for photochemical modeling, all SMOKE inputs are subject to extensive quality assurance procedures performed by ARB staff. Annual and forecasted emissions are carefully reviewed before input into SMOKE. ARB and district staff review data used to calculate emissions along with other associated data, such as the location of facilities and assignment of SCC to each process. Growth and control information are reviewed and updated as needed.

The next check is to compare annual average emissions from CEPAM with planning inventory totals to ensure data integrity. The planning and modeling inventories start with the same annual average emissions. The planning inventory is developed for an average summer day and an average winter day, whereas the modeling inventory is developed by month. Both inventory types use the same temporal data described in Section 2.2. The summer planning inventory uses the monthly throughputs from May through October. Similarly, the winter planning inventory uses the monthly throughputs from November through April. The modeling inventory produces emissions for a weekday, Saturday and Sunday for each month.

Annual emissions totals are plotted using the same gridding inputs as used in SMOKE in order to visually inspect and analyze the spatial allocation of emissions independent of temporal allocation and chemical speciation. Spatial plots by source category like the one shown in Figure 3 are carefully screened for proper spatial distribution of emissions.




Figure 3 Example of a spatial plot by source category

Before air quality model-ready emissions files are generated by SMOKE, the run configurations and parameters set within the SMOKE environment are checked for consistency for both the reference and future years.

To aid in the quality assurance process, SMOKE is configured to generate inventory reports of temporally, chemically, and spatially-resolved emissions inventories. ARB staff utilize the SMOKE reports by checking emissions totals by source category and

region, creating and analyzing time series plots, and comparing aggregate emissions totals with the pre-SMOKE emissions totals obtained from CEPAM. A screenshot capture of a portion of such report can be seen in Figure 4.

```
# Processed as Area sources
# Base inventory year      2012
# No gridding matrix applied
# No speciation matrix applied
# Temporal factors applied for episode from
#   Wednesday Aug. 8, 2012 at 030000 to
#   Thursday Aug. 9, 2012 at 060000
# Annual total data basis in report
#
#Date      , Region      , SCC      , [tons/day] , [tons/day] , [tons/day] , [tons/day] , [tons/day] , [tons/day]
#CO      , NOX      , TOG      , NH3      , SOX      , PM
08/09/2012, 0LC006017LAK, 00600005294212000010, 0.19098E-01, 0.46288E-01, 0.44956E-02, 0.00000E+00, 0.16055E-03, 0.16051E-02
08/09/2012, 0LC006017LAK, 00600005294212000011, 0.94998E-02, 0.21652E-01, 0.30532E-02, 0.00000E+00, 0.00000E+00, 0.11252E-02
08/09/2012, 0LC006017LAK, 00600011011003000000, 0.00000E+00, 0.00000E+00, 0.00000E+00, 0.63987E-03, 0.00000E+00, 0.60000E+00
08/09/2012, 0LC006017LAK, 00600012012202420000, 0.00000E+00, 0.00000E+00, 0.00000E+00, 0.29915E-01, 0.00000E+00, 0.60000E+00
08/09/2012, 0LC006017LAK, 00600019917002400000, 0.00000E+00, 0.00000E+00, 0.00000E+00, 0.13904E-01, 0.00000E+00, 0.60000E+00
08/09/2012, 0LC006017LAK, 00600021020033000000, 0.00000E+00, 0.00000E+00, 0.13736E-01, 0.00000E+00, 0.00000E+00, 0.60000E+00
08/09/2012, 0LC006017LAK, 00600021020081500000, 0.00000E+00, 0.00000E+00, 0.31439E-02, 0.00000E+00, 0.00000E+00, 0.60000E+00
08/09/2012, 0LC006017LAK, 00600022020405000000, 0.00000E+00, 0.00000E+00, 0.31245E-01, 0.00000E+00, 0.00000E+00, 0.60000E+00
08/09/2012, 0LC006017LAK, 00600022020430220000, 0.00000E+00, 0.00000E+00, 0.72951E-03, 0.00000E+00, 0.00000E+00, 0.60000E+00
08/09/2012, 0LC006017LAK, 00600022020430830000, 0.00000E+00, 0.00000E+00, 0.36475E-03, 0.00000E+00, 0.00000E+00, 0.60000E+00
08/09/2012, 0LC006017LAK, 00600022020432040000, 0.00000E+00, 0.00000E+00, 0.36475E-03, 0.00000E+00, 0.00000E+00, 0.60000E+00
```

Figure 4 Screen capture of a SMOKE-generated QA report

**4.1.1. Area and Point Sources Temporal Profiles:** Checks for missing or invalid temporal assignments are conducted to ensure accurate temporal allocation of emissions. Special attention is paid to checking monthly throughputs and appropriate monthly temporal distribution of emissions for each source category. In addition, checks for time-invariant temporal assignments are done for certain source categories and suitable alternate temporal assignments are determined and applied. For the agricultural source sector (e.g. agricultural pesticides/fertilizers, farming operations, fugitive windblown dust, managed burning and disposal, and farm equipment), replacement temporal assignments are extracted from the Agricultural Emissions Temporal and Spatial Allocation Tool (AgTool). (Anderson, et al., 2012). The AgTool is a database management system capable of temporally and spatially allocating emissions from the agricultural source sector. It was developed by Sierra Research, Inc. and its subcontractor Alpine Geophysics, LLC along with collaboration from ARB and the San Joaquin Valley Air Pollution Control District (SJVAPCD). Temporal allocation data outputs from the AgTool, were compiled using input data

provided by the UC Cooperative Extension, U.S. Department of Agriculture (USDA), and the CA Department of Pesticide Regulation (DPR).

Further improvements to temporal profiles used in the allocation of area source emissions are performed using suitable alternate temporal assignments determined by ARB staff. Select sources from manufacturing and industrial, degreasing, petroleum marketing, mineral processes, consumer products, residential fuel combustion, farming operations, aircraft, and commercial harbor craft sectors are among the source categories included in the application of adjustments to temporal allocation.

#### **4.2. On-road Emissions**

There are several processes to conduct quality assurance of the on-road mobile source modeling inventory at various stages of the inventory processing. The specific steps taken are described below:

1. Generate an ITN spatial plot to check if there were any missing network activities.
2. Generate a time series plot for each county to check the diurnal pattern of network activities.
3. Generate time series plots for the DTIM output files by county and by SCC to check the diurnal pattern.
4. Generate time series plots for the on-road mobile source files after scaling to EMFAC 2014 emissions (MEDS files) by county and SCC to check the diurnal pattern.
5. Compare the statewide daily total emissions for the MEDS files and the EMFAC 2014 emissions files to ensure that the emissions are the same.
6. Generate the spatial plot for the MEDS file to check if there were any missing emissions.
7. Generate time series and spatial plots again to check the final MEDS files.

### 4.3. Day-specific Sources

**4.3.1. Wildfires and Prescribed Burns:** To check for potential wildfire activity data gaps in the CALFIRE interagency fire perimeters geodatabase, staff examined geospatial fire activity data reported in the national Geospatial Multi-Agency Coordination ([www.geomac.gov](http://www.geomac.gov)) wildland fire geodatabase. California wildfires reported to GeoMAC were accounted for in the CALFIRE geodatabase.

Prescribed burns are performed by land and fire management agencies primarily to reduce wildfire risk to local communities associated with high loads of vegetation fuels in adjacent wildlands. Vegetation is burned during winter, in-situ or in piles following mechanical treatment. Public land management agencies also perform prescribed burning to restore the natural role of fire in selected ecosystems. To check for potential prescribed burn activity data gaps in the CALFIRE interagency fire perimeters geodatabase, staff queried data for calendar year 2012 reported to ARB's Prescribed Fire Information Reporting System (PFIRS) (<https://ssl.arb.ca.gov/pfirs/index.php>). Staff discovered that CALFIRE data accounted for 38 prescribed burn projects, while PFIRS reported 453 projects. Only one burn project was accounted for in both datasets. Burn project area for CALFIRE data totaled approximately 3,780 acres, while burned acres reported to PFIRS totaled 9,097 acres. Burn projects reported to PFIRS were located in the Sierra Nevada Mountains and northern Coast Range.

Records for 651 prescribed wildland burn events reported for 2012 were downloaded from PFIRS and imported to a geodatabase. Data fields included event ("Unit") name, burned area, latitude/longitude, start and end dates. A series of geoprocessing steps were used to map and overlay prescribed burns as points on the statewide vegetation fuels (FCCS) and moisture raster datasets, to retrieve associated fuel loadings and moisture values for use as input to FOFEM. Prescribed burn points were also overlaid on the statewide 4-km modeling grid to assign grid cell IDs to each

burn. Emission estimates for each prescribed burn event were generated by FOFEM and summarized in an Access database.

**4.3.2. Paved Road Dust:** The average daily emissions inventory was adjusted with day-specific precipitation data to produce a day-specific emissions inventory. Total emissions by county before the adjustment were compared to CEPAM for a reasonable match. After the adjustment, the day-specific total emissions by county were compared to CEPAM using time series plots. These plots were verified to confirm that there were only two values for every county/air basin/district: high values and low values. The high values are emissions that were not affected by rain adjustment, while the low values are emissions that were affected by the 25% rain adjustment reduction. Additionally the day-specific total was also compared to other inventory years to verify the expected growth trend.

**4.3.3. Unpaved Road Dust:** Unpaved road dust followed the same quality assurance process as paved road dust, except that total removal rather than 25% reduction is applied whenever precipitation is greater than 0.01”.

**4.3.4. Agricultural Burning:** Checks were done to verify the quality of the agricultural burn data. The day-specific emissions from agricultural burning were compared to the emissions from CEPAM for each county to check for reasonableness. Time series plots were reviewed for each county to see that days when burning occurred matched the days provided by the local air district. For each county, a few individual fires were calculated by hand starting from the raw data through all the steps to the final MEDS files to make sure the calculations were done correctly. Spatial plots were made to double check the locations of each burn.

**4.3.5. Refinery Fire:** The calculations in the MEDS files were verified by hand to make sure the emissions and stack data matched what was provided by the BAAQMD.

#### **4.4. Additional QA**

In addition to the QA described above, comparisons are made between annual average inventories from CEPAM and modeling inventories. The modeling inventory shows emissions by month and subsequently calculates the annual average for comparison with CEPAM emissions. Annual average inventories and modeling inventories can be different, but differences should be well understood. For example, modeling inventories are adjusted to reflect different days of the week for on-road motor vehicles as detailed in Section 3.4; since weekend travel is generally less than weekday travel, modeling inventory emissions are usually lower when compared to annual average inventories from CEPAM. Figure 5 provides a screen capture of a report that summarizes different emission categories for San Luis Obispo County. Please note that this table is only an example since emissions have been updated from what is displayed here.

County:40 Spec:NOx

EIC	Description	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual	CEPAM	Difference
10	electric utilities	0.12	0.11	0.1	0.06	0.09	0.13	0.13	0.16	0.14	0.16	0.14	0.13	0.12	0.12	0.00
20	cogeneration	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.00
30	oil and gas production (combustion)	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.00
40	petroleum refining (combustion)	0.3	0.3	0.26	0.3	0.33	0.33	0.33	0.33	0.33	0.33	0.33	0.26	0.31	0.31	0.00
50	manufacturing and industrial	0.06	0.06	0.06	0.06	0.07	0.06	0.07	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.00
52	food and agricultural processing	0.19	0.19	0.19	0.34	0.34	0.34	0.38	0.38	0.38	0.18	0.18	0.18	0.27	0.27	0.00
60	service and commercial	0.91	0.92	0.92	0.92	0.92	0.9	0.9	0.91	0.91	0.91	0.92	0.91	0.91	0.91	0.00
99	other (fuel combustion)	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.00
110	sewage treatment	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
120	landfills	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
130	incinerators	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
140	soil remediation	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
199	other (waste disposal)	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
210	laundering	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
220	degreasing	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
230	coatings and related process solvents	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
240	printing	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
250	adhesives and sealants	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
299	other (cleaning and surface coatings)	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
310	oil and gas production	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
320	petroleum refining	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00
330	petroleum marketing	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
399	other (petroleum production and marketing)	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
410	chemical	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
420	food and agriculture	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
430	mineral processes	0.03	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.04	0.04	0.00
440	metal processes	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
450	wood and paper	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
460	glass and related products	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
470	electronics	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
499	other (industrial processes)	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
510	consumer products	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
520	architectural coatings and related process sol	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
530	pesticides/fertilizers	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
540	asphalt paving / roofing	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
610	residential fuel combustion	0.73	0.73	0.68	0.65	0.57	0.57	0.57	0.57	0.57	0.65	0.7	0.73	0.64	0.64	0.00
620	farming operations	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
630	construction and demolition	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
640	paved road dust	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
645	unpaved road dust	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
650	fugitive windblown dust	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
660	fires	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
670	managed burning and disposal	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.00
690	cooking	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
699	other (miscellaneous processes)	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
700	on-road vehicles	9.34	9.32	9.36	9.17	9.06	8.81	8.69	8.77	8.63	8.79	9.3	9.23	9.04	9.60	0.56
810	aircraft	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.00
820	trains	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.93	0.74
830	ships and commercial boats	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
833	ocean going vessels	11.23	11.23	11.23	11.23	11.23	11.23	11.23	11.23	11.23	11.23	11.23	11.23	11.23	11.52	0.29
835	commercial harbor craft	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	0.83	-0.29
840	recreational boats	0.05	0.05	0.17	0.18	0.16	0.47	0.46	0.43	0.12	0.11	0.11	0.06	0.2	0.20	0.00
850	off-road recreational vehicles	0.03	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.04	0.04	0.00
860	off-road equipment	1.08	1.24	1.21	1.24	1.25	1.28	1.25	1.25	1.28	1.21	1.19	1.12	1.21	1.21	0.00
870	farm equipment	1.08	1.22	1.72	1.77	2.21	2.21	2.16	2.21	2.17	1.52	1.14	1.06	1.71	1.71	0.00
890	fuel storage and handling	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
920	geogenic sources	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0.00
***	Total	26.78	27.05	27.59	27.61	27.93	28.05	27.88	28.01	27.55	26.67	27.01	26.67	27.42	28.73	1.31

Notes:

CEPAM refers to annual average emissions from 2016 SIP Baseline Emission Inventory Tool with external adjustments: <http://outapp.arb.ca.gov/cefs/2016oz>  
 Monthly gridded emissions comes from GeoVAST mo-yr/avg tabular summary - gid 319

**On-road vehicles:** The modeling inventory adjusts on-road by day of week as well as day-specific temperatures and relative humidity - Fridays are higher wit time series plots shows weekdays are ~9-10 tpd

**Trains:** The modeling inventory reflects the revised locomotive emissions; the planning inventory reflects the previous emission estimates

**OGV model produces gridded OGV emissions,** which can vary from planning inventory (these emissions include OC1 and OC2 offshore air basins)

**CHC** The modeling inventory reflects the revised commercial harbor craft emissions; the planning inventory reflects the previous emission estimates

Figure 5 Screenshot of comparison of inventories report



Staff also review how modeling emissions vary over a year. Figure 6 provides an example of a modeling inventory time series plot for San Luis Obispo County for area-wide sources, on-road sources and off-road sources. Again, this figure is only an example.




Figure 6 Daily variation of NOx emissions for mobile sources for San Luis Obispo

#### **4.5. Model ready files QA**

Prior to developing the modeling inventory emissions files used in the photochemical models, the same model-ready emissions files developed for the individual source categories (e.g. on-road, area, point, day-specific sources) are checked for quality assurance. Extensive quality assurance procedures are already performed by ARB staff on the intermediate emissions files (e.g. MEDS, SMOKE-generated reports), however, further checks are needed to ensure data integrity is preserved when the model-ready emissions files are generated from those intermediate emissions files.

Comparisons of the totals for both the intermediate and model-ready emissions files are made. Emissions totals are aggregated spatially, temporally, and chemically to single-layer, statewide, daily values by inventory pollutant. Spatial plots are also generated for both the intermediate and model-ready emissions files using the same graphical utilities and aggregated to the same spatial, temporal, and chemical resolution to allow equal comparison of emissions. Any discrepancies in the emissions totals are reconciled before proceeding with the development of the model-ready inventory emissions files.

Before combining the model-ready emissions files of the individual source category inventories into a single model-ready inventory, they are checked for completeness. Day-specific source inventories (when necessary) should have emissions for every day in the modeling period. Likewise, source inventories with emissions files that use averaged temporal allocation (e.g. day-of-week, weekday/weekend, monthly) should have model-ready emissions files to represent every day in the modeling period. In particular, it is important that during these checks source inventories with missing files are identified and resolved. Once all constituent source inventories are complete, they are used to develop the model-ready inventory used in photochemical modeling. When the modeling inventory files are generated, log files are also generated documenting what each daily model-ready emissions file is comprised of as an additional means of verifying that each daily model-ready inventory is complete.

## Bibliography

Anderson, C., Carlson, T., Dulla, R. & Wilkinson, J., 2012. *AGTOOL Agricultural sector emissions allocation model – Data development*. CCOS\_11-2, Sacramento, CA: Sierra Research Inc.

ARB, 2011. *Heavy Duty Diesel Vehicle Exhaust PM Speciation Profile*. [Online]

Available at:

<http://www.arb.ca.gov/ei/speciate/profilereference/HDDV%20PM%20Profiles%20Final.pdf>

[Accessed 2016 17 August].

ARB-Miscellaneous Methodologies, 2013. *District Miscellaneous Processes Methodologies - Managed Burning and Disposal*. [Online]

Available at: <http://www.arb.ca.gov/ei/areasrc/distmiscprocwstburndis.htm>

[Accessed 10 7 2015].

ARB-MSEI, 2015. *Mobile Source Emissions Inventory -- Categories*. [Online]

Available at: <http://www.arb.ca.gov/msei/categories.htm#emfac2014>

[Accessed 10 07 2015].

ARB-PTSD, 2011. *Appendix D: Emissions estimation methodology for Ocean-going vessels*. [Online]

Available at: <http://www.arb.ca.gov/regact/2011/ogv11/ogv11appd.pdf>

[Accessed 10 7 2015].

Baek, B. a. S. C., 2015. *Final Summary Report: Development of SMOKE version 4.0*. Under contract ITS:12-764, Sacramento, CA: s.n.

Caldwell, P. C. H. -N. B. D. a. B. G., 2009. Evaluation of a WRF dynamical downscaling simulation over California. *Climatic change*, 95(3), pp. 499-521.

Clinton, N., Gong, P. & Scott, K. 2., 2006. Quantification of pollutants emitted from very large wildland fires in Southern California. *Atmospheric Environment*, Volume 40, pp. 3686-3695.

Dun and Bradstreet, 2015. *Market insight database*. [Online]

Available at: <http://www.dnb.com>

[Accessed 15 June 2015].

FactFinder, U. C. B. / A., 2011. "B25040: House heating fuel. 2013 ACS 5-year estimates". [Online]

Available at: <http://factfinder2.census.gov>

[Accessed March 2015].

Fovell, R., 2008. *Validation of WRF Surface meteorology Simulations, presentation November 4th 2008*. Sacramento: CCOS Technical Committee meeting, California Air Resources Board.

Funk, t., Stiefer, P. & Chinkin, L., 2001. *Development of gridded spatial allocation factors for the state of California*. Sacramento: STI.

Gong, P., Clinton, N. & Pu, R. T. Y. a. S. J., 2003. *Extension and input refinement to the ARB wildland fire emissions estimation model. Final report, contract number 00-729*, Sacramento, CA.: Air Resources Board.

Guenther, A. et al., 2006. Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature. *Chem. Phys.*, 6(11), pp. 3181-3210.

Homer, C. et al., 2015. Completion of the 2011 National Land Cover Database for the conterminous United States-Representing a decade of land cover change information. *Photogrammetric Engineering and Remote Sensing*, 81(5), pp. 345-354.

Jackson, B., 2012. *Preparation of a gridded emissions Inventory for Mexico*. [Online]

Available at: [https://www.arb.ca.gov/eos/public/arb\\_mexico\\_ei.pdf](https://www.arb.ca.gov/eos/public/arb_mexico_ei.pdf)

[Accessed 27 09 2016].

Karl, T. et al., 2013. Airborne flux measurements of BVOCs above Californian oak forests: experimental investigation of surface and entrainment fluxes, OH densities and Dahmkohler numbers. *J. Atmos. Sci.*, 70(10), pp. 3277-3287.

Lutes, D., Keane, R. & Reinhardt, E. 2., 2012. *FOFEM 6 User Guide*. USDA Forest Service, Rocky Mountain Research Station, Missoula Fire Sciences Laboratory, Missoula, MT 59808: s.n.

Mc. Rae, G. J., 1980. A simple procedure for calculating atmospheric water vapor concentration. *J. Air Pollution Control Association*, Volume 30, p. 384.

Misztal, P. et al., 2015. Evaluation of regional isoprene emission estimates in California based on direct airborne flux measurements. *Submitted to Atmos. Chem. Phys.*.

Misztal, P. K. et al., 2014. Airborne flux measurements of biogenic volatile organic compounds over California. *Atmos. Chem. Phys.*, Volume 14, pp. 10631-10647.

Ottmar, R., Sandberg, D. & Riccardi, C. a. P. S., 2007. An overview of the Fuel Characteristic Classification System – Quantifying, classifying, and creating fuelbeds for resource planning.. *Canadian Journal of Forest Research*, 37(12), pp. 2383-2393.

Reid, S., Penfold, B. & Chinkin, L., 2006. *Emission inventory for the Central California Ozone Study (CCOS) review of spatial variations of area, non-road mobile, and point sources of emissions*, Petaluma, CA: STI.

Scott, K. & Benjamin, M., 2003. Development of a biogenic volatile organic compounds emission inventory for the SCOS97-NARSTO domain. *Atmos. Environ.*, Volume 37, Supplement 2, pp. 39-49.

Systems Applications Inc., 2001. *DTIM 4 User's Guide*. Prepared for the California Department of Transportation, Sacramento: Systems Applications, Inc..

U.S. EPA, 2011. AP 42, Compilation of Air Pollutant Emission Factors. In: *Volume 1. Chapter 13: Miscellaneous Sources. Section 13.2.1 (Paved Road)*. Fifth ed. s.l.:U.S. EPA.

U.S. EPA, 2014. *Draft Modeling Guidance for Demonstrating Attainment of Air Quality Goals for Ozone, PM<sub>2.5</sub>, and RegionalHaze*, North Carolina: U.S. EPA.

U.S. EPA, 2016. <https://www3.epa.gov/ttn/emc/cem.html>. [Online]  
Available at: <https://www3.epa.gov/ttn/emc/cem.html>  
[Accessed 16 August 2016].

Valade, A., 2009. *Evaluation of the simulation of the 2006 California heat wave by WRF using satellite*, Davis, California: U.C Davis.

WRAP, 2005. *2002 Fire Emission Inventory for the WRAP Region Phase I – Essential Documentation*. [Online]  
Available at:  
[http://www.wrapair.org/forums/fejf/documents/emissions/WRAP\\_2002%20EI%20Report\\_20050107.pdf](http://www.wrapair.org/forums/fejf/documents/emissions/WRAP_2002%20EI%20Report_20050107.pdf)  
[Accessed 28 4 2016].

**Appendix A: Day of week redistribution factors by vehicle type and county**

The factors shown in Table 9 represent the “day of week” factors for each county for a broad vehicle class: LD is Light Duty, LM is Light and Medium Duty Trucks, and HH is Heavy- Heavy Duty Trucks.

Table 9 Day of week adjustment by vehicle class and county

County	Day of Week	LD	LM	HH
Alameda	Sunday	0.797	0.496	0.324
Alameda	Monday	0.948	0.919	0.893
Alameda	Tues/Wed/Thurs	1	1	1
Alameda	Friday	1.051	1.014	0.959
Alameda	Saturday	0.929	0.618	0.369
Alameda	Holiday	0.797	0.866	0.829
Alpine	Sunday	1.201	0.821	0.415
Alpine	Monday	1.007	0.945	0.908
Alpine	Tues/Wed/Thurs	1	1	1
Alpine	Friday	1.247	1.082	1.007
Alpine	Saturday	1.219	0.803	0.442
Alpine	Holiday	1.118	0.935	0.832
Amador	Sunday	1.201	0.821	0.415
Amador	Monday	1.007	0.945	0.908
Amador	Tues/Wed/Thurs	1	1	1
Amador	Friday	1.247	1.082	1.007
Amador	Saturday	1.219	0.803	0.442
Amador	Holiday	1.118	0.935	0.832
Butte	Sunday	0.651	0.442	0.41
Butte	Monday	0.964	0.96	0.871
Butte	Tues/Wed/Thurs	1	1	1
Butte	Friday	1.008	1.015	0.962
Butte	Saturday	0.771	0.604	0.503
Butte	Holiday	0.73	0.657	0.606
Calaveras	Sunday	1.201	0.821	0.415
Calaveras	Monday	1.007	0.945	0.908
Calaveras	Tues/Wed/Thurs	1	1	1
Calaveras	Friday	1.247	1.082	1.007
Calaveras	Saturday	1.219	0.803	0.442
Calaveras	Holiday	1.118	0.935	0.832
Colusa	Sunday	0.651	0.442	0.41
Colusa	Monday	0.964	0.96	0.871
Colusa	Tues/Wed/Thurs	1	1	1
Colusa	Friday	1.008	1.015	0.962
Colusa	Saturday	0.771	0.604	0.503
Colusa	Holiday	0.73	0.657	0.606
Contra Costa	Sunday	0.779	0.519	0.376
Contra Costa	Monday	0.943	0.927	0.873
Contra Costa	Tues/Wed/Thurs	1	1	1
Contra Costa	Friday	1.048	1.023	0.982
Contra Costa	Saturday	0.924	0.665	0.471
Contra Costa	Holiday	0.788	0.827	0.799
Del Norte	Sunday	0.85	0.493	0.326
Del Norte	Monday	0.961	0.95	0.915
Del Norte	Tues/Wed/Thurs	1	1	1
Del Norte	Friday	1.031	1.004	0.932
Del Norte	Saturday	0.924	0.619	0.376
Del Norte	Holiday	0.77	0.619	0.527
El Dorado	Sunday	0.972	0.668	0.602
El Dorado	Monday	0.988	0.977	0.943
El Dorado	Tues/Wed/Thurs	1	1	1
El Dorado	Friday	1.178	1.101	0.963
El Dorado	Saturday	1.037	0.786	0.575
El Dorado	Holiday	0.971	0.933	0.921
Fresno	Sunday	0.851	0.443	0.396
Fresno	Monday	1.016	0.934	0.878
Fresno	Tues/Wed/Thurs	1	1	1
Fresno	Friday	1.155	1.026	0.927
Fresno	Saturday	0.946	0.563	0.478
Fresno	Holiday	0.799	0.774	0.784
Glenn	Sunday	0.651	0.442	0.41

County	Day of Week	LD	LM	HH
Glenn	Monday	0.964	0.96	0.871
Glenn	Tues/Wed/Thurs	1	1	1
Glenn	Friday	1.008	1.015	0.962
Glenn	Saturday	0.771	0.604	0.503
Glenn	Holiday	0.73	0.657	0.606
Humboldt	Sunday	0.85	0.493	0.326
Humboldt	Monday	0.961	0.95	0.915
Humboldt	Tues/Wed/Thurs	1	1	1
Humboldt	Friday	1.031	1.004	0.932
Humboldt	Saturday	0.924	0.619	0.376
Humboldt	Holiday	0.77	0.619	0.527
Imperial	Sunday	1.082	0.608	0.396
Imperial	Monday	1.004	0.931	0.948
Imperial	Tues/Wed/Thurs	1	1	1
Imperial	Friday	1.109	1.161	0.983
Imperial	Saturday	1.065	0.687	0.522
Imperial	Holiday	1.024	0.814	0.673
Inyo	Sunday	1.201	0.821	0.415
Inyo	Monday	1.007	0.945	0.908
Inyo	Tues/Wed/Thurs	1	1	1
Inyo	Friday	1.247	1.082	1.007
Inyo	Saturday	1.219	0.803	0.442
Inyo	Holiday	1.118	0.935	0.832
Kern	Sunday	1.114	0.63	0.416
Kern	Monday	1.061	0.942	0.849
Kern	Tues/Wed/Thurs	1	1	1
Kern	Friday	1.253	1.044	0.9
Kern	Saturday	1.1	0.734	0.535
Kern	Holiday	0.986	0.911	0.837
Kings	Sunday	0.663	0.358	0.355
Kings	Monday	0.961	0.909	0.89
Kings	Tues/Wed/Thurs	1	1	1
Kings	Friday	1.045	0.982	0.947
Kings	Saturday	0.807	0.52	0.454
Kings	Holiday	0.669	0.665	0.758
Lake	Sunday	0.85	0.493	0.326
Lake	Monday	0.961	0.95	0.915
Lake	Tues/Wed/Thurs	1	1	1
Lake	Friday	1.031	1.004	0.932
Lake	Saturday	0.924	0.619	0.376
Lake	Holiday	0.77	0.619	0.527
Lassen	Sunday	0.941	0.703	0.587
Lassen	Monday	0.993	0.942	0.798
Lassen	Tues/Wed/Thurs	1	1	1
Lassen	Friday	1.094	1.07	0.882
Lassen	Saturday	0.962	0.766	0.658
Lassen	Holiday	0.968	0.744	0.608
Los Angeles	Sunday	0.858	0.489	0.398
Los Angeles	Monday	0.973	0.936	0.878
Los Angeles	Tues/Wed/Thurs	1	1	1
Los Angeles	Friday	1.047	1.005	0.918
Los Angeles	Saturday	0.979	0.641	0.509
Los Angeles	Holiday	0.863	0.808	0.801
Madera	Sunday	1.017	0.478	0.4
Madera	Monday	1.024	0.942	0.902
Madera	Tues/Wed/Thurs	1	1	1
Madera	Friday	1.176	1.022	0.96
Madera	Saturday	1.105	0.602	0.476
Madera	Holiday	0.866	0.833	0.832
Marin	Sunday	0.779	0.519	0.376
Marin	Monday	0.943	0.927	0.873
Marin	Tues/Wed/Thurs	1	1	1
Marin	Friday	1.048	1.023	0.982
Marin	Saturday	0.924	0.665	0.471
Marin	Holiday	0.788	0.827	0.799
Mariposa	Sunday	1.201	0.821	0.415
Mariposa	Monday	1.007	0.945	0.908
Mariposa	Tues/Wed/Thurs	1	1	1
Mariposa	Friday	1.247	1.082	1.007
Mariposa	Saturday	1.219	0.803	0.442
Mariposa	Holiday	1.118	0.935	0.832
Mendocino	Sunday	0.85	0.493	0.326
Mendocino	Monday	0.961	0.95	0.915
Mendocino	Tues/Wed/Thurs	1	1	1



County	Day of Week	LD	LM	HH
Mendocino	Friday	1.031	1.004	0.932
Mendocino	Saturday	0.924	0.619	0.376
Mendocino	Holiday	0.77	0.619	0.527
Merced	Sunday	1.002	0.593	0.421
Merced	Monday	1.009	0.958	0.904
Merced	Tues/Wed/Thurs	1	1	1
Merced	Friday	1.185	1.103	0.97
Merced	Saturday	1.055	0.713	0.477
Merced	Holiday	0.977	0.897	0.797
Modoc	Sunday	1.133	0.801	0.638
Modoc	Monday	1.159	0.961	0.634
Modoc	Tues/Wed/Thurs	1	1	1
Modoc	Friday	1.202	1.109	0.767
Modoc	Saturday	1.041	0.819	0.745
Modoc	Holiday	1.087	0.992	0.704
Mono	Sunday	1.201	0.821	0.415
Mono	Monday	1.007	0.945	0.908
Mono	Tues/Wed/Thurs	1	1	1
Mono	Friday	1.247	1.082	1.007
Mono	Saturday	1.219	0.803	0.442
Mono	Holiday	1.118	0.935	0.832
Monterey	Sunday	1.2	0.603	0.342
Monterey	Monday	1.106	0.988	0.876
Monterey	Tues/Wed/Thurs	1	1	1
Monterey	Friday	1.116	1.093	0.995
Monterey	Saturday	1.023	0.724	0.7
Monterey	Holiday	1.083	0.755	0.607
Napa	Sunday	1.028	0.624	0.392
Napa	Monday	0.989	0.95	0.895
Napa	Tues/Wed/Thurs	1	1	1
Napa	Friday	1.126	1.041	0.988
Napa	Saturday	1.118	0.743	0.44
Napa	Holiday	0.952	0.905	0.847
Nevada	Sunday	0.972	0.668	0.602
Nevada	Monday	0.988	0.977	0.943
Nevada	Tues/Wed/Thurs	1	1	1
Nevada	Friday	1.178	1.101	0.963
Nevada	Saturday	1.037	0.786	0.575
Nevada	Holiday	0.971	0.933	0.921
Orange	Sunday	0.808	0.415	0.327
Orange	Monday	0.962	0.92	0.891
Orange	Tues/Wed/Thurs	1	1	1
Orange	Friday	1.038	1.025	0.988
Orange	Saturday	0.94	0.587	0.433
Orange	Holiday	0.831	0.774	0.796
Placer	Sunday	0.972	0.668	0.602
Placer	Monday	0.988	0.977	0.943
Placer	Tues/Wed/Thurs	1	1	1
Placer	Friday	1.178	1.101	0.963
Placer	Saturday	1.037	0.786	0.575
Placer	Holiday	0.971	0.933	0.921
Plumas	Sunday	0.651	0.442	0.41
Plumas	Monday	0.964	0.96	0.871
Plumas	Tues/Wed/Thurs	1	1	1
Plumas	Friday	1.008	1.015	0.962
Plumas	Saturday	0.771	0.604	0.503
Plumas	Holiday	0.73	0.657	0.606
Riverside	Sunday	0.894	0.489	0.383
Riverside	Monday	0.974	0.941	0.887
Riverside	Tues/Wed/Thurs	1	1	1
Riverside	Friday	1.085	1.028	0.977
Riverside	Saturday	1.011	0.629	0.491
Riverside	Holiday	0.933	0.848	0.844
Sacramento	Sunday	0.774	0.49	0.431
Sacramento	Monday	0.963	0.954	0.913
Sacramento	Tues/Wed/Thurs	1	1	1
Sacramento	Friday	1.065	1.039	0.973
Sacramento	Saturday	0.884	0.622	0.502
Sacramento	Holiday	0.809	0.832	0.852
San Benito	Sunday	1.2	0.603	0.342
San Benito	Monday	1.106	0.988	0.876
San Benito	Tues/Wed/Thurs	1	1	1
San Benito	Friday	1.116	1.093	0.995
San Benito	Saturday	1.023	0.724	0.7

County	Day of Week	LD	LM	HH
San Benito	Holiday	1.083	0.755	0.607
San Bernardino	Sunday	0.89	0.56	0.532
San Bernardino	Monday	0.988	0.931	0.913
San Bernardino	Tues/Wed/Thurs	1	1	1
San Bernardino	Friday	1.094	1.069	1.012
San Bernardino	Saturday	0.97	0.743	0.634
San Bernardino	Holiday	0.942	0.818	0.831
San Diego	Sunday	0.796	0.532	0.341
San Diego	Monday	0.963	0.928	0.882
San Diego	Tues/Wed/Thurs	1	1	1
San Diego	Friday	1.067	1.022	0.982
San Diego	Saturday	0.928	0.665	0.446
San Diego	Holiday	0.808	0.785	0.785
San Francisco	Sunday	0.852	0.522	0.39
San Francisco	Monday	0.928	0.897	0.888
San Francisco	Tues/Wed/Thurs	1	1	1
San Francisco	Friday	1.05	1.002	0.98
San Francisco	Saturday	0.957	0.639	0.452
San Francisco	Holiday	0.783	0.811	0.84
San Joaquin	Sunday	0.933	0.5	0.393
San Joaquin	Monday	0.984	0.918	0.908
San Joaquin	Tues/Wed/Thurs	1	1	1
San Joaquin	Friday	1.128	1.086	0.976
San Joaquin	Saturday	1.035	0.657	0.466
San Joaquin	Holiday	0.907	0.77	0.757
San Luis Obispo	Sunday	1.038	0.629	0.413
San Luis Obispo	Monday	1.064	0.97	0.935
San Luis Obispo	Tues/Wed/Thurs	1	1	1
San Luis Obispo	Friday	1.113	1.094	1.047
San Luis Obispo	Saturday	0.99	0.725	0.563
San Luis Obispo	Holiday	0.967	0.714	0.669
San Mateo	Sunday	0.714	0.439	0.324
San Mateo	Monday	0.926	0.89	0.887
San Mateo	Tues/Wed/Thurs	1	1	1
San Mateo	Friday	1.02	0.983	0.978
San Mateo	Saturday	0.835	0.55	0.402
San Mateo	Holiday	0.78	0.742	0.767
Santa Barbara	Sunday	0.81	0.388	0.301
Santa Barbara	Monday	1.044	0.952	0.912
Santa Barbara	Tues/Wed/Thurs	1	1	1
Santa Barbara	Friday	1.08	1.011	0.996
Santa Barbara	Saturday	0.829	0.542	0.562
Santa Barbara	Holiday	0.811	0.535	0.545
Santa Clara	Sunday	0.734	0.489	0.343
Santa Clara	Monday	0.954	0.909	0.906
Santa Clara	Tues/Wed/Thurs	1	1	1
Santa Clara	Friday	1.042	1.004	0.953
Santa Clara	Saturday	0.853	0.614	0.4
Santa Clara	Holiday	0.765	0.834	0.807
Santa Cruz	Sunday	0.846	0.526	0.468
Santa Cruz	Monday	0.935	0.923	0.947
Santa Cruz	Tues/Wed/Thurs	1	1	1
Santa Cruz	Friday	1.027	1.012	1.036
Santa Cruz	Saturday	0.935	0.652	0.541
Santa Cruz	Holiday	0.9	0.896	0.875
Shasta	Sunday	1.076	0.823	0.627
Shasta	Monday	0.939	1.007	0.66
Shasta	Tues/Wed/Thurs	1	1	1
Shasta	Friday	1.078	1.156	0.774
Shasta	Saturday	1.117	0.863	0.719
Shasta	Holiday	0.902	0.837	0.602
Sierra	Sunday	0.972	0.668	0.602
Sierra	Monday	0.988	0.977	0.943
Sierra	Tues/Wed/Thurs	1	1	1
Sierra	Friday	1.178	1.101	0.963
Sierra	Saturday	1.037	0.786	0.575
Sierra	Holiday	0.971	0.933	0.921
Siskiyou	Sunday	1.133	0.801	0.638
Siskiyou	Monday	1.159	0.961	0.634
Siskiyou	Tues/Wed/Thurs	1	1	1
Siskiyou	Friday	1.202	1.109	0.767
Siskiyou	Saturday	1.041	0.819	0.745
Siskiyou	Holiday	1.087	0.992	0.704
Solano	Sunday	1.008	0.589	0.36

County	Day of Week	LD	LM	HH
Solano	Monday	0.979	0.948	0.887
Solano	Tues/Wed/Thurs	1	1	1
Solano	Friday	1.13	1.033	0.969
Solano	Saturday	1.091	0.719	0.416
Solano	Holiday	0.909	0.896	0.844
Sonoma	Sunday	0.779	0.519	0.376
Sonoma	Monday	0.943	0.927	0.873
Sonoma	Tues/Wed/Thurs	1	1	1
Sonoma	Friday	1.048	1.023	0.982
Sonoma	Saturday	0.924	0.665	0.471
Sonoma	Holiday	0.788	0.827	0.799
Stanislaus	Sunday	1.002	0.593	0.421
Stanislaus	Monday	1.009	0.958	0.904
Stanislaus	Tues/Wed/Thurs	1	1	1
Stanislaus	Friday	1.185	1.103	0.97
Stanislaus	Saturday	1.055	0.713	0.477
Stanislaus	Holiday	0.977	0.897	0.797
Sutter	Sunday	0.972	0.668	0.602
Sutter	Monday	0.988	0.977	0.943
Sutter	Tues/Wed/Thurs	1	1	1
Sutter	Friday	1.178	1.101	0.963
Sutter	Saturday	1.037	0.786	0.575
Sutter	Holiday	0.971	0.933	0.921
Tehama	Sunday	1.076	0.823	0.627
Tehama	Monday	0.939	1.007	0.66
Tehama	Tues/Wed/Thurs	1	1	1
Tehama	Friday	1.078	1.156	0.774
Tehama	Saturday	1.117	0.863	0.719
Tehama	Holiday	0.902	0.837	0.602
Trinity	Sunday	1.133	0.801	0.638
Trinity	Monday	1.159	0.961	0.634
Trinity	Tues/Wed/Thurs	1	1	1
Trinity	Friday	1.202	1.109	0.767
Trinity	Saturday	1.041	0.819	0.745
Trinity	Holiday	1.087	0.992	0.704
Tulare	Sunday	1.029	0.429	0.185
Tulare	Monday	1.052	0.936	0.912
Tulare	Tues/Wed/Thurs	1	1	1
Tulare	Friday	1.099	1.02	0.97
Tulare	Saturday	0.993	0.67	0.503
Tulare	Holiday	0.942	0.585	0.567
Tuolumne	Sunday	1.201	0.821	0.415
Tuolumne	Monday	1.007	0.945	0.908
Tuolumne	Tues/Wed/Thurs	1	1	1
Tuolumne	Friday	1.247	1.082	1.007
Tuolumne	Saturday	1.219	0.803	0.442
Tuolumne	Holiday	1.118	0.935	0.832
Ventura	Sunday	0.772	0.406	0.491
Ventura	Monday	0.956	0.924	0.932
Ventura	Tues/Wed/Thurs	1	1	1
Ventura	Friday	1.036	0.992	1.004
Ventura	Saturday	0.888	0.554	0.637
Ventura	Holiday	0.817	0.785	0.863
Yolo	Sunday	0.902	0.563	0.357
Yolo	Monday	0.972	0.954	0.932
Yolo	Tues/Wed/Thurs	1	1	1
Yolo	Friday	1.099	1.045	0.973
Yolo	Saturday	0.992	0.669	0.426
Yolo	Holiday	0.895	0.883	0.861
Yuba	Sunday	0.972	0.668	0.602
Yuba	Monday	0.988	0.977	0.943
Yuba	Tues/Wed/Thurs	1	1	1
Yuba	Friday	1.178	1.101	0.963
Yuba	Saturday	1.037	0.786	0.575
Yuba	Holiday	0.971	0.933	0.921

**Appendix B: Hour of Day Profiles by vehicle type and county**

The factors shown in Table 10 represent the “day of week” factors for each county for a broad vehicle class: LD is Light Duty, LM is Light and Medium Duty Trucks, and HH is Heavy- Heavy Duty Trucks.

**Table 10 Hour of Day Profiles by vehicle type and county**

Day of Week	Hour	Alameda			Alpine			Amador			Butte			Calaveras			Colusa			Contra Costa			
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	
Sunday	0	0.020	0.041	0.061	0.010	0.014	0.032	0.010	0.014	0.032	0.015	0.010	0.014	0.032	0.015	0.010	0.014	0.032	0.015	0.010	0.014	0.032	0.015
Sunday	1	0.013	0.039	0.056	0.007	0.011	0.024	0.007	0.011	0.024	0.010	0.006	0.011	0.024	0.010	0.006	0.011	0.024	0.010	0.006	0.011	0.024	0.010
Sunday	2	0.010	0.039	0.052	0.005	0.011	0.022	0.005	0.011	0.022	0.007	0.004	0.012	0.022	0.007	0.004	0.012	0.022	0.007	0.004	0.012	0.022	0.007
Sunday	3	0.007	0.038	0.049	0.004	0.010	0.021	0.004	0.010	0.021	0.006	0.004	0.012	0.021	0.006	0.004	0.012	0.021	0.006	0.004	0.012	0.021	0.006
Sunday	4	0.007	0.037	0.046	0.004	0.010	0.020	0.004	0.010	0.020	0.006	0.005	0.017	0.020	0.006	0.005	0.017	0.020	0.006	0.005	0.017	0.020	0.006
Sunday	5	0.010	0.038	0.044	0.007	0.013	0.021	0.007	0.013	0.021	0.010	0.011	0.029	0.021	0.010	0.011	0.029	0.021	0.010	0.011	0.029	0.021	0.010
Sunday	6	0.016	0.038	0.043	0.012	0.019	0.026	0.012	0.019	0.026	0.016	0.017	0.037	0.026	0.016	0.017	0.037	0.026	0.016	0.017	0.037	0.026	0.016
Sunday	7	0.022	0.039	0.042	0.019	0.023	0.029	0.019	0.023	0.029	0.023	0.029	0.051	0.023	0.023	0.029	0.051	0.023	0.023	0.029	0.051	0.023	0.023
Sunday	8	0.032	0.040	0.041	0.032	0.035	0.038	0.032	0.035	0.038	0.033	0.043	0.071	0.032	0.035	0.038	0.033	0.043	0.071	0.033	0.040	0.042	0.033
Sunday	9	0.046	0.043	0.041	0.051	0.051	0.053	0.051	0.051	0.053	0.047	0.063	0.091	0.051	0.051	0.053	0.047	0.063	0.091	0.048	0.046	0.044	0.044
Sunday	10	0.059	0.046	0.041	0.067	0.067	0.071	0.067	0.067	0.071	0.057	0.075	0.084	0.067	0.075	0.084	0.057	0.075	0.084	0.062	0.051	0.045	0.045
Sunday	11	0.065	0.047	0.039	0.080	0.081	0.085	0.080	0.081	0.085	0.067	0.083	0.079	0.080	0.081	0.085	0.067	0.083	0.079	0.067	0.053	0.046	0.046
Sunday	12	0.069	0.048	0.038	0.083	0.081	0.076	0.083	0.081	0.076	0.074	0.090	0.070	0.083	0.081	0.076	0.074	0.090	0.070	0.070	0.054	0.046	0.046
Sunday	13	0.071	0.049	0.036	0.085	0.082	0.074	0.085	0.082	0.074	0.088	0.089	0.061	0.085	0.082	0.074	0.088	0.089	0.061	0.073	0.055	0.050	0.050
Sunday	14	0.072	0.049	0.035	0.085	0.083	0.069	0.085	0.083	0.069	0.079	0.081	0.057	0.085	0.083	0.069	0.079	0.081	0.057	0.073	0.055	0.047	0.047
Sunday	15	0.071	0.049	0.034	0.084	0.081	0.066	0.084	0.081	0.066	0.080	0.079	0.053	0.084	0.081	0.066	0.080	0.079	0.053	0.073	0.053	0.041	0.041
Sunday	16	0.070	0.048	0.033	0.082	0.079	0.060	0.082	0.079	0.060	0.082	0.079	0.045	0.082	0.079	0.060	0.082	0.079	0.045	0.072	0.052	0.039	0.039
Sunday	17	0.069	0.048	0.034	0.076	0.070	0.053	0.076	0.070	0.053	0.075	0.066	0.043	0.076	0.070	0.053	0.075	0.066	0.043	0.070	0.050	0.038	0.038
Sunday	18	0.063	0.045	0.033	0.064	0.056	0.043	0.064	0.056	0.043	0.066	0.054	0.039	0.064	0.056	0.043	0.066	0.054	0.039	0.063	0.047	0.036	0.036
Sunday	19	0.057	0.043	0.035	0.049	0.043	0.035	0.049	0.043	0.035	0.055	0.042	0.037	0.049	0.043	0.035	0.055	0.042	0.037	0.056	0.044	0.035	0.035
Sunday	20	0.052	0.041	0.036	0.038	0.033	0.024	0.038	0.033	0.024	0.045	0.031	0.030	0.038	0.033	0.024	0.045	0.031	0.030	0.051	0.041	0.036	0.036
Sunday	21	0.045	0.037	0.039	0.026	0.022	0.020	0.026	0.022	0.020	0.035	0.022	0.024	0.026	0.022	0.020	0.035	0.022	0.024	0.042	0.038	0.037	0.037
Sunday	22	0.033	0.032	0.043	0.017	0.014	0.017	0.017	0.014	0.017	0.023	0.013	0.018	0.017	0.014	0.017	0.023	0.013	0.018	0.030	0.032	0.039	0.039
Sunday	23	0.021	0.027	0.049	0.010	0.010	0.020	0.010	0.010	0.020	0.014	0.008	0.015	0.010	0.010	0.020	0.014	0.008	0.015	0.019	0.027	0.043	0.043
Monday	0	0.009	0.026	0.032	0.006	0.010	0.017	0.006	0.010	0.017	0.006	0.002	0.006	0.006	0.010	0.017	0.006	0.002	0.006	0.007	0.023	0.029	0.029
Monday	1	0.004	0.027	0.032	0.004	0.009	0.016	0.004	0.009	0.016	0.004	0.002	0.007	0.004	0.009	0.016	0.004	0.002	0.007	0.003	0.022	0.028	0.028
Monday	2	0.003	0.028	0.033	0.003	0.009	0.016	0.003	0.009	0.016	0.003	0.002	0.016	0.003	0.009	0.016	0.003	0.002	0.016	0.002	0.022	0.029	0.029
Monday	3	0.005	0.030	0.035	0.005	0.011	0.019	0.005	0.011	0.019	0.003	0.004	0.012	0.005	0.011	0.019	0.003	0.004	0.003	0.023	0.023	0.030	0.030
Monday	4	0.014	0.033	0.039	0.008	0.017	0.024	0.008	0.017	0.024	0.007	0.009	0.021	0.008	0.017	0.024	0.007	0.009	0.021	0.012	0.028	0.035	0.035
Monday	5	0.034	0.039	0.044	0.019	0.028	0.036	0.019	0.028	0.036	0.018	0.024	0.037	0.019	0.028	0.036	0.018	0.024	0.037	0.033	0.041	0.042	0.042
Monday	6	0.051	0.046	0.048	0.036	0.041	0.050	0.036	0.041	0.050	0.041	0.051	0.055	0.036	0.041	0.050	0.041	0.051	0.055	0.054	0.051	0.048	0.048
Monday	7	0.064	0.053	0.053	0.051	0.044	0.065	0.051	0.044	0.065	0.048	0.069	0.066	0.051	0.044	0.065	0.048	0.069	0.066	0.066	0.058	0.053	0.053
Monday	8	0.064	0.055	0.053	0.053	0.056	0.068	0.053	0.056	0.068	0.067	0.077	0.077	0.053	0.056	0.068	0.067	0.077	0.077	0.062	0.060	0.055	0.055
Monday	9	0.058	0.054	0.054	0.059	0.065	0.080	0.059	0.065	0.080	0.057	0.071	0.080	0.059	0.065	0.080	0.057	0.071	0.080	0.055	0.056	0.054	0.054
Monday	10	0.053	0.053	0.054	0.067	0.074	0.087	0.067	0.074	0.087	0.057	0.071	0.077	0.067	0.074	0.087	0.057	0.071	0.077	0.052	0.054	0.054	0.054
Monday	11	0.051	0.054	0.054	0.071	0.075	0.082	0.071	0.075	0.082	0.060	0.074	0.073	0.071	0.075	0.082	0.060	0.074	0.073	0.053	0.055	0.054	0.054
Monday	12	0.052	0.056	0.054	0.074	0.074	0.080	0.074	0.074	0.080	0.063	0.072	0.071	0.074	0.074	0.080	0.063	0.072	0.071	0.054	0.056	0.054	0.054

Day of Week	Hour	Alameda			Alpine			Amador			Butte			Calaveras			Colusa			Contra Costa		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Monday	13	0.054	0.057	0.054	0.074	0.075	0.075	0.074	0.075	0.075	0.063	0.072	0.068	0.074	0.075	0.075	0.063	0.072	0.068	0.056	0.056	0.054
Monday	14	0.061	0.059	0.053	0.077	0.076	0.065	0.077	0.076	0.065	0.067	0.077	0.064	0.077	0.076	0.065	0.067	0.077	0.064	0.063	0.059	0.056
Monday	15	0.066	0.059	0.051	0.082	0.076	0.058	0.081	0.073	0.045	0.078	0.080	0.056	0.082	0.076	0.058	0.078	0.080	0.056	0.069	0.063	0.058
Monday	16	0.069	0.057	0.048	0.081	0.073	0.045	0.081	0.073	0.045	0.086	0.077	0.049	0.081	0.073	0.045	0.086	0.077	0.049	0.072	0.060	0.052
Monday	17	0.070	0.053	0.044	0.071	0.059	0.035	0.071	0.059	0.035	0.087	0.062	0.041	0.071	0.059	0.035	0.087	0.062	0.041	0.073	0.056	0.047
Monday	18	0.062	0.045	0.037	0.052	0.042	0.023	0.052	0.042	0.023	0.051	0.038	0.030	0.052	0.042	0.023	0.051	0.038	0.030	0.061	0.045	0.039
Monday	19	0.048	0.035	0.031	0.037	0.030	0.017	0.037	0.030	0.017	0.036	0.024	0.024	0.037	0.030	0.017	0.036	0.024	0.024	0.045	0.033	0.031
Monday	20	0.036	0.028	0.026	0.027	0.022	0.013	0.027	0.022	0.013	0.026	0.018	0.023	0.027	0.022	0.013	0.026	0.018	0.023	0.035	0.026	0.026
Monday	21	0.031	0.022	0.023	0.020	0.016	0.010	0.020	0.016	0.010	0.020	0.012	0.021	0.020	0.016	0.010	0.020	0.012	0.021	0.031	0.022	0.024
Monday	22	0.024	0.018	0.023	0.015	0.012	0.009	0.015	0.012	0.009	0.015	0.007	0.015	0.015	0.012	0.009	0.015	0.007	0.015	0.023	0.017	0.023
Monday	23	0.016	0.015	0.025	0.009	0.007	0.010	0.009	0.007	0.010	0.008	0.004	0.015	0.009	0.007	0.010	0.008	0.004	0.015	0.014	0.014	0.025
Tues/Wed/Thurs	0	0.008	0.026	0.034	0.005	0.009	0.017	0.005	0.009	0.017	0.006	0.003	0.010	0.005	0.009	0.017	0.006	0.003	0.010	0.006	0.022	0.031
Tues/Wed/Thurs	1	0.004	0.027	0.034	0.003	0.008	0.017	0.003	0.008	0.017	0.003	0.002	0.011	0.003	0.008	0.017	0.003	0.002	0.011	0.003	0.021	0.030
Tues/Wed/Thurs	2	0.003	0.028	0.035	0.002	0.009	0.017	0.002	0.009	0.017	0.003	0.002	0.013	0.002	0.009	0.017	0.003	0.002	0.013	0.002	0.021	0.030
Tues/Wed/Thurs	3	0.005	0.030	0.037	0.003	0.010	0.022	0.003	0.010	0.022	0.003	0.003	0.015	0.003	0.010	0.022	0.003	0.003	0.015	0.003	0.023	0.031
Tues/Wed/Thurs	4	0.014	0.034	0.041	0.006	0.014	0.025	0.006	0.014	0.025	0.006	0.008	0.022	0.006	0.014	0.025	0.006	0.008	0.022	0.011	0.028	0.036
Tues/Wed/Thurs	5	0.035	0.040	0.046	0.018	0.027	0.039	0.018	0.027	0.039	0.017	0.024	0.037	0.018	0.027	0.039	0.017	0.024	0.037	0.034	0.040	0.044
Tues/Wed/Thurs	6	0.055	0.047	0.050	0.037	0.042	0.052	0.037	0.042	0.052	0.041	0.053	0.054	0.037	0.042	0.052	0.041	0.053	0.054	0.056	0.052	0.049
Tues/Wed/Thurs	7	0.067	0.054	0.053	0.053	0.047	0.064	0.053	0.047	0.064	0.053	0.069	0.066	0.053	0.047	0.064	0.053	0.069	0.066	0.068	0.059	0.054
Tues/Wed/Thurs	8	0.064	0.056	0.054	0.054	0.056	0.070	0.054	0.056	0.070	0.066	0.077	0.077	0.054	0.056	0.070	0.066	0.077	0.063	0.060	0.056	0.056
Tues/Wed/Thurs	9	0.057	0.054	0.055	0.059	0.068	0.083	0.059	0.068	0.083	0.057	0.071	0.080	0.059	0.068	0.083	0.057	0.071	0.080	0.055	0.055	0.053
Tues/Wed/Thurs	10	0.051	0.053	0.054	0.064	0.069	0.081	0.064	0.069	0.081	0.056	0.071	0.077	0.064	0.069	0.081	0.056	0.071	0.077	0.051	0.053	0.052
Tues/Wed/Thurs	11	0.049	0.054	0.054	0.068	0.069	0.077	0.068	0.069	0.077	0.058	0.071	0.074	0.068	0.069	0.077	0.058	0.071	0.074	0.050	0.054	0.052
Tues/Wed/Thurs	12	0.050	0.055	0.054	0.069	0.071	0.074	0.069	0.071	0.074	0.062	0.070	0.069	0.069	0.071	0.074	0.062	0.070	0.069	0.052	0.055	0.053
Tues/Wed/Thurs	13	0.053	0.056	0.053	0.072	0.073	0.074	0.072	0.073	0.074	0.063	0.073	0.067	0.072	0.073	0.074	0.063	0.073	0.067	0.054	0.056	0.054
Tues/Wed/Thurs	14	0.060	0.058	0.052	0.077	0.076	0.067	0.077	0.076	0.067	0.066	0.076	0.063	0.077	0.076	0.067	0.066	0.076	0.063	0.062	0.059	0.054
Tues/Wed/Thurs	15	0.064	0.058	0.050	0.084	0.078	0.058	0.084	0.078	0.058	0.079	0.080	0.056	0.084	0.078	0.058	0.079	0.080	0.056	0.067	0.063	0.056
Tues/Wed/Thurs	16	0.067	0.056	0.047	0.082	0.074	0.048	0.082	0.074	0.048	0.087	0.076	0.045	0.082	0.074	0.048	0.087	0.076	0.045	0.070	0.060	0.051
Tues/Wed/Thurs	17	0.067	0.052	0.042	0.074	0.061	0.036	0.074	0.061	0.036	0.088	0.062	0.040	0.074	0.061	0.036	0.088	0.062	0.040	0.071	0.057	0.046
Tues/Wed/Thurs	18	0.061	0.044	0.036	0.053	0.044	0.023	0.053	0.044	0.023	0.054	0.039	0.031	0.053	0.044	0.023	0.054	0.039	0.031	0.062	0.047	0.039
Tues/Wed/Thurs	19	0.050	0.035	0.030	0.038	0.031	0.016	0.038	0.031	0.016	0.036	0.026	0.023	0.038	0.031	0.016	0.036	0.026	0.023	0.048	0.035	0.031
Tues/Wed/Thurs	20	0.038	0.027	0.025	0.030	0.025	0.012	0.030	0.025	0.012	0.028	0.019	0.021	0.030	0.025	0.012	0.028	0.019	0.021	0.038	0.027	0.026
Tues/Wed/Thurs	21	0.033	0.022	0.022	0.023	0.018	0.010	0.023	0.018	0.010	0.021	0.013	0.020	0.023	0.018	0.010	0.021	0.013	0.020	0.033	0.022	0.024
Tues/Wed/Thurs	22	0.026	0.017	0.022	0.017	0.013	0.010	0.017	0.013	0.010	0.014	0.007	0.016	0.017	0.013	0.010	0.014	0.007	0.016	0.024	0.017	0.022
Tues/Wed/Thurs	23	0.016	0.014	0.023	0.010	0.008	0.010	0.010	0.008	0.010	0.009	0.004	0.013	0.010	0.008	0.010	0.009	0.004	0.013	0.015	0.013	0.024
Friday	0	0.009	0.027	0.036	0.005	0.009	0.019	0.005	0.009	0.019	0.007	0.003	0.011	0.005	0.009	0.019	0.007	0.003	0.011	0.008	0.022	0.033
Friday	1	0.005	0.028	0.037	0.003	0.008	0.019	0.003	0.008	0.019	0.004	0.003	0.012	0.003	0.008	0.019	0.004	0.003	0.012	0.004	0.021	0.031
Friday	2	0.004	0.029	0.038	0.002	0.008	0.019	0.002	0.008	0.019	0.004	0.003	0.015	0.002	0.008	0.019	0.004	0.003	0.015	0.003	0.022	0.032
Friday	3	0.005	0.031	0.039	0.002	0.008	0.021	0.002	0.008	0.021	0.004	0.004	0.017	0.002	0.008	0.021	0.004	0.004	0.017	0.004	0.023	0.033
Friday	4	0.013	0.034	0.043	0.005	0.013	0.024	0.005	0.013	0.024	0.006	0.007	0.024	0.005	0.013	0.024	0.006	0.007	0.024	0.010	0.028	0.036
Friday	5	0.032	0.040	0.048	0.013	0.023	0.037	0.013	0.023	0.037	0.015	0.022	0.039	0.013	0.023	0.037	0.015	0.022	0.039	0.030	0.039	0.044
Friday	6	0.049	0.046	0.052	0.026	0.035	0.049	0.026	0.035	0.049	0.035	0.045	0.055	0.026	0.035	0.049	0.035	0.045	0.055	0.050	0.049	0.050
Friday	7	0.060	0.052	0.055	0.039	0.040	0.060	0.039	0.040	0.060	0.063	0.063	0.064	0.039	0.040	0.060	0.063	0.063	0.064	0.063	0.057	0.055
Friday	8	0.059	0.054	0.056	0.043	0.049	0.068	0.043	0.049	0.068	0.058	0.072	0.074	0.043	0.049	0.068	0.058	0.072	0.074	0.059	0.057	0.056
Friday	9	0.054	0.053	0.056	0.049	0.057	0.073	0.049	0.057	0.073	0.052	0.068	0.075	0.049	0.057	0.073	0.052	0.068	0.075	0.053	0.054	0.054
Friday	10	0.051	0.053	0.056	0.058	0.063	0.078	0.058	0.063	0.078	0.055	0.071	0.074	0.058	0.063	0.078	0.055	0.071	0.074	0.051	0.053	0.053
Friday	11	0.052	0.055	0.055	0.064	0.069	0.077	0.064	0.069	0.077	0.060	0.074	0.074	0.064	0.069	0.077	0.060	0.074	0.074	0.053	0.055	0.054
Friday	12	0.054	0.056	0.055	0.066	0.071	0.076	0.066	0.071	0.076	0.063	0.072	0.069	0.066	0.071	0.076	0.063	0.072	0.069	0.056	0.057	0.055
Friday	13	0.056	0.057	0.054	0.071	0.074	0.077	0.071	0.074	0.077	0.065	0.076	0.069	0.071	0.074	0.077	0.065	0.076	0.069	0.058	0.058	0.056
Friday	14	0.061	0.058	0.052	0.076	0.077	0.070	0.076	0.077	0.070	0.069	0.078	0.063	0.076	0.077	0.070	0.069	0.078	0.063	0.064	0.064	0.056
Friday	15	0.063	0.058	0.049	0.083	0.079	0.060	0.083	0.079	0.060	0.078	0.080	0.055	0.083	0.079	0.060	0.078	0.080	0.055	0.066	0.062	0.056
Friday	16	0.064	0.055	0.045	0.083	0.077	0.050	0.083	0.077	0.050	0.085	0.075	0.047	0.083	0.077	0.050	0.085	0.075	0.047	0.067	0.059	0.050

Day of Week	Hour	Alameda			Alpine			Amador			Butte			Calaveras			Colusa			Contra Costa			
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	
Friday	17	0.064	0.051	0.040	0.064	0.038	0.038	0.075	0.064	0.038	0.061	0.039	0.082	0.075	0.064	0.038	0.082	0.061	0.039	0.067	0.055	0.046	
Friday	18	0.059	0.044	0.034	0.062	0.051	0.025	0.062	0.051	0.025	0.041	0.029	0.059	0.041	0.029	0.041	0.029	0.059	0.041	0.029	0.060	0.047	0.039
Friday	19	0.052	0.035	0.027	0.050	0.039	0.018	0.050	0.039	0.018	0.042	0.024	0.042	0.050	0.039	0.018	0.042	0.024	0.024	0.049	0.049	0.036	0.030
Friday	20	0.042	0.028	0.022	0.041	0.030	0.013	0.041	0.030	0.013	0.032	0.021	0.032	0.041	0.030	0.013	0.032	0.021	0.021	0.040	0.040	0.029	0.023
Friday	21	0.036	0.023	0.019	0.036	0.025	0.010	0.036	0.025	0.010	0.027	0.015	0.027	0.036	0.025	0.010	0.027	0.015	0.020	0.035	0.023	0.020	0.020
Friday	22	0.032	0.019	0.017	0.030	0.019	0.011	0.030	0.019	0.011	0.030	0.016	0.030	0.019	0.011	0.030	0.016	0.016	0.030	0.030	0.019	0.019	0.019
Friday	23	0.014	0.018	0.018	0.018	0.009	0.009	0.018	0.012	0.009	0.014	0.007	0.015	0.018	0.012	0.009	0.014	0.007	0.015	0.022	0.015	0.020	0.020
Saturday	1	0.016	0.033	0.052	0.010	0.015	0.027	0.010	0.015	0.027	0.012	0.007	0.021	0.010	0.015	0.027	0.012	0.007	0.021	0.015	0.030	0.044	0.044
Saturday	2	0.010	0.033	0.051	0.007	0.012	0.023	0.007	0.012	0.023	0.008	0.005	0.016	0.007	0.012	0.023	0.008	0.005	0.016	0.009	0.027	0.040	0.040
Saturday	3	0.008	0.033	0.049	0.005	0.011	0.022	0.005	0.011	0.022	0.006	0.004	0.020	0.005	0.011	0.022	0.006	0.004	0.020	0.006	0.026	0.039	0.039
Saturday	4	0.006	0.034	0.048	0.004	0.010	0.025	0.004	0.010	0.025	0.005	0.004	0.022	0.004	0.010	0.025	0.005	0.004	0.022	0.005	0.025	0.037	0.037
Saturday	5	0.008	0.035	0.048	0.005	0.013	0.028	0.005	0.013	0.028	0.006	0.008	0.024	0.005	0.013	0.028	0.006	0.008	0.024	0.006	0.027	0.037	0.037
Saturday	6	0.014	0.037	0.049	0.010	0.021	0.034	0.010	0.021	0.034	0.012	0.017	0.039	0.010	0.021	0.034	0.012	0.017	0.039	0.013	0.030	0.040	0.040
Saturday	7	0.023	0.039	0.050	0.017	0.028	0.039	0.017	0.028	0.039	0.021	0.028	0.049	0.017	0.028	0.039	0.021	0.028	0.049	0.023	0.035	0.042	0.042
Saturday	8	0.033	0.041	0.051	0.029	0.036	0.053	0.029	0.036	0.053	0.034	0.041	0.058	0.029	0.036	0.053	0.034	0.041	0.058	0.034	0.041	0.047	0.047
Saturday	9	0.045	0.044	0.052	0.044	0.045	0.060	0.044	0.045	0.060	0.045	0.057	0.067	0.044	0.045	0.060	0.045	0.057	0.067	0.046	0.047	0.049	0.049
Saturday	10	0.054	0.047	0.052	0.059	0.061	0.071	0.059	0.061	0.071	0.054	0.068	0.074	0.059	0.061	0.071	0.054	0.068	0.074	0.055	0.051	0.050	0.050
Saturday	11	0.060	0.050	0.051	0.073	0.074	0.078	0.073	0.074	0.078	0.063	0.080	0.073	0.073	0.074	0.078	0.063	0.080	0.073	0.061	0.054	0.051	0.051
Saturday	12	0.066	0.053	0.048	0.081	0.077	0.083	0.081	0.077	0.083	0.068	0.082	0.071	0.081	0.077	0.083	0.068	0.082	0.071	0.065	0.056	0.052	0.052
Saturday	13	0.066	0.053	0.045	0.075	0.072	0.060	0.075	0.072	0.060	0.074	0.079	0.062	0.075	0.072	0.060	0.074	0.079	0.062	0.067	0.059	0.058	0.058
Saturday	14	0.066	0.053	0.042	0.075	0.068	0.055	0.075	0.068	0.055	0.074	0.076	0.057	0.075	0.068	0.055	0.074	0.076	0.057	0.068	0.058	0.057	0.057
Saturday	15	0.066	0.053	0.040	0.075	0.068	0.052	0.075	0.068	0.052	0.073	0.074	0.052	0.075	0.068	0.052	0.073	0.074	0.052	0.068	0.057	0.051	0.051
Saturday	16	0.065	0.051	0.037	0.072	0.070	0.047	0.072	0.070	0.047	0.067	0.045	0.067	0.072	0.070	0.047	0.067	0.045	0.067	0.068	0.056	0.047	0.047
Saturday	17	0.065	0.050	0.034	0.066	0.063	0.040	0.066	0.063	0.040	0.069	0.058	0.040	0.066	0.063	0.040	0.069	0.058	0.040	0.067	0.054	0.044	0.044
Saturday	18	0.060	0.046	0.031	0.058	0.052	0.031	0.058	0.052	0.031	0.058	0.047	0.034	0.058	0.052	0.031	0.058	0.047	0.034	0.060	0.048	0.036	0.036
Saturday	19	0.050	0.041	0.028	0.047	0.041	0.026	0.047	0.041	0.026	0.046	0.036	0.029	0.047	0.041	0.026	0.046	0.036	0.029	0.049	0.041	0.029	0.029
Saturday	20	0.043	0.036	0.025	0.038	0.031	0.020	0.038	0.031	0.020	0.040	0.028	0.024	0.038	0.031	0.020	0.040	0.028	0.024	0.043	0.036	0.024	0.024
Saturday	21	0.042	0.033	0.024	0.031	0.025	0.016	0.031	0.025	0.016	0.036	0.022	0.023	0.031	0.025	0.016	0.036	0.022	0.023	0.041	0.033	0.024	0.024
Saturday	22	0.039	0.029	0.023	0.025	0.020	0.018	0.025	0.020	0.018	0.029	0.016	0.017	0.025	0.020	0.018	0.029	0.016	0.017	0.037	0.029	0.023	0.023
Saturday	23	0.029	0.025	0.023	0.016	0.013	0.018	0.016	0.013	0.018	0.020	0.011	0.017	0.016	0.013	0.018	0.020	0.011	0.017	0.028	0.024	0.022	0.022
Holiday	0	0.015	0.028	0.035	0.008	0.011	0.020	0.008	0.011	0.020	0.004	0.004	0.012	0.008	0.011	0.020	0.004	0.004	0.012	0.013	0.027	0.034	0.034
Holiday	1	0.008	0.029	0.035	0.005	0.009	0.018	0.005	0.009	0.018	0.006	0.004	0.011	0.005	0.009	0.018	0.006	0.004	0.011	0.007	0.026	0.033	0.033
Holiday	2	0.006	0.031	0.036	0.003	0.010	0.018	0.003	0.010	0.018	0.004	0.003	0.012	0.003	0.010	0.018	0.004	0.003	0.012	0.004	0.025	0.033	0.033
Holiday	3	0.005	0.032	0.037	0.004	0.010	0.021	0.004	0.010	0.021	0.004	0.005	0.015	0.004	0.010	0.021	0.004	0.005	0.015	0.003	0.025	0.033	0.033
Holiday	4	0.009	0.035	0.040	0.005	0.012	0.020	0.005	0.012	0.020	0.007	0.009	0.024	0.005	0.012	0.020	0.007	0.009	0.024	0.007	0.029	0.035	0.035
Holiday	5	0.019	0.037	0.043	0.009	0.018	0.031	0.009	0.018	0.031	0.014	0.020	0.037	0.009	0.018	0.031	0.014	0.020	0.037	0.017	0.034	0.039	0.039
Holiday	6	0.029	0.042	0.045	0.018	0.023	0.038	0.018	0.023	0.038	0.030	0.036	0.047	0.018	0.023	0.038	0.030	0.036	0.047	0.029	0.040	0.044	0.044
Holiday	7	0.038	0.046	0.048	0.029	0.031	0.043	0.029	0.031	0.043	0.044	0.052	0.061	0.029	0.031	0.043	0.044	0.052	0.061	0.038	0.045	0.047	0.047
Holiday	8	0.046	0.049	0.051	0.041	0.044	0.056	0.041	0.044	0.056	0.052	0.066	0.075	0.041	0.044	0.056	0.052	0.066	0.075	0.045	0.050	0.051	0.051
Holiday	9	0.049	0.050	0.052	0.058	0.057	0.075	0.058	0.057	0.075	0.053	0.071	0.081	0.058	0.057	0.075	0.053	0.071	0.081	0.049	0.053	0.052	0.052
Holiday	10	0.055	0.053	0.053	0.076	0.083	0.087	0.076	0.083	0.087	0.066	0.076	0.081	0.076	0.083	0.087	0.066	0.076	0.081	0.056	0.056	0.053	0.053
Holiday	11	0.060	0.056	0.054	0.084	0.086	0.088	0.084	0.086	0.088	0.066	0.076	0.081	0.084	0.086	0.088	0.066	0.076	0.081	0.062	0.059	0.055	0.055
Holiday	12	0.064	0.058	0.055	0.085	0.087	0.089	0.085	0.087	0.089	0.071	0.078	0.074	0.085	0.087	0.089	0.071	0.078	0.074	0.067	0.061	0.056	0.056
Holiday	13	0.066	0.059	0.054	0.083	0.081	0.078	0.083	0.081	0.078	0.071	0.076	0.065	0.083	0.081	0.078	0.071	0.076	0.065	0.070	0.062	0.056	0.056
Holiday	14	0.069	0.060	0.053	0.080	0.074	0.068	0.080	0.074	0.068	0.070	0.078	0.060	0.080	0.074	0.068	0.070	0.078	0.060	0.073	0.062	0.057	0.057
Holiday	15	0.069	0.058	0.051	0.078	0.074	0.060	0.078	0.074	0.060	0.075	0.075	0.053	0.078	0.074	0.060	0.075	0.075	0.053	0.071	0.061	0.054	0.054
Holiday	16	0.068	0.056	0.047	0.078	0.072	0.049	0.078	0.072	0.049	0.079	0.070	0.044	0.078	0.072	0.049	0.079	0.070	0.044	0.070	0.057	0.050	0.050
Holiday	17	0.066	0.051	0.043	0.071	0.066	0.041	0.071	0.066	0.041	0.074	0.064	0.041	0.071	0.066	0.041	0.074	0.064	0.041	0.067	0.053	0.044	0.044
Holiday	18	0.060	0.044	0.034	0.057	0.049	0.033	0.057	0.049	0.033	0.058	0.044	0.034	0.057	0.049	0.033	0.058	0.044	0.034	0.059	0.045	0.038	0.038
Holiday	19	0.052	0.036	0.031	0.043	0.040	0.022	0.043	0.040	0.022	0.047	0.033	0.026	0.043	0.040	0.022	0.047	0.033	0.026	0.051	0.036	0.031	0.031
Holiday	20	0.046	0.030	0.027	0.033	0.026	0.013	0.033	0.026														

Day of Week	Alameda			Alpine			Amador			Butte			Calaveras			Colusa			Contra Costa			
	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	
Holiday	21	0.042	0.025	0.024	0.018	0.011	0.024	0.018	0.011	0.030	0.018	0.021	0.024	0.018	0.011	0.030	0.018	0.021	0.041	0.026	0.026	
Holiday	22	0.035	0.020	0.024	0.017	0.012	0.009	0.017	0.012	0.024	0.011	0.017	0.017	0.012	0.009	0.024	0.011	0.017	0.033	0.021	0.025	
Holiday	23	0.024	0.016	0.026	0.010	0.008	0.010	0.010	0.008	0.014	0.007	0.014	0.010	0.008	0.010	0.014	0.007	0.014	0.021	0.017	0.026	
Day of Week	Hour	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Sunday	0	0.013	0.011	0.008	0.013	0.020	0.031	0.015	0.033	0.043	0.015	0.010	0.015	0.013	0.011	0.008	0.026	0.015	0.017	0.010	0.014	0.032
Sunday	1	0.013	0.008	0.010	0.008	0.016	0.028	0.010	0.030	0.040	0.010	0.006	0.011	0.013	0.008	0.010	0.026	0.013	0.016	0.007	0.011	0.024
Sunday	2	0.012	0.006	0.008	0.006	0.013	0.026	0.008	0.027	0.037	0.007	0.004	0.012	0.012	0.006	0.008	0.025	0.009	0.014	0.005	0.011	0.022
Sunday	3	0.014	0.005	0.007	0.005	0.012	0.025	0.005	0.025	0.034	0.006	0.004	0.012	0.014	0.005	0.007	0.025	0.008	0.015	0.004	0.010	0.021
Sunday	4	0.014	0.004	0.011	0.005	0.012	0.025	0.006	0.024	0.034	0.006	0.005	0.017	0.014	0.004	0.011	0.027	0.010	0.015	0.004	0.010	0.020
Sunday	5	0.017	0.009	0.019	0.008	0.015	0.027	0.010	0.026	0.034	0.010	0.011	0.029	0.017	0.009	0.019	0.030	0.015	0.017	0.007	0.013	0.021
Sunday	6	0.021	0.014	0.028	0.013	0.020	0.030	0.017	0.029	0.036	0.016	0.017	0.037	0.021	0.014	0.028	0.032	0.019	0.021	0.012	0.019	0.026
Sunday	7	0.026	0.020	0.036	0.022	0.028	0.034	0.022	0.032	0.037	0.023	0.029	0.051	0.026	0.020	0.036	0.033	0.026	0.029	0.019	0.023	0.029
Sunday	8	0.031	0.032	0.043	0.034	0.041	0.040	0.032	0.038	0.040	0.033	0.043	0.071	0.031	0.032	0.043	0.037	0.039	0.035	0.032	0.035	0.038
Sunday	9	0.040	0.050	0.054	0.048	0.055	0.046	0.044	0.046	0.044	0.047	0.063	0.091	0.040	0.050	0.054	0.040	0.053	0.047	0.051	0.051	0.053
Sunday	10	0.047	0.064	0.067	0.064	0.068	0.052	0.055	0.052	0.046	0.057	0.075	0.084	0.047	0.064	0.067	0.043	0.063	0.057	0.067	0.067	0.071
Sunday	11	0.055	0.079	0.062	0.075	0.075	0.055	0.063	0.057	0.047	0.067	0.083	0.079	0.055	0.079	0.062	0.046	0.071	0.065	0.080	0.081	0.085
Sunday	12	0.061	0.087	0.065	0.082	0.079	0.058	0.071	0.062	0.049	0.074	0.090	0.070	0.061	0.087	0.065	0.048	0.075	0.068	0.083	0.081	0.076
Sunday	13	0.065	0.092	0.064	0.084	0.079	0.058	0.076	0.064	0.049	0.078	0.089	0.061	0.065	0.092	0.064	0.052	0.078	0.068	0.085	0.082	0.074
Sunday	14	0.067	0.087	0.065	0.084	0.077	0.057	0.077	0.063	0.048	0.079	0.081	0.057	0.067	0.087	0.065	0.053	0.074	0.065	0.085	0.083	0.069
Sunday	15	0.072	0.086	0.067	0.082	0.073	0.057	0.077	0.061	0.047	0.080	0.079	0.053	0.072	0.086	0.067	0.056	0.071	0.061	0.084	0.081	0.066
Sunday	16	0.077	0.086	0.072	0.079	0.068	0.055	0.075	0.059	0.046	0.079	0.075	0.045	0.077	0.086	0.067	0.056	0.068	0.058	0.082	0.079	0.060
Sunday	17	0.070	0.075	0.058	0.072	0.062	0.053	0.073	0.056	0.045	0.075	0.066	0.043	0.070	0.075	0.058	0.059	0.067	0.055	0.076	0.070	0.053
Sunday	18	0.067	0.059	0.054	0.060	0.052	0.049	0.066	0.050	0.044	0.066	0.054	0.039	0.067	0.059	0.054	0.059	0.062	0.055	0.064	0.056	0.043
Sunday	19	0.062	0.045	0.050	0.050	0.043	0.045	0.057	0.044	0.042	0.042	0.042	0.037	0.062	0.045	0.050	0.057	0.051	0.051	0.049	0.043	0.035
Sunday	20	0.054	0.035	0.047	0.041	0.035	0.042	0.050	0.038	0.041	0.045	0.031	0.030	0.054	0.035	0.047	0.052	0.041	0.049	0.038	0.033	0.024
Sunday	21	0.045	0.024	0.039	0.031	0.026	0.039	0.040	0.033	0.040	0.035	0.022	0.024	0.045	0.024	0.039	0.047	0.032	0.044	0.026	0.022	0.020
Sunday	22	0.033	0.015	0.033	0.021	0.019	0.036	0.030	0.028	0.040	0.023	0.013	0.018	0.033	0.024	0.033	0.039	0.023	0.042	0.017	0.014	0.017
Sunday	23	0.022	0.009	0.032	0.013	0.015	0.033	0.020	0.023	0.039	0.014	0.008	0.015	0.022	0.009	0.032	0.031	0.018	0.038	0.010	0.010	0.020
Monday	0	0.010	0.003	0.007	0.008	0.014	0.027	0.009	0.019	0.024	0.006	0.002	0.006	0.010	0.003	0.007	0.025	0.010	0.016	0.006	0.010	0.017
Monday	1	0.009	0.002	0.007	0.005	0.012	0.025	0.005	0.018	0.023	0.004	0.002	0.007	0.009	0.002	0.007	0.025	0.008	0.016	0.004	0.009	0.016
Monday	2	0.010	0.003	0.010	0.004	0.012	0.025	0.004	0.018	0.023	0.003	0.002	0.010	0.010	0.003	0.010	0.024	0.008	0.017	0.003	0.009	0.016
Monday	3	0.012	0.006	0.012	0.006	0.014	0.027	0.005	0.020	0.025	0.003	0.004	0.012	0.012	0.006	0.012	0.030	0.014	0.019	0.005	0.011	0.019
Monday	4	0.014	0.009	0.013	0.011	0.019	0.030	0.011	0.023	0.027	0.007	0.009	0.021	0.014	0.009	0.013	0.030	0.022	0.025	0.008	0.017	0.024
Monday	5	0.022	0.022	0.026	0.023	0.030	0.036	0.024	0.034	0.033	0.018	0.024	0.037	0.022	0.022	0.026	0.034	0.036	0.031	0.019	0.028	0.036
Monday	6	0.037	0.047	0.044	0.042	0.047	0.043	0.044	0.047	0.041	0.041	0.051	0.055	0.037	0.047	0.044	0.036	0.043	0.034	0.036	0.041	0.050
Monday	7	0.045	0.058	0.058	0.060	0.061	0.048	0.069	0.064	0.048	0.078	0.069	0.066	0.045	0.058	0.058	0.040	0.056	0.039	0.051	0.044	0.065
Monday	8	0.047	0.062	0.067	0.059	0.062	0.050	0.063	0.062	0.049	0.067	0.077	0.077	0.047	0.065	0.048	0.041	0.045	0.045	0.053	0.056	0.068
Monday	9	0.050	0.065	0.078	0.056	0.061	0.050	0.055	0.056	0.047	0.057	0.071	0.080	0.050	0.065	0.078	0.043	0.064	0.051	0.059	0.065	0.080
Monday	10	0.051	0.065	0.080	0.058	0.064	0.051	0.055	0.056	0.048	0.057	0.071	0.077	0.051	0.065	0.080	0.044	0.069	0.058	0.067	0.074	0.087
Monday	11	0.056	0.067	0.083	0.062	0.068	0.053	0.057	0.059	0.050	0.060	0.074	0.073	0.058	0.067	0.083	0.048	0.071	0.066	0.071	0.075	0.082
Monday	12	0.058	0.069	0.081	0.066	0.068	0.054	0.061	0.061	0.052	0.063	0.072	0.071	0.058	0.069	0.081	0.048	0.068	0.067	0.074	0.074	0.080
Monday	13	0.063	0.074	0.076	0.067	0.067	0.054	0.063	0.062	0.054	0.063	0.072	0.068	0.063	0.074	0.076	0.050	0.070	0.067	0.074	0.075	0.075
Monday	14	0.067	0.076	0.074	0.070	0.069	0.055	0.069	0.065	0.056	0.067	0.077	0.064	0.067	0.076	0.074	0.051	0.069	0.066	0.077	0.076	0.068
Monday	15	0.073	0.087	0.062	0.073	0.069	0.055	0.074	0.068	0.058	0.078	0.080	0.056	0.073	0.087	0.062	0.057	0.072	0.062	0.082	0.076	0.058
Monday	16	0.076	0.084	0.053	0.075	0.067	0.054	0.079	0.068	0.059	0.086	0.077	0.049	0.076	0.084	0.053	0.054	0.063	0.061	0.081	0.073	0.045
Monday	17	0.075	0.075	0.040	0.073	0.061	0.052	0.076	0.062	0.057	0.087	0.062	0.041	0.075	0.075	0.040	0.057	0.054	0.055	0.071	0.059	0.035

Day of Week	Hour	Del Norte			El Dorado			Fresno			Glenn			Humboldt			Imperial			Inyo		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Monday	18	0.057	0.047	0.032	0.056	0.046	0.045	0.053	0.043	0.050	0.038	0.030	0.057	0.047	0.032	0.054	0.040	0.047	0.052	0.042	0.023	
Monday	19	0.050	0.031	0.039	0.040	0.031	0.039	0.037	0.030	0.043	0.036	0.024	0.050	0.031	0.029	0.052	0.032	0.041	0.037	0.030	0.017	
Monday	20	0.043	0.020	0.021	0.031	0.022	0.035	0.030	0.023	0.039	0.026	0.018	0.036	0.018	0.021	0.047	0.022	0.037	0.020	0.022	0.013	
Monday	21	0.035	0.015	0.020	0.025	0.017	0.032	0.024	0.018	0.035	0.020	0.012	0.035	0.015	0.020	0.045	0.018	0.031	0.020	0.016	0.010	
Monday	22	0.025	0.009	0.014	0.017	0.012	0.030	0.018	0.013	0.032	0.013	0.007	0.025	0.009	0.014	0.038	0.013	0.026	0.015	0.012	0.009	
Monday	23	0.016	0.005	0.013	0.012	0.009	0.030	0.012	0.010	0.029	0.008	0.015	0.016	0.005	0.013	0.030	0.014	0.025	0.009	0.007	0.010	
Tues/Wed/Thurs	0	0.004	0.004	0.008	0.008	0.014	0.029	0.007	0.018	0.027	0.006	0.011	0.010	0.004	0.008	0.024	0.011	0.023	0.005	0.009	0.017	
Tues/Wed/Thurs	1	0.009	0.003	0.008	0.004	0.011	0.027	0.004	0.017	0.027	0.003	0.002	0.009	0.003	0.008	0.025	0.009	0.020	0.003	0.008	0.017	
Tues/Wed/Thurs	2	0.010	0.002	0.012	0.004	0.011	0.027	0.003	0.017	0.027	0.003	0.013	0.010	0.002	0.012	0.026	0.008	0.020	0.002	0.009	0.017	
Tues/Wed/Thurs	3	0.015	0.005	0.014	0.005	0.013	0.029	0.004	0.019	0.028	0.003	0.015	0.011	0.005	0.014	0.027	0.012	0.022	0.003	0.010	0.022	
Tues/Wed/Thurs	4	0.015	0.010	0.018	0.010	0.018	0.031	0.009	0.023	0.031	0.006	0.022	0.015	0.010	0.021	0.029	0.018	0.025	0.006	0.014	0.025	
Tues/Wed/Thurs	5	0.024	0.024	0.035	0.022	0.029	0.037	0.024	0.032	0.036	0.017	0.024	0.024	0.024	0.035	0.034	0.036	0.032	0.018	0.027	0.039	
Tues/Wed/Thurs	6	0.037	0.048	0.048	0.042	0.047	0.044	0.044	0.047	0.044	0.041	0.053	0.037	0.048	0.048	0.036	0.046	0.039	0.037	0.042	0.052	
Tues/Wed/Thurs	7	0.045	0.059	0.065	0.060	0.061	0.050	0.070	0.064	0.051	0.077	0.069	0.045	0.059	0.065	0.040	0.057	0.044	0.053	0.047	0.064	
Tues/Wed/Thurs	8	0.047	0.063	0.069	0.060	0.062	0.051	0.065	0.063	0.051	0.066	0.077	0.047	0.063	0.069	0.041	0.065	0.048	0.054	0.056	0.070	
Tues/Wed/Thurs	9	0.050	0.064	0.074	0.055	0.060	0.050	0.055	0.057	0.049	0.057	0.071	0.050	0.050	0.064	0.074	0.041	0.062	0.059	0.068	0.083	
Tues/Wed/Thurs	10	0.051	0.065	0.075	0.056	0.061	0.051	0.054	0.056	0.050	0.056	0.071	0.051	0.065	0.075	0.044	0.066	0.057	0.064	0.069	0.081	
Tues/Wed/Thurs	11	0.055	0.065	0.076	0.059	0.064	0.052	0.055	0.058	0.051	0.058	0.071	0.055	0.065	0.076	0.046	0.067	0.061	0.068	0.069	0.077	
Tues/Wed/Thurs	12	0.057	0.068	0.076	0.061	0.065	0.053	0.058	0.060	0.051	0.062	0.070	0.057	0.068	0.076	0.048	0.067	0.064	0.069	0.071	0.074	
Tues/Wed/Thurs	13	0.061	0.070	0.071	0.064	0.066	0.053	0.061	0.062	0.053	0.063	0.073	0.067	0.070	0.071	0.049	0.069	0.063	0.072	0.073	0.074	
Tues/Wed/Thurs	14	0.066	0.074	0.068	0.068	0.068	0.053	0.068	0.065	0.054	0.066	0.076	0.066	0.074	0.068	0.052	0.069	0.061	0.077	0.076	0.067	
Tues/Wed/Thurs	15	0.073	0.084	0.062	0.073	0.069	0.053	0.074	0.067	0.056	0.079	0.080	0.056	0.073	0.084	0.062	0.055	0.071	0.084	0.078	0.058	
Tues/Wed/Thurs	16	0.078	0.086	0.053	0.075	0.067	0.052	0.080	0.067	0.056	0.087	0.076	0.045	0.078	0.086	0.053	0.057	0.056	0.082	0.074	0.048	
Tues/Wed/Thurs	17	0.077	0.078	0.041	0.074	0.063	0.050	0.078	0.063	0.054	0.088	0.062	0.040	0.077	0.078	0.041	0.056	0.051	0.074	0.061	0.036	
Tues/Wed/Thurs	18	0.059	0.047	0.030	0.059	0.048	0.044	0.055	0.045	0.047	0.054	0.039	0.031	0.059	0.047	0.030	0.041	0.045	0.053	0.044	0.023	
Tues/Wed/Thurs	19	0.048	0.031	0.027	0.043	0.034	0.038	0.039	0.032	0.040	0.036	0.026	0.023	0.048	0.031	0.027	0.052	0.032	0.038	0.031	0.016	
Tues/Wed/Thurs	20	0.041	0.021	0.020	0.035	0.025	0.034	0.032	0.024	0.035	0.028	0.019	0.021	0.041	0.021	0.020	0.050	0.024	0.030	0.025	0.012	
Tues/Wed/Thurs	21	0.036	0.017	0.020	0.029	0.019	0.031	0.027	0.019	0.028	0.021	0.013	0.020	0.036	0.021	0.020	0.050	0.016	0.023	0.018	0.010	
Tues/Wed/Thurs	22	0.025	0.009	0.014	0.020	0.013	0.029	0.020	0.014	0.028	0.014	0.007	0.016	0.025	0.009	0.014	0.039	0.016	0.027	0.013	0.010	
Tues/Wed/Thurs	23	0.017	0.005	0.012	0.013	0.009	0.028	0.013	0.010	0.025	0.009	0.004	0.013	0.017	0.005	0.012	0.031	0.013	0.010	0.008	0.010	
Friday	0	0.009	0.004	0.008	0.007	0.014	0.032	0.007	0.019	0.030	0.007	0.003	0.011	0.009	0.004	0.008	0.023	0.009	0.005	0.009	0.019	
Friday	1	0.009	0.003	0.009	0.005	0.011	0.030	0.004	0.018	0.030	0.004	0.003	0.012	0.009	0.003	0.009	0.024	0.009	0.003	0.008	0.019	
Friday	2	0.009	0.003	0.011	0.004	0.011	0.030	0.003	0.017	0.029	0.004	0.003	0.015	0.009	0.003	0.011	0.024	0.009	0.002	0.008	0.019	
Friday	3	0.011	0.005	0.016	0.005	0.012	0.030	0.004	0.019	0.031	0.004	0.004	0.017	0.011	0.005	0.016	0.026	0.011	0.002	0.008	0.021	
Friday	4	0.013	0.009	0.022	0.008	0.016	0.033	0.009	0.023	0.034	0.006	0.024	0.024	0.013	0.009	0.022	0.028	0.017	0.005	0.013	0.024	
Friday	5	0.021	0.021	0.039	0.017	0.026	0.038	0.020	0.032	0.039	0.015	0.022	0.039	0.021	0.021	0.039	0.032	0.031	0.013	0.023	0.037	
Friday	6	0.033	0.041	0.054	0.033	0.040	0.045	0.037	0.044	0.046	0.035	0.045	0.055	0.047	0.060	0.078	0.040	0.040	0.026	0.035	0.049	
Friday	7	0.039	0.052	0.065	0.049	0.054	0.050	0.059	0.060	0.053	0.063	0.064	0.064	0.039	0.052	0.065	0.036	0.052	0.039	0.040	0.060	
Friday	8	0.044	0.059	0.074	0.051	0.057	0.052	0.057	0.059	0.053	0.058	0.072	0.074	0.044	0.059	0.074	0.039	0.058	0.043	0.049	0.068	
Friday	9	0.047	0.060	0.078	0.050	0.057	0.052	0.052	0.056	0.052	0.052	0.068	0.075	0.047	0.060	0.078	0.040	0.059	0.049	0.057	0.073	
Friday	10	0.048	0.067	0.075	0.054	0.061	0.054	0.053	0.057	0.052	0.055	0.071	0.074	0.048	0.067	0.075	0.043	0.063	0.058	0.063	0.078	
Friday	11	0.054	0.068	0.077	0.060	0.066	0.055	0.056	0.059	0.053	0.060	0.074	0.074	0.054	0.068	0.077	0.045	0.066	0.064	0.069	0.077	
Friday	12	0.060	0.072	0.079	0.063	0.067	0.055	0.059	0.061	0.053	0.063	0.072	0.069	0.060	0.072	0.079	0.046	0.063	0.066	0.071	0.076	
Friday	13	0.063	0.075	0.072	0.066	0.068	0.054	0.062	0.063	0.054	0.065	0.076	0.069	0.063	0.075	0.049	0.066	0.063	0.071	0.074	0.077	
Friday	14	0.068	0.078	0.067	0.070	0.070	0.054	0.068	0.066	0.055	0.069	0.078	0.063	0.068	0.078	0.067	0.051	0.067	0.076	0.077	0.070	
Friday	15	0.073	0.083	0.060	0.073	0.070	0.052	0.073	0.067	0.055	0.078	0.080	0.055	0.073	0.083	0.060	0.054	0.069	0.083	0.079	0.060	
Friday	16	0.076	0.082	0.049	0.074	0.067	0.050	0.077	0.067	0.053	0.085	0.075	0.057	0.076	0.082	0.049	0.056	0.067	0.083	0.077	0.050	
Friday	17	0.074	0.072	0.038	0.072	0.063	0.047	0.074	0.061	0.050	0.082	0.061	0.039	0.074	0.072	0.038	0.058	0.060	0.075	0.064	0.038	
Friday	18	0.060	0.050	0.026	0.063	0.051	0.042	0.060	0.047	0.043	0.059	0.041	0.029	0.060	0.050	0.026	0.057	0.051	0.062	0.051	0.025	
Friday	19	0.052	0.034	0.024	0.050	0.039	0.035	0.046	0.034	0.036	0.042	0.028	0.024	0.052	0.034	0.024	0.057	0.043	0.050	0.039	0.018	
Friday	20	0.043	0.022	0.017	0.041	0.029	0.030	0.038	0.026	0.030	0.032	0.021	0.021	0.043	0.022	0.017	0.053	0.033	0.041	0.030	0.013	
Friday	21	0.040	0.018	0.016	0.037	0.023	0.028	0.034	0.020	0.026	0.027	0.015	0.020	0.040	0.018	0.016	0.049	0.025	0.036	0.025	0.010	



Day of Week	Hour	Del Norte			El Dorado			Fresno			Glenn			Humboldt			Imperial			Inyo		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Friday	22	0.031	0.012	0.011	0.030	0.017	0.026	0.028	0.015	0.023	0.021	0.011	0.016	0.031	0.012	0.011	0.042	0.017	0.023	0.030	0.019	0.011
Friday	23	0.022	0.007	0.012	0.019	0.011	0.024	0.020	0.011	0.020	0.014	0.007	0.015	0.022	0.007	0.012	0.034	0.014	0.020	0.018	0.012	0.009
Saturday	0	0.012	0.008	0.014	0.013	0.019	0.038	0.010	0.028	0.041	0.012	0.007	0.021	0.012	0.008	0.014	0.025	0.018	0.036	0.010	0.015	0.027
Saturday	1	0.013	0.006	0.014	0.008	0.015	0.034	0.010	0.025	0.038	0.008	0.005	0.014	0.013	0.006	0.014	0.027	0.015	0.030	0.007	0.012	0.023
Saturday	2	0.013	0.004	0.011	0.006	0.014	0.032	0.008	0.024	0.037	0.006	0.004	0.020	0.013	0.004	0.011	0.027	0.012	0.024	0.005	0.011	0.022
Saturday	3	0.012	0.004	0.014	0.006	0.013	0.031	0.006	0.023	0.036	0.005	0.004	0.022	0.012	0.004	0.014	0.028	0.015	0.027	0.004	0.010	0.025
Saturday	4	0.014	0.008	0.020	0.007	0.014	0.032	0.009	0.024	0.037	0.006	0.008	0.024	0.014	0.008	0.020	0.031	0.019	0.030	0.005	0.013	0.028
Saturday	5	0.020	0.016	0.034	0.011	0.018	0.034	0.016	0.029	0.040	0.012	0.017	0.039	0.020	0.016	0.034	0.034	0.035	0.037	0.010	0.021	0.034
Saturday	6	0.025	0.025	0.043	0.019	0.026	0.039	0.026	0.036	0.045	0.021	0.028	0.049	0.025	0.025	0.043	0.035	0.038	0.043	0.017	0.028	0.039
Saturday	7	0.030	0.031	0.058	0.032	0.038	0.046	0.036	0.043	0.049	0.034	0.041	0.058	0.030	0.038	0.050	0.040	0.050	0.050	0.029	0.036	0.053
Saturday	8	0.036	0.041	0.070	0.045	0.051	0.052	0.045	0.050	0.052	0.045	0.057	0.067	0.036	0.041	0.070	0.040	0.057	0.055	0.044	0.045	0.060
Saturday	9	0.043	0.053	0.079	0.057	0.062	0.056	0.053	0.055	0.054	0.054	0.068	0.074	0.043	0.053	0.079	0.043	0.064	0.058	0.059	0.061	0.071
Saturday	10	0.052	0.069	0.082	0.067	0.071	0.060	0.060	0.061	0.056	0.063	0.080	0.073	0.052	0.069	0.082	0.044	0.066	0.064	0.073	0.074	0.078
Saturday	11	0.054	0.076	0.075	0.074	0.076	0.061	0.066	0.064	0.056	0.068	0.082	0.071	0.054	0.076	0.075	0.045	0.064	0.069	0.081	0.077	0.083
Saturday	12	0.061	0.080	0.070	0.075	0.075	0.060	0.069	0.065	0.056	0.074	0.083	0.068	0.061	0.080	0.070	0.046	0.063	0.066	0.078	0.077	0.075
Saturday	13	0.063	0.082	0.064	0.075	0.074	0.057	0.069	0.063	0.054	0.074	0.079	0.062	0.063	0.082	0.064	0.049	0.063	0.063	0.075	0.072	0.060
Saturday	14	0.065	0.081	0.062	0.074	0.071	0.055	0.070	0.063	0.053	0.074	0.076	0.057	0.065	0.081	0.062	0.051	0.062	0.059	0.075	0.068	0.055
Saturday	15	0.067	0.080	0.054	0.072	0.068	0.051	0.069	0.060	0.049	0.073	0.074	0.052	0.067	0.080	0.054	0.053	0.062	0.053	0.075	0.068	0.052
Saturday	16	0.071	0.081	0.051	0.070	0.064	0.048	0.067	0.057	0.046	0.073	0.067	0.045	0.071	0.081	0.051	0.053	0.057	0.047	0.072	0.070	0.047
Saturday	17	0.068	0.072	0.037	0.066	0.057	0.044	0.063	0.051	0.042	0.069	0.058	0.039	0.068	0.072	0.037	0.054	0.054	0.039	0.066	0.063	0.040
Saturday	18	0.062	0.053	0.032	0.056	0.047	0.038	0.056	0.044	0.036	0.058	0.047	0.034	0.062	0.053	0.032	0.055	0.048	0.034	0.058	0.052	0.031
Saturday	19	0.059	0.040	0.029	0.046	0.037	0.033	0.047	0.036	0.031	0.046	0.036	0.029	0.059	0.040	0.029	0.049	0.030	0.030	0.047	0.041	0.026
Saturday	20	0.051	0.032	0.021	0.040	0.030	0.028	0.041	0.031	0.027	0.040	0.028	0.024	0.051	0.032	0.021	0.049	0.032	0.026	0.038	0.031	0.020
Saturday	21	0.047	0.026	0.023	0.035	0.025	0.025	0.038	0.027	0.023	0.036	0.022	0.023	0.047	0.026	0.023	0.045	0.025	0.023	0.031	0.025	0.016
Saturday	22	0.037	0.019	0.020	0.028	0.019	0.023	0.034	0.024	0.019	0.029	0.016	0.017	0.037	0.019	0.020	0.030	0.020	0.020	0.025	0.020	0.018
Saturday	23	0.028	0.014	0.021	0.020	0.014	0.021	0.024	0.019	0.019	0.020	0.011	0.017	0.028	0.014	0.021	0.036	0.018	0.016	0.016	0.013	0.018
Holiday	0	0.010	0.004	0.009	0.010	0.016	0.028	0.013	0.023	0.029	0.010	0.004	0.012	0.010	0.004	0.009	0.027	0.013	0.019	0.008	0.011	0.020
Holiday	1	0.014	0.004	0.008	0.006	0.013	0.027	0.007	0.022	0.027	0.006	0.004	0.011	0.014	0.004	0.008	0.028	0.008	0.018	0.005	0.009	0.018
Holiday	2	0.010	0.003	0.014	0.004	0.012	0.026	0.005	0.022	0.027	0.004	0.003	0.014	0.010	0.003	0.014	0.026	0.008	0.018	0.003	0.010	0.018
Holiday	3	0.014	0.005	0.012	0.005	0.013	0.027	0.004	0.021	0.028	0.004	0.005	0.015	0.014	0.005	0.012	0.027	0.010	0.018	0.004	0.010	0.021
Holiday	4	0.014	0.006	0.017	0.008	0.016	0.029	0.008	0.024	0.030	0.007	0.009	0.024	0.014	0.006	0.017	0.030	0.016	0.022	0.005	0.012	0.020
Holiday	5	0.019	0.018	0.028	0.014	0.023	0.032	0.016	0.031	0.034	0.014	0.020	0.037	0.019	0.018	0.028	0.030	0.026	0.029	0.009	0.018	0.031
Holiday	6	0.028	0.034	0.042	0.025	0.033	0.036	0.028	0.039	0.038	0.030	0.036	0.047	0.028	0.034	0.042	0.032	0.032	0.031	0.018	0.023	0.038
Holiday	7	0.039	0.045	0.052	0.036	0.044	0.042	0.040	0.046	0.041	0.044	0.052	0.061	0.039	0.045	0.052	0.042	0.042	0.037	0.029	0.031	0.043
Holiday	8	0.041	0.051	0.059	0.046	0.053	0.048	0.045	0.049	0.043	0.052	0.066	0.075	0.041	0.051	0.059	0.040	0.055	0.044	0.041	0.044	0.056
Holiday	9	0.044	0.057	0.066	0.054	0.059	0.050	0.049	0.052	0.047	0.053	0.071	0.081	0.044	0.057	0.066	0.042	0.061	0.054	0.058	0.057	0.075
Holiday	10	0.050	0.069	0.075	0.065	0.069	0.053	0.057	0.059	0.049	0.059	0.076	0.081	0.050	0.069	0.075	0.045	0.067	0.060	0.076	0.083	0.087
Holiday	11	0.056	0.072	0.077	0.074	0.074	0.057	0.065	0.063	0.051	0.066	0.076	0.071	0.056	0.072	0.077	0.047	0.070	0.068	0.084	0.086	0.088
Holiday	12	0.058	0.080	0.078	0.077	0.074	0.056	0.070	0.067	0.054	0.071	0.078	0.074	0.058	0.080	0.078	0.046	0.069	0.070	0.085	0.087	0.089
Holiday	13	0.063	0.077	0.069	0.076	0.074	0.058	0.072	0.067	0.056	0.071	0.076	0.065	0.063	0.077	0.069	0.053	0.080	0.070	0.083	0.081	0.078
Holiday	14	0.068	0.083	0.067	0.075	0.073	0.056	0.074	0.066	0.055	0.070	0.078	0.060	0.068	0.083	0.067	0.051	0.075	0.068	0.080	0.074	0.068
Holiday	15	0.071	0.082	0.064	0.074	0.070	0.055	0.076	0.067	0.056	0.075	0.075	0.053	0.071	0.082	0.064	0.054	0.067	0.062	0.078	0.074	0.060
Holiday	16	0.075	0.083	0.061	0.072	0.066	0.054	0.076	0.064	0.055	0.079	0.070	0.044	0.075	0.083	0.061	0.056	0.066	0.057	0.078	0.072	0.049
Holiday	17	0.072	0.076	0.044	0.068	0.059	0.051	0.072	0.058	0.052	0.074	0.064	0.041	0.072	0.076	0.044	0.056	0.061	0.054	0.080	0.074	0.068
Holiday	18	0.054	0.048	0.040	0.057	0.049	0.045	0.058	0.046	0.049	0.058	0.044	0.034	0.054	0.048	0.040	0.052	0.047	0.045	0.057	0.049	0.033
Holiday	19	0.056	0.036	0.029	0.047	0.036	0.041	0.047	0.035	0.043	0.047	0.033	0.026	0.056	0.036	0.029	0.053	0.039	0.040	0.043	0.040	0.022
Holiday	20	0.049	0.025	0.029	0.037	0.029	0.037	0.039	0.028	0.040	0.038	0.025	0.025	0.039	0.028	0.040	0.038	0.025	0.025	0.033	0.036	0.013
Holiday	21	0.040	0.019	0.023	0.030	0.020	0.033	0.032	0.022	0.036	0.030	0.018	0.021	0.040	0.019	0.023	0.046	0.022	0.030	0.024	0.018	0.011
Holiday	22	0.029	0.012	0.018	0.023	0.015	0.031	0.026	0.017	0.032	0.024	0.011	0.017	0.029	0.012	0.018	0.042	0.020	0.027	0.017	0.012	0.009
Holiday	23	0.025	0.010	0.019	0.015	0.015	0.029	0.018	0.013	0.029	0.014	0.007	0.014	0.025	0.010	0.019	0.032	0.019	0.025	0.010	0.008	0.010

Day of Week	Hour	Kern			Kings			Lake			Lassen			Los Angeles			Madera			Marin		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Sunday	0	0.014	0.028	0.041	0.016	0.031	0.042	0.013	0.011	0.008	0.020	0.007	0.015	0.025	0.043	0.051	0.014	0.037	0.044	0.019	0.038	0.053
Sunday	1	0.010	0.024	0.038	0.010	0.025	0.038	0.013	0.008	0.010	0.020	0.005	0.014	0.018	0.033	0.044	0.008	0.032	0.040	0.012	0.034	0.047
Sunday	2	0.007	0.022	0.034	0.007	0.026	0.036	0.012	0.006	0.008	0.020	0.003	0.012	0.014	0.028	0.040	0.005	0.028	0.037	0.008	0.031	0.043
Sunday	3	0.006	0.020	0.033	0.005	0.022	0.031	0.014	0.005	0.007	0.021	0.004	0.011	0.009	0.022	0.035	0.004	0.026	0.035	0.006	0.030	0.040
Sunday	4	0.007	0.021	0.033	0.004	0.020	0.031	0.014	0.004	0.011	0.024	0.007	0.011	0.008	0.021	0.034	0.004	0.025	0.034	0.006	0.029	0.038
Sunday	5	0.012	0.024	0.033	0.008	0.023	0.031	0.017	0.009	0.019	0.028	0.012	0.015	0.012	0.024	0.035	0.009	0.027	0.034	0.010	0.031	0.038
Sunday	6	0.016	0.027	0.034	0.018	0.029	0.036	0.021	0.014	0.028	0.030	0.017	0.026	0.018	0.029	0.037	0.016	0.030	0.036	0.016	0.033	0.039
Sunday	7	0.024	0.032	0.035	0.023	0.030	0.035	0.026	0.020	0.036	0.034	0.032	0.037	0.025	0.034	0.039	0.022	0.033	0.036	0.023	0.036	0.040
Sunday	8	0.032	0.039	0.038	0.034	0.040	0.040	0.031	0.032	0.043	0.037	0.045	0.053	0.035	0.040	0.042	0.033	0.039	0.040	0.033	0.040	0.042
Sunday	9	0.042	0.045	0.040	0.048	0.049	0.046	0.040	0.050	0.054	0.044	0.064	0.064	0.047	0.050	0.045	0.046	0.044	0.048	0.048	0.046	0.044
Sunday	10	0.051	0.051	0.042	0.059	0.057	0.049	0.047	0.064	0.067	0.046	0.076	0.076	0.057	0.056	0.047	0.056	0.052	0.046	0.062	0.051	0.045
Sunday	11	0.059	0.056	0.045	0.071	0.064	0.052	0.055	0.079	0.062	0.050	0.088	0.079	0.062	0.059	0.047	0.065	0.057	0.048	0.067	0.053	0.046
Sunday	12	0.066	0.060	0.046	0.084	0.077	0.057	0.061	0.087	0.065	0.053	0.088	0.075	0.065	0.060	0.046	0.071	0.059	0.049	0.070	0.054	0.046
Sunday	13	0.071	0.063	0.047	0.083	0.077	0.056	0.065	0.092	0.064	0.054	0.082	0.069	0.068	0.058	0.044	0.076	0.059	0.048	0.073	0.055	0.050
Sunday	14	0.075	0.065	0.047	0.080	0.072	0.055	0.067	0.087	0.065	0.059	0.075	0.067	0.068	0.058	0.044	0.076	0.059	0.048	0.073	0.055	0.047
Sunday	15	0.078	0.064	0.048	0.076	0.065	0.052	0.072	0.086	0.067	0.060	0.076	0.064	0.067	0.055	0.043	0.076	0.058	0.047	0.072	0.052	0.039
Sunday	16	0.077	0.063	0.048	0.074	0.062	0.050	0.077	0.086	0.072	0.063	0.074	0.058	0.065	0.052	0.042	0.077	0.058	0.047	0.072	0.052	0.039
Sunday	17	0.074	0.060	0.047	0.068	0.056	0.046	0.070	0.075	0.058	0.063	0.063	0.058	0.063	0.049	0.040	0.074	0.055	0.046	0.070	0.050	0.038
Sunday	18	0.069	0.055	0.046	0.059	0.044	0.042	0.067	0.059	0.054	0.061	0.053	0.050	0.059	0.045	0.040	0.068	0.048	0.043	0.063	0.047	0.036
Sunday	19	0.061	0.049	0.046	0.050	0.037	0.037	0.062	0.045	0.050	0.059	0.051	0.041	0.056	0.042	0.039	0.060	0.043	0.041	0.056	0.044	0.035
Sunday	20	0.053	0.042	0.045	0.043	0.032	0.037	0.054	0.035	0.047	0.051	0.034	0.036	0.052	0.040	0.040	0.052	0.039	0.040	0.051	0.041	0.036
Sunday	21	0.042	0.035	0.044	0.036	0.028	0.035	0.045	0.024	0.039	0.044	0.025	0.031	0.047	0.038	0.041	0.042	0.034	0.039	0.042	0.038	0.037
Sunday	22	0.032	0.030	0.045	0.028	0.022	0.034	0.033	0.015	0.033	0.035	0.016	0.024	0.036	0.034	0.042	0.038	0.038	0.030	0.030	0.034	0.039
Sunday	23	0.021	0.025	0.046	0.015	0.015	0.033	0.022	0.009	0.032	0.024	0.009	0.018	0.024	0.029	0.042	0.018	0.023	0.037	0.019	0.027	0.043
Monday	0	0.013	0.022	0.025	0.005	0.013	0.019	0.010	0.003	0.007	0.020	0.005	0.012	0.012	0.018	0.025	0.007	0.021	0.024	0.007	0.023	0.029
Monday	1	0.009	0.019	0.024	0.002	0.012	0.019	0.009	0.002	0.007	0.021	0.003	0.010	0.006	0.015	0.023	0.003	0.020	0.024	0.003	0.022	0.028
Monday	2	0.008	0.019	0.024	0.001	0.014	0.020	0.010	0.003	0.010	0.022	0.003	0.012	0.006	0.015	0.023	0.002	0.020	0.024	0.002	0.022	0.029
Monday	3	0.011	0.022	0.026	0.001	0.012	0.019	0.012	0.006	0.012	0.023	0.004	0.012	0.007	0.017	0.024	0.004	0.023	0.026	0.003	0.023	0.030
Monday	4	0.021	0.029	0.028	0.003	0.015	0.021	0.014	0.009	0.013	0.026	0.011	0.013	0.016	0.024	0.030	0.012	0.028	0.029	0.012	0.028	0.035
Monday	5	0.040	0.041	0.033	0.012	0.021	0.027	0.022	0.022	0.026	0.037	0.047	0.021	0.038	0.042	0.038	0.029	0.036	0.036	0.033	0.041	0.042
Monday	6	0.047	0.046	0.034	0.034	0.040	0.038	0.037	0.047	0.044	0.038	0.049	0.030	0.054	0.056	0.044	0.050	0.051	0.044	0.054	0.051	0.048
Monday	7	0.056	0.054	0.038	0.070	0.071	0.056	0.045	0.058	0.058	0.041	0.051	0.035	0.061	0.062	0.049	0.072	0.063	0.051	0.066	0.058	0.053
Monday	8	0.050	0.052	0.038	0.073	0.071	0.056	0.047	0.062	0.067	0.043	0.058	0.047	0.059	0.061	0.049	0.063	0.059	0.049	0.062	0.060	0.055
Monday	9	0.049	0.052	0.039	0.061	0.062	0.053	0.050	0.065	0.078	0.045	0.073	0.058	0.054	0.058	0.049	0.058	0.056	0.049	0.055	0.056	0.054
Monday	10	0.052	0.053	0.042	0.059	0.062	0.054	0.051	0.065	0.080	0.047	0.076	0.068	0.052	0.057	0.050	0.057	0.057	0.051	0.052	0.054	0.053
Monday	11	0.057	0.056	0.044	0.059	0.063	0.056	0.056	0.067	0.083	0.052	0.073	0.077	0.054	0.058	0.052	0.060	0.062	0.055	0.053	0.055	0.054
Monday	12	0.061	0.059	0.046	0.062	0.064	0.056	0.058	0.069	0.081	0.053	0.068	0.073	0.054	0.058	0.052	0.060	0.062	0.055	0.054	0.056	0.054
Monday	13	0.064	0.060	0.049	0.064	0.067	0.058	0.063	0.074	0.076	0.056	0.065	0.066	0.055	0.058	0.052	0.061	0.061	0.054	0.056	0.056	0.054
Monday	14	0.068	0.063	0.052	0.073	0.071	0.064	0.067	0.076	0.074	0.058	0.072	0.066	0.059	0.060	0.052	0.066	0.062	0.057	0.063	0.059	0.056
Monday	15	0.074	0.067	0.057	0.078	0.072	0.064	0.073	0.087	0.062	0.059	0.079	0.061	0.062	0.060	0.052	0.071	0.064	0.058	0.069	0.063	0.058
Monday	16	0.073	0.065	0.058	0.086	0.073	0.062	0.076	0.084	0.053	0.061	0.069	0.053	0.063	0.058	0.051	0.062	0.057	0.072	0.072	0.060	0.052
Monday	17	0.067	0.058	0.057	0.087	0.070	0.062	0.075	0.075	0.040	0.059	0.067	0.054	0.064	0.055	0.050	0.074	0.058	0.055	0.073	0.056	0.047
Monday	18	0.050	0.044	0.053	0.056	0.046	0.053	0.057	0.047	0.032	0.056	0.043	0.048	0.059	0.047	0.047	0.052	0.041	0.047	0.061	0.045	0.039
Monday	19	0.037	0.034	0.049	0.037	0.028	0.038	0.050	0.031	0.029	0.050	0.030	0.044	0.049	0.036	0.042	0.037	0.030	0.039	0.045	0.033	0.031
Monday	20	0.032	0.028	0.048	0.029	0.021	0.033	0.043	0.020	0.021	0.043	0.021	0.036	0.039	0.028	0.038	0.030	0.022	0.034	0.035	0.026	0.026
Monday	21	0.026	0.023	0.048	0.023	0.015	0.029	0.035	0.015	0.020	0.040	0.016	0.037	0.034	0.023	0.037	0.025	0.017	0.031	0.031	0.022	0.024
Monday	22	0.021	0.018	0.044	0.016	0.010	0.024	0.025	0.009	0.014	0.030	0.009	0.035	0.027	0.020	0.036	0.019	0.014	0.027	0.023	0.017	0.023
Monday	23	0.014	0.015	0.042	0.009	0.007	0.021	0.016	0.005	0.013	0.022	0.006	0.030	0.017	0.016	0.035	0.012	0.011	0.024	0.014	0.014	0.025

Day of Week	Hour	Kern		Kings		Lake		Lassen		Los Angeles		Madera		Marin		
		LD	HH	LD	HH	LD	HH	LD	HH	LD	HH	LD	HH	LD	HH	
Tues/Wed/Thurs	0	0.010	0.021	0.013	0.022	0.010	0.004	0.004	0.004	0.011	0.019	0.005	0.020	0.006	0.022	0.031
Tues/Wed/Thurs	1	0.006	0.019	0.031	0.002	0.009	0.003	0.008	0.008	0.006	0.016	0.001	0.029	0.003	0.021	0.030
Tues/Wed/Thurs	2	0.006	0.019	0.031	0.000	0.010	0.002	0.012	0.012	0.005	0.016	0.001	0.019	0.002	0.021	0.030
Tues/Wed/Thurs	3	0.009	0.022	0.031	0.000	0.011	0.005	0.014	0.025	0.007	0.017	0.002	0.028	0.003	0.023	0.031
Tues/Wed/Thurs	4	0.019	0.029	0.034	0.003	0.015	0.010	0.021	0.028	0.015	0.025	0.033	0.010	0.011	0.028	0.036
Tues/Wed/Thurs	5	0.039	0.041	0.037	0.012	0.024	0.024	0.035	0.039	0.037	0.042	0.041	0.027	0.034	0.040	0.044
Tues/Wed/Thurs	6	0.048	0.046	0.039	0.035	0.048	0.048	0.048	0.041	0.054	0.056	0.047	0.050	0.056	0.052	0.049
Tues/Wed/Thurs	7	0.058	0.053	0.042	0.070	0.045	0.059	0.065	0.041	0.061	0.062	0.051	0.074	0.068	0.059	0.054
Tues/Wed/Thurs	8	0.052	0.052	0.042	0.073	0.047	0.063	0.069	0.044	0.059	0.062	0.051	0.065	0.063	0.060	0.056
Tues/Wed/Thurs	9	0.049	0.050	0.041	0.060	0.050	0.064	0.074	0.067	0.054	0.058	0.050	0.057	0.055	0.055	0.053
Tues/Wed/Thurs	10	0.050	0.051	0.042	0.057	0.060	0.054	0.067	0.067	0.052	0.057	0.051	0.055	0.051	0.053	0.052
Tues/Wed/Thurs	11	0.054	0.054	0.044	0.058	0.055	0.065	0.076	0.049	0.052	0.057	0.051	0.056	0.050	0.054	0.052
Tues/Wed/Thurs	12	0.059	0.056	0.046	0.060	0.057	0.068	0.076	0.051	0.053	0.057	0.051	0.057	0.052	0.055	0.053
Tues/Wed/Thurs	13	0.062	0.058	0.047	0.061	0.061	0.070	0.071	0.064	0.055	0.058	0.050	0.059	0.054	0.056	0.054
Tues/Wed/Thurs	14	0.068	0.062	0.050	0.071	0.066	0.074	0.068	0.062	0.059	0.059	0.050	0.065	0.062	0.059	0.054
Tues/Wed/Thurs	15	0.075	0.067	0.053	0.077	0.073	0.084	0.062	0.058	0.060	0.058	0.049	0.072	0.067	0.063	0.056
Tues/Wed/Thurs	16	0.075	0.066	0.054	0.086	0.078	0.086	0.053	0.070	0.062	0.056	0.048	0.078	0.070	0.060	0.051
Tues/Wed/Thurs	17	0.070	0.060	0.053	0.087	0.077	0.078	0.041	0.060	0.062	0.053	0.046	0.079	0.071	0.057	0.046
Tues/Wed/Thurs	18	0.052	0.046	0.048	0.059	0.051	0.059	0.047	0.043	0.058	0.046	0.043	0.055	0.062	0.047	0.039
Tues/Wed/Thurs	19	0.039	0.036	0.044	0.039	0.032	0.038	0.027	0.049	0.051	0.036	0.039	0.040	0.048	0.035	0.031
Tues/Wed/Thurs	20	0.033	0.030	0.042	0.032	0.041	0.021	0.020	0.044	0.042	0.028	0.036	0.033	0.038	0.027	0.026
Tues/Wed/Thurs	21	0.029	0.025	0.041	0.026	0.036	0.017	0.018	0.038	0.037	0.024	0.028	0.028	0.033	0.022	0.024
Tues/Wed/Thurs	22	0.023	0.020	0.039	0.018	0.025	0.009	0.014	0.029	0.030	0.020	0.033	0.021	0.024	0.017	0.022
Tues/Wed/Thurs	23	0.015	0.017	0.038	0.010	0.017	0.005	0.012	0.022	0.019	0.016	0.032	0.013	0.015	0.013	0.024
Friday	0	0.009	0.021	0.035	0.006	0.014	0.024	0.008	0.004	0.012	0.021	0.032	0.005	0.020	0.022	0.033
Friday	1	0.007	0.019	0.034	0.002	0.012	0.024	0.009	0.004	0.005	0.017	0.030	0.002	0.004	0.021	0.031
Friday	2	0.006	0.019	0.034	0.001	0.011	0.022	0.009	0.003	0.003	0.017	0.030	0.001	0.003	0.022	0.032
Friday	3	0.018	0.021	0.035	0.001	0.013	0.024	0.011	0.005	0.007	0.018	0.031	0.003	0.004	0.023	0.033
Friday	4	0.005	0.027	0.037	0.002	0.015	0.025	0.013	0.009	0.014	0.025	0.035	0.008	0.010	0.028	0.036
Friday	5	0.031	0.037	0.040	0.011	0.021	0.021	0.039	0.033	0.033	0.040	0.044	0.022	0.030	0.039	0.044
Friday	6	0.039	0.043	0.043	0.031	0.039	0.043	0.054	0.035	0.034	0.040	0.044	0.039	0.050	0.049	0.050
Friday	7	0.048	0.050	0.045	0.063	0.064	0.057	0.039	0.040	0.046	0.060	0.053	0.059	0.063	0.057	0.055
Friday	8	0.045	0.050	0.045	0.067	0.069	0.059	0.044	0.044	0.061	0.060	0.054	0.054	0.059	0.057	0.056
Friday	9	0.045	0.049	0.046	0.057	0.062	0.057	0.047	0.047	0.068	0.060	0.052	0.058	0.053	0.054	0.054
Friday	10	0.049	0.053	0.047	0.057	0.063	0.056	0.048	0.046	0.068	0.071	0.052	0.058	0.051	0.053	0.054
Friday	11	0.054	0.055	0.048	0.059	0.065	0.058	0.054	0.049	0.075	0.077	0.053	0.059	0.053	0.055	0.054
Friday	12	0.058	0.057	0.049	0.061	0.064	0.058	0.060	0.072	0.079	0.070	0.054	0.059	0.056	0.057	0.055
Friday	13	0.063	0.060	0.050	0.062	0.066	0.058	0.063	0.075	0.072	0.065	0.056	0.059	0.058	0.058	0.056
Friday	14	0.068	0.063	0.051	0.070	0.069	0.058	0.068	0.078	0.067	0.060	0.057	0.059	0.064	0.059	0.056
Friday	15	0.072	0.067	0.053	0.073	0.069	0.060	0.073	0.083	0.060	0.055	0.058	0.071	0.066	0.062	0.056
Friday	16	0.073	0.064	0.052	0.079	0.073	0.060	0.076	0.082	0.049	0.054	0.059	0.062	0.067	0.059	0.050
Friday	17	0.070	0.059	0.050	0.079	0.065	0.055	0.074	0.072	0.038	0.046	0.059	0.051	0.067	0.055	0.046
Friday	18	0.060	0.048	0.044	0.061	0.050	0.047	0.060	0.050	0.043	0.043	0.057	0.046	0.060	0.047	0.039
Friday	19	0.049	0.039	0.039	0.045	0.034	0.036	0.052	0.034	0.024	0.036	0.037	0.035	0.049	0.036	0.030
Friday	20	0.042	0.032	0.035	0.036	0.023	0.028	0.043	0.022	0.017	0.046	0.045	0.029	0.040	0.029	0.023
Friday	21	0.037	0.027	0.032	0.031	0.017	0.024	0.040	0.018	0.016	0.041	0.040	0.024	0.035	0.023	0.020
Friday	22	0.031	0.023	0.029	0.031	0.019	0.031	0.031	0.018	0.026	0.036	0.036	0.021	0.030	0.030	0.019
Friday	23	0.021	0.018	0.027	0.017	0.008	0.016	0.022	0.007	0.012	0.024	0.027	0.017	0.022	0.015	0.020
Saturday	0	0.016	0.028	0.043	0.013	0.012	0.035	0.012	0.008	0.014	0.025	0.020	0.031	0.015	0.030	0.044
Saturday	1	0.011	0.023	0.041	0.008	0.013	0.006	0.014	0.009	0.007	0.015	0.020	0.008	0.009	0.027	0.040
Saturday	2	0.009	0.022	0.040	0.005	0.017	0.031	0.013	0.004	0.011	0.025	0.011	0.023	0.006	0.026	0.039
Saturday	3	0.009	0.021	0.040	0.003	0.012	0.004	0.014	0.026	0.004	0.017	0.008	0.020	0.005	0.025	0.037

Day of Week	Hour	Kern		Kings		Lake		Lassen		Los Angeles		Madera		Marin	
		LD	HH	LD	HH	LD	HH	LD	HH	LD	HH	LD	HH	LD	HH
Saturday	4	0.014	0.025	0.041	0.031	0.014	0.008	0.020	0.007	0.022	0.038	0.008	0.027	0.006	0.027
Saturday	5	0.027	0.034	0.044	0.033	0.020	0.016	0.034	0.022	0.023	0.042	0.017	0.032	0.013	0.030
Saturday	6	0.034	0.038	0.045	0.041	0.025	0.025	0.043	0.035	0.033	0.046	0.026	0.039	0.023	0.035
Saturday	7	0.042	0.045	0.047	0.048	0.030	0.031	0.058	0.039	0.041	0.037	0.036	0.045	0.034	0.041
Saturday	8	0.050	0.052	0.050	0.049	0.036	0.041	0.070	0.044	0.057	0.052	0.047	0.052	0.046	0.047
Saturday	9	0.056	0.056	0.052	0.053	0.043	0.053	0.079	0.047	0.065	0.053	0.057	0.057	0.055	0.051
Saturday	10	0.060	0.057	0.053	0.061	0.052	0.069	0.082	0.050	0.080	0.060	0.062	0.062	0.061	0.054
Saturday	11	0.063	0.059	0.053	0.067	0.054	0.076	0.075	0.050	0.078	0.060	0.067	0.063	0.065	0.056
Saturday	12	0.065	0.061	0.052	0.071	0.061	0.080	0.070	0.053	0.075	0.062	0.068	0.062	0.066	0.058
Saturday	13	0.066	0.061	0.050	0.060	0.063	0.082	0.064	0.055	0.070	0.064	0.062	0.062	0.067	0.059
Saturday	14	0.067	0.060	0.049	0.070	0.065	0.081	0.062	0.068	0.068	0.048	0.068	0.059	0.067	0.058
Saturday	15	0.067	0.060	0.048	0.070	0.067	0.080	0.054	0.063	0.059	0.045	0.068	0.056	0.068	0.057
Saturday	16	0.064	0.056	0.044	0.070	0.071	0.081	0.051	0.057	0.064	0.055	0.062	0.054	0.068	0.056
Saturday	17	0.058	0.052	0.041	0.066	0.068	0.072	0.037	0.055	0.064	0.060	0.064	0.050	0.067	0.054
Saturday	18	0.051	0.046	0.036	0.059	0.062	0.053	0.032	0.052	0.049	0.044	0.057	0.042	0.060	0.048
Saturday	19	0.044	0.037	0.032	0.049	0.059	0.040	0.029	0.048	0.039	0.031	0.037	0.029	0.049	0.041
Saturday	20	0.039	0.033	0.028	0.043	0.051	0.032	0.021	0.046	0.034	0.030	0.046	0.043	0.043	0.036
Saturday	21	0.035	0.029	0.026	0.040	0.047	0.026	0.023	0.039	0.026	0.043	0.030	0.027	0.041	0.033
Saturday	22	0.030	0.024	0.024	0.037	0.037	0.019	0.020	0.031	0.020	0.042	0.029	0.024	0.037	0.029
Saturday	23	0.023	0.020	0.020	0.024	0.028	0.014	0.021	0.023	0.010	0.017	0.033	0.026	0.028	0.024
Holiday	0	0.015	0.023	0.028	0.011	0.010	0.004	0.009	0.007	0.015	0.017	0.024	0.031	0.013	0.027
Holiday	1	0.009	0.021	0.028	0.006	0.014	0.004	0.008	0.003	0.012	0.011	0.020	0.028	0.007	0.026
Holiday	2	0.007	0.020	0.028	0.002	0.010	0.003	0.014	0.003	0.011	0.009	0.019	0.027	0.004	0.025
Holiday	3	0.008	0.021	0.028	0.001	0.014	0.005	0.012	0.002	0.016	0.007	0.019	0.028	0.003	0.025
Holiday	4	0.013	0.024	0.028	0.002	0.014	0.006	0.017	0.004	0.015	0.012	0.023	0.030	0.007	0.029
Holiday	5	0.027	0.032	0.032	0.010	0.019	0.018	0.028	0.031	0.020	0.021	0.024	0.033	0.017	0.034
Holiday	6	0.033	0.037	0.033	0.026	0.028	0.034	0.042	0.033	0.025	0.028	0.034	0.041	0.029	0.040
Holiday	7	0.039	0.043	0.036	0.043	0.039	0.045	0.052	0.038	0.036	0.044	0.042	0.047	0.038	0.045
Holiday	8	0.043	0.047	0.037	0.050	0.041	0.051	0.059	0.044	0.054	0.043	0.045	0.050	0.045	0.050
Holiday	9	0.050	0.050	0.040	0.051	0.044	0.057	0.066	0.046	0.071	0.064	0.048	0.053	0.049	0.053
Holiday	10	0.055	0.055	0.042	0.060	0.050	0.069	0.075	0.051	0.088	0.073	0.054	0.058	0.056	0.056
Holiday	11	0.064	0.060	0.047	0.067	0.056	0.072	0.077	0.053	0.082	0.075	0.058	0.061	0.062	0.059
Holiday	12	0.068	0.061	0.050	0.073	0.058	0.080	0.078	0.055	0.082	0.072	0.061	0.063	0.067	0.061
Holiday	13	0.071	0.066	0.051	0.075	0.063	0.077	0.069	0.054	0.078	0.063	0.063	0.064	0.070	0.062
Holiday	14	0.073	0.064	0.052	0.076	0.068	0.083	0.067	0.060	0.077	0.067	0.064	0.064	0.073	0.062
Holiday	15	0.075	0.067	0.055	0.072	0.071	0.082	0.064	0.054	0.081	0.062	0.065	0.061	0.071	0.061
Holiday	16	0.072	0.064	0.055	0.075	0.075	0.083	0.061	0.062	0.077	0.063	0.064	0.057	0.070	0.057
Holiday	17	0.066	0.059	0.054	0.071	0.072	0.076	0.044	0.061	0.066	0.050	0.063	0.053	0.067	0.053
Holiday	18	0.056	0.046	0.049	0.059	0.054	0.048	0.040	0.057	0.043	0.042	0.058	0.046	0.059	0.045
Holiday	19	0.047	0.042	0.050	0.047	0.056	0.036	0.029	0.052	0.035	0.041	0.052	0.038	0.050	0.036
Holiday	20	0.039	0.033	0.046	0.040	0.049	0.025	0.029	0.043	0.022	0.034	0.047	0.032	0.046	0.031
Holiday	21	0.031	0.027	0.046	0.034	0.040	0.019	0.023	0.041	0.024	0.036	0.042	0.028	0.041	0.026
Holiday	22	0.025	0.021	0.043	0.030	0.029	0.012	0.018	0.031	0.011	0.026	0.037	0.025	0.033	0.021
Holiday	23	0.016	0.018	0.041	0.018	0.025	0.010	0.019	0.022	0.009	0.026	0.025	0.020	0.021	0.017

Day of Week	Hour	Mariposa			Mendocino			Merced			Modoc			Mono			Monterey			Napa		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Sunday	0	0.010	0.014	0.032	0.013	0.011	0.008	0.014	0.025	0.037	0.019	0.009	0.017	0.010	0.014	0.032	0.019	0.010	0.010	0.017	0.035	0.054
Sunday	1	0.007	0.011	0.024	0.013	0.008	0.010	0.009	0.019	0.032	0.021	0.007	0.014	0.007	0.011	0.024	0.020	0.008	0.023	0.011	0.030	0.047
Sunday	2	0.005	0.011	0.022	0.012	0.006	0.008	0.007	0.016	0.029	0.022	0.006	0.013	0.005	0.011	0.022	0.020	0.007	0.021	0.007	0.028	0.044
Sunday	3	0.004	0.010	0.021	0.014	0.005	0.007	0.005	0.015	0.028	0.022	0.005	0.013	0.004	0.010	0.021	0.020	0.007	0.019	0.006	0.026	0.043
Sunday	4	0.004	0.010	0.020	0.014	0.004	0.011	0.006	0.016	0.028	0.023	0.006	0.013	0.004	0.010	0.020	0.024	0.012	0.019	0.006	0.025	0.038
Sunday	5	0.007	0.013	0.021	0.017	0.009	0.019	0.010	0.019	0.029	0.025	0.008	0.016	0.007	0.013	0.021	0.026	0.017	0.021	0.009	0.027	0.038
Sunday	6	0.012	0.019	0.026	0.021	0.014	0.028	0.015	0.023	0.031	0.028	0.014	0.024	0.012	0.019	0.026	0.029	0.024	0.024	0.014	0.030	0.038
Sunday	7	0.019	0.023	0.029	0.026	0.020	0.036	0.021	0.029	0.035	0.030	0.022	0.034	0.019	0.023	0.029	0.031	0.030	0.034	0.020	0.033	0.039
Sunday	8	0.032	0.035	0.038	0.031	0.032	0.043	0.031	0.038	0.040	0.033	0.036	0.048	0.032	0.035	0.038	0.035	0.038	0.040	0.031	0.038	0.042
Sunday	9	0.051	0.051	0.053	0.040	0.050	0.054	0.043	0.050	0.047	0.036	0.052	0.062	0.051	0.051	0.053	0.038	0.049	0.047	0.047	0.046	0.046
Sunday	10	0.067	0.067	0.071	0.047	0.064	0.067	0.055	0.060	0.051	0.040	0.071	0.075	0.067	0.067	0.071	0.041	0.057	0.060	0.060	0.054	0.046
Sunday	11	0.080	0.081	0.085	0.055	0.079	0.062	0.063	0.065	0.054	0.044	0.082	0.086	0.080	0.081	0.085	0.047	0.068	0.066	0.066	0.056	0.047
Sunday	12	0.083	0.081	0.076	0.061	0.087	0.065	0.070	0.070	0.055	0.049	0.089	0.088	0.083	0.081	0.076	0.051	0.074	0.063	0.067	0.056	0.045
Sunday	13	0.085	0.082	0.074	0.065	0.092	0.064	0.075	0.071	0.056	0.054	0.090	0.080	0.085	0.082	0.074	0.053	0.073	0.070	0.070	0.056	0.042
Sunday	14	0.085	0.083	0.069	0.067	0.087	0.065	0.077	0.069	0.055	0.058	0.089	0.072	0.085	0.083	0.069	0.059	0.078	0.071	0.057	0.057	0.038
Sunday	15	0.084	0.081	0.066	0.072	0.086	0.067	0.078	0.070	0.053	0.063	0.087	0.069	0.084	0.081	0.066	0.061	0.078	0.066	0.071	0.052	0.037
Sunday	16	0.082	0.079	0.060	0.077	0.086	0.072	0.077	0.067	0.052	0.064	0.081	0.059	0.082	0.079	0.060	0.064	0.074	0.072	0.055	0.036	0.036
Sunday	17	0.076	0.070	0.053	0.070	0.075	0.058	0.075	0.062	0.049	0.065	0.066	0.051	0.076	0.070	0.053	0.063	0.068	0.071	0.052	0.037	0.035
Sunday	18	0.064	0.056	0.043	0.067	0.059	0.054	0.068	0.055	0.046	0.065	0.055	0.044	0.064	0.064	0.043	0.064	0.060	0.068	0.051	0.036	0.036
Sunday	19	0.049	0.043	0.035	0.062	0.045	0.050	0.061	0.047	0.042	0.062	0.043	0.036	0.049	0.043	0.035	0.060	0.052	0.062	0.048	0.048	0.037
Sunday	20	0.038	0.033	0.024	0.054	0.035	0.047	0.051	0.039	0.040	0.057	0.032	0.028	0.038	0.033	0.024	0.055	0.043	0.056	0.046	0.046	0.038
Sunday	21	0.026	0.022	0.040	0.045	0.024	0.039	0.041	0.028	0.036	0.049	0.022	0.023	0.026	0.026	0.022	0.020	0.050	0.046	0.046	0.038	0.038
Sunday	22	0.017	0.014	0.017	0.033	0.015	0.033	0.029	0.024	0.038	0.041	0.015	0.019	0.017	0.014	0.017	0.039	0.022	0.033	0.033	0.032	0.043
Sunday	23	0.010	0.010	0.020	0.022	0.009	0.032	0.019	0.019	0.037	0.028	0.012	0.016	0.010	0.010	0.020	0.030	0.016	0.020	0.020	0.027	0.050
Monday	0	0.006	0.010	0.017	0.010	0.003	0.007	0.011	0.017	0.023	0.023	0.007	0.013	0.006	0.013	0.023	0.024	0.006	0.010	0.010	0.024	0.031
Monday	1	0.004	0.009	0.016	0.009	0.002	0.007	0.007	0.015	0.022	0.023	0.006	0.011	0.004	0.009	0.016	0.024	0.007	0.005	0.023	0.031	0.031
Monday	2	0.003	0.009	0.016	0.010	0.003	0.010	0.006	0.015	0.022	0.025	0.007	0.011	0.003	0.009	0.016	0.025	0.009	0.004	0.022	0.030	0.030
Monday	3	0.005	0.011	0.019	0.012	0.006	0.012	0.009	0.018	0.025	0.027	0.010	0.011	0.008	0.017	0.024	0.023	0.011	0.005	0.023	0.032	0.032
Monday	4	0.008	0.017	0.024	0.014	0.009	0.013	0.018	0.027	0.032	0.030	0.015	0.012	0.005	0.017	0.024	0.035	0.023	0.014	0.030	0.037	0.037
Monday	5	0.019	0.028	0.036	0.022	0.022	0.026	0.030	0.039	0.039	0.033	0.022	0.018	0.019	0.028	0.036	0.039	0.042	0.039	0.041	0.041	0.044
Monday	6	0.036	0.041	0.050	0.037	0.047	0.044	0.044	0.051	0.045	0.036	0.034	0.024	0.036	0.041	0.050	0.044	0.060	0.050	0.050	0.049	0.051
Monday	7	0.051	0.044	0.065	0.045	0.058	0.058	0.058	0.058	0.050	0.040	0.043	0.030	0.051	0.044	0.065	0.041	0.056	0.059	0.058	0.056	0.056
Monday	8	0.053	0.056	0.068	0.047	0.062	0.067	0.053	0.058	0.051	0.043	0.054	0.039	0.053	0.056	0.068	0.043	0.058	0.055	0.055	0.056	0.055
Monday	9	0.059	0.065	0.080	0.050	0.065	0.078	0.051	0.059	0.053	0.045	0.067	0.048	0.059	0.065	0.080	0.045	0.065	0.053	0.053	0.057	0.058
Monday	10	0.067	0.074	0.087	0.051	0.065	0.080	0.054	0.062	0.056	0.050	0.074	0.054	0.067	0.074	0.087	0.046	0.065	0.055	0.055	0.060	0.058
Monday	11	0.071	0.075	0.082	0.056	0.067	0.083	0.057	0.064	0.057	0.052	0.075	0.059	0.071	0.075	0.082	0.050	0.066	0.057	0.057	0.058	0.058
Monday	12	0.074	0.074	0.080	0.058	0.069	0.081	0.060	0.064	0.058	0.055	0.078	0.059	0.074	0.074	0.080	0.052	0.068	0.058	0.060	0.060	0.059
Monday	13	0.074	0.075	0.075	0.063	0.074	0.076	0.061	0.064	0.058	0.057	0.081	0.060	0.074	0.075	0.075	0.056	0.069	0.059	0.059	0.059	0.055
Monday	14	0.077	0.076	0.065	0.067	0.076	0.074	0.067	0.066	0.058	0.057	0.081	0.065	0.077	0.076	0.065	0.057	0.070	0.064	0.064	0.058	0.053
Monday	15	0.082	0.076	0.058	0.073	0.087	0.062	0.072	0.065	0.057	0.059	0.080	0.063	0.082	0.076	0.058	0.058	0.070	0.068	0.068	0.058	0.050
Monday	16	0.081	0.073	0.045	0.076	0.084	0.053	0.075	0.063	0.055	0.060	0.072	0.064	0.081	0.073	0.045	0.059	0.067	0.071	0.071	0.058	0.046
Monday	17	0.071	0.059	0.035	0.075	0.075	0.040	0.074	0.055	0.051	0.057	0.059	0.066	0.071	0.059	0.035	0.058	0.062	0.070	0.070	0.054	0.042
Monday	18	0.057	0.042	0.023	0.047	0.032	0.032	0.055	0.042	0.042	0.053	0.045	0.063	0.052	0.042	0.023	0.055	0.043	0.055	0.055	0.041	0.035
Monday	19	0.037	0.030	0.017	0.050	0.031	0.029	0.042	0.031	0.036	0.048	0.032	0.060	0.037	0.030	0.017	0.045	0.029	0.048	0.043	0.032	0.028
Monday	20	0.027	0.022	0.013	0.043	0.020	0.021	0.034	0.023	0.031	0.042	0.022	0.054	0.027	0.022	0.013	0.041	0.022	0.035	0.035	0.026	0.024
Monday	21	0.020	0.016	0.010	0.035	0.015	0.020	0.027	0.018	0.028	0.036	0.016	0.046	0.020	0.016	0.010	0.035	0.017	0.030	0.030	0.022	0.021
Monday	22	0.015	0.012	0.009	0.025	0.009	0.014	0.020	0.014	0.027	0.029	0.012	0.039	0.015	0.012	0.009	0.026	0.011	0.023	0.023	0.018	0.022
Monday	23	0.009	0.007	0.010	0.016	0.005	0.013	0.014	0.011	0.025	0.020	0.008	0.031	0.009	0.007	0.010	0.020	0.006	0.016	0.015	0.015	0.025
Tues/Wed/Thurs	0	0.005	0.009	0.017	0.010	0.004	0.008	0.008	0.016	0.025	0.023	0.007	0.018	0.005	0.009	0.017	0.020	0.006	0.005	0.023	0.033	0.033
Tues/Wed/Thurs	1	0.003	0.008	0.017	0.009	0.003	0.008	0.005	0.014	0.024	0.024	0.006	0.015	0.003	0.008	0.017	0.022	0.006	0.005	0.021	0.031	0.031
Tues/Wed/Thurs	2	0.002	0.009	0.017	0.010	0.002	0.012	0.005	0.014	0.025	0.027	0.006	0.013	0.002	0.009	0.017	0.023	0.007	0.004	0.021	0.031	0.031
Tues/Wed/Thurs	3	0.003	0.010	0.022	0.011	0.005	0.014	0.008	0.018	0.028	0.029	0.009	0.013	0.003	0.010	0.022	0.025	0.010	0.005	0.022	0.022	0.032

Day of Week	Hour	Mariposa			Mendocino			Merced			Modoc			Mono			Monterey			Napa		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Tues/Wed/Thurs	4	0.006	0.014	0.025	0.015	0.020	0.021	0.017	0.026	0.034	0.014	0.016	0.006	0.014	0.025	0.030	0.019	0.024	0.013	0.028	0.039	
Tues/Wed/Thurs	5	0.018	0.027	0.039	0.024	0.024	0.035	0.030	0.039	0.047	0.035	0.020	0.018	0.027	0.039	0.037	0.037	0.029	0.036	0.040	0.046	
Tues/Wed/Thurs	6	0.037	0.042	0.052	0.037	0.048	0.048	0.044	0.050	0.047	0.033	0.027	0.037	0.042	0.052	0.043	0.037	0.038	0.048	0.048	0.051	
Tues/Wed/Thurs	7	0.053	0.047	0.064	0.045	0.059	0.065	0.059	0.059	0.052	0.046	0.036	0.053	0.047	0.064	0.042	0.045	0.046	0.059	0.056	0.056	
Tues/Wed/Thurs	8	0.054	0.056	0.070	0.047	0.063	0.069	0.055	0.058	0.052	0.056	0.046	0.054	0.056	0.070	0.045	0.062	0.050	0.056	0.057	0.057	
Tues/Wed/Thurs	9	0.059	0.068	0.083	0.050	0.064	0.074	0.051	0.059	0.054	0.066	0.057	0.059	0.068	0.083	0.046	0.063	0.055	0.052	0.055	0.056	
Tues/Wed/Thurs	10	0.064	0.069	0.081	0.051	0.065	0.075	0.052	0.060	0.056	0.071	0.065	0.064	0.069	0.081	0.047	0.061	0.058	0.053	0.057	0.057	
Tues/Wed/Thurs	11	0.068	0.069	0.077	0.055	0.065	0.076	0.054	0.061	0.057	0.076	0.070	0.068	0.069	0.077	0.049	0.065	0.060	0.053	0.058	0.057	
Tues/Wed/Thurs	12	0.069	0.071	0.074	0.057	0.068	0.076	0.057	0.062	0.057	0.076	0.070	0.069	0.071	0.074	0.051	0.066	0.060	0.055	0.058	0.056	
Tues/Wed/Thurs	13	0.072	0.073	0.074	0.061	0.070	0.071	0.060	0.063	0.056	0.077	0.069	0.072	0.074	0.074	0.054	0.069	0.059	0.057	0.060	0.055	
Tues/Wed/Thurs	14	0.077	0.076	0.067	0.066	0.074	0.068	0.066	0.065	0.056	0.081	0.067	0.077	0.076	0.067	0.058	0.072	0.059	0.064	0.061	0.053	
Tues/Wed/Thurs	15	0.084	0.078	0.058	0.073	0.084	0.062	0.073	0.066	0.055	0.078	0.064	0.084	0.078	0.058	0.059	0.072	0.057	0.069	0.061	0.050	
Tues/Wed/Thurs	16	0.082	0.074	0.048	0.078	0.086	0.053	0.077	0.064	0.053	0.057	0.072	0.082	0.074	0.048	0.060	0.070	0.053	0.072	0.058	0.046	
Tues/Wed/Thurs	17	0.074	0.061	0.036	0.077	0.078	0.041	0.076	0.057	0.049	0.060	0.057	0.074	0.061	0.036	0.058	0.063	0.051	0.072	0.055	0.041	
Tues/Wed/Thurs	18	0.053	0.044	0.023	0.059	0.047	0.030	0.058	0.044	0.041	0.046	0.053	0.053	0.044	0.023	0.052	0.044	0.046	0.058	0.044	0.035	
Tues/Wed/Thurs	19	0.038	0.031	0.016	0.048	0.031	0.027	0.044	0.032	0.034	0.048	0.033	0.044	0.038	0.031	0.016	0.043	0.024	0.047	0.035	0.028	
Tues/Wed/Thurs	20	0.030	0.025	0.012	0.041	0.021	0.020	0.036	0.025	0.030	0.045	0.025	0.038	0.030	0.025	0.012	0.043	0.024	0.039	0.029	0.024	
Tues/Wed/Thurs	21	0.023	0.018	0.010	0.036	0.017	0.020	0.028	0.019	0.026	0.038	0.018	0.032	0.023	0.018	0.010	0.038	0.018	0.033	0.022	0.021	
Tues/Wed/Thurs	22	0.017	0.013	0.010	0.025	0.009	0.014	0.021	0.014	0.025	0.032	0.014	0.026	0.017	0.013	0.010	0.029	0.011	0.025	0.018	0.022	
Tues/Wed/Thurs	23	0.010	0.008	0.010	0.017	0.005	0.012	0.015	0.012	0.023	0.025	0.010	0.021	0.010	0.008	0.010	0.022	0.008	0.017	0.015	0.025	
Friday	0	0.005	0.009	0.019	0.009	0.004	0.008	0.008	0.016	0.027	0.021	0.007	0.019	0.005	0.009	0.019	0.020	0.006	0.009	0.022	0.034	
Friday	1	0.003	0.008	0.019	0.009	0.003	0.009	0.006	0.014	0.025	0.023	0.006	0.017	0.003	0.006	0.019	0.020	0.006	0.005	0.022	0.032	
Friday	2	0.002	0.008	0.019	0.009	0.003	0.011	0.005	0.014	0.026	0.024	0.007	0.016	0.002	0.008	0.019	0.022	0.007	0.004	0.021	0.034	
Friday	3	0.002	0.008	0.021	0.011	0.005	0.016	0.008	0.017	0.029	0.026	0.009	0.016	0.002	0.008	0.021	0.024	0.009	0.005	0.022	0.034	
Friday	4	0.005	0.013	0.024	0.013	0.009	0.022	0.014	0.024	0.035	0.029	0.013	0.019	0.005	0.013	0.024	0.028	0.018	0.011	0.026	0.039	
Friday	5	0.013	0.023	0.037	0.021	0.021	0.039	0.024	0.035	0.042	0.032	0.018	0.023	0.013	0.023	0.037	0.035	0.033	0.029	0.038	0.046	
Friday	6	0.026	0.035	0.049	0.033	0.041	0.054	0.036	0.045	0.047	0.033	0.030	0.032	0.026	0.035	0.049	0.041	0.050	0.039	0.045	0.052	
Friday	7	0.039	0.040	0.060	0.039	0.052	0.065	0.049	0.053	0.052	0.037	0.039	0.039	0.039	0.049	0.060	0.049	0.046	0.048	0.051	0.057	
Friday	8	0.043	0.049	0.068	0.044	0.059	0.074	0.047	0.054	0.053	0.040	0.051	0.049	0.043	0.049	0.068	0.041	0.056	0.047	0.051	0.057	
Friday	9	0.049	0.057	0.073	0.047	0.060	0.078	0.047	0.056	0.055	0.045	0.063	0.054	0.049	0.057	0.073	0.045	0.058	0.047	0.055	0.058	
Friday	10	0.058	0.063	0.078	0.048	0.067	0.075	0.051	0.060	0.058	0.048	0.060	0.058	0.058	0.063	0.078	0.047	0.062	0.052	0.057	0.059	
Friday	11	0.064	0.069	0.077	0.054	0.068	0.077	0.054	0.062	0.060	0.049	0.072	0.063	0.064	0.069	0.077	0.050	0.067	0.055	0.058	0.059	
Friday	12	0.066	0.071	0.076	0.060	0.072	0.079	0.057	0.063	0.060	0.052	0.074	0.063	0.066	0.071	0.076	0.051	0.067	0.059	0.060	0.058	
Friday	13	0.071	0.074	0.077	0.063	0.075	0.072	0.061	0.065	0.059	0.055	0.077	0.062	0.071	0.074	0.077	0.062	0.064	0.061	0.061	0.052	
Friday	14	0.076	0.077	0.070	0.068	0.078	0.067	0.068	0.067	0.058	0.059	0.080	0.063	0.076	0.077	0.070	0.060	0.075	0.067	0.061	0.051	
Friday	15	0.083	0.079	0.060	0.073	0.083	0.060	0.074	0.067	0.056	0.063	0.081	0.061	0.083	0.079	0.060	0.060	0.074	0.069	0.061	0.048	
Friday	16	0.083	0.077	0.050	0.076	0.082	0.049	0.076	0.064	0.053	0.058	0.075	0.059	0.083	0.077	0.050	0.070	0.055	0.069	0.058	0.048	
Friday	17	0.075	0.064	0.038	0.074	0.072	0.038	0.075	0.058	0.048	0.059	0.063	0.055	0.075	0.064	0.038	0.060	0.064	0.068	0.051	0.040	
Friday	18	0.062	0.051	0.025	0.060	0.050	0.026	0.064	0.048	0.040	0.054	0.052	0.051	0.062	0.051	0.025	0.054	0.049	0.060	0.046	0.034	
Friday	19	0.050	0.039	0.018	0.052	0.034	0.024	0.052	0.037	0.032	0.050	0.036	0.046	0.050	0.039	0.018	0.050	0.036	0.054	0.039	0.027	
Friday	20	0.041	0.030	0.013	0.043	0.022	0.017	0.043	0.029	0.026	0.046	0.030	0.041	0.041	0.030	0.013	0.045	0.028	0.048	0.033	0.023	
Friday	21	0.036	0.025	0.010	0.040	0.018	0.016	0.035	0.022	0.022	0.040	0.022	0.036	0.036	0.025	0.010	0.038	0.021	0.039	0.026	0.019	
Friday	22	0.030	0.019	0.011	0.031	0.012	0.011	0.027	0.016	0.020	0.031	0.016	0.031	0.030	0.019	0.011	0.031	0.015	0.031	0.020	0.020	
Friday	23	0.018	0.012	0.009	0.022	0.007	0.012	0.020	0.012	0.018	0.025	0.012	0.025	0.018	0.012	0.009	0.023	0.010	0.022	0.016	0.021	
Saturday	0	0.010	0.015	0.027	0.012	0.008	0.014	0.015	0.026	0.040	0.026	0.013	0.020	0.010	0.015	0.027	0.023	0.011	0.014	0.029	0.051	
Saturday	1	0.007	0.012	0.023	0.013	0.006	0.014	0.010	0.020	0.035	0.026	0.008	0.016	0.007	0.012	0.023	0.025	0.010	0.009	0.024	0.044	
Saturday	2	0.005	0.011	0.022	0.013	0.004	0.011	0.008	0.018	0.032	0.027	0.007	0.015	0.005	0.011	0.022	0.025	0.009	0.007	0.022	0.041	
Saturday	3	0.004	0.010	0.025	0.012	0.004	0.014	0.008	0.019	0.032	0.030	0.007	0.014	0.004	0.010	0.025	0.027	0.011	0.006	0.023	0.040	
Saturday	4	0.005	0.013	0.028	0.014	0.008	0.020	0.011	0.021	0.035	0.029	0.009	0.016	0.005	0.013	0.028	0.031	0.020	0.007	0.023	0.041	
Saturday	5	0.010	0.021	0.034	0.020	0.016	0.034	0.017	0.028	0.039	0.033	0.015	0.019	0.010	0.021	0.034	0.038	0.029	0.013	0.029	0.045	
Saturday	6	0.017	0.028	0.039	0.025	0.025	0.043	0.025	0.036	0.045	0.036	0.023	0.025	0.017	0.028	0.039	0.038	0.047	0.021	0.033	0.047	
Saturday	7	0.029	0.036	0.053	0.030	0.031	0.058	0.034	0.044	0.050	0.038	0.033	0.036	0.029	0.036	0.053	0.042	0.047	0.030	0.038	0.053	

Day of Week	Hour	Mariposa			Mendocino			Merced			Modoc			Mono			Monterey			Napa			
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	
Saturday	8	0.044	0.045	0.060	0.036	0.041	0.070	0.044	0.053	0.055	0.041	0.047	0.047	0.044	0.045	0.060	0.043	0.055	0.043	0.050	0.042	0.046	0.052
Saturday	9	0.059	0.061	0.071	0.043	0.053	0.079	0.054	0.061	0.060	0.045	0.063	0.059	0.059	0.061	0.071	0.047	0.062	0.062	0.055	0.054	0.054	0.058
Saturday	10	0.073	0.074	0.078	0.052	0.069	0.082	0.062	0.068	0.063	0.049	0.075	0.067	0.073	0.074	0.078	0.047	0.067	0.062	0.062	0.063	0.058	0.055
Saturday	11	0.081	0.077	0.083	0.054	0.076	0.075	0.067	0.071	0.064	0.050	0.084	0.073	0.081	0.077	0.083	0.049	0.068	0.063	0.063	0.068	0.060	0.052
Saturday	12	0.078	0.077	0.075	0.061	0.080	0.070	0.069	0.070	0.062	0.053	0.083	0.071	0.078	0.077	0.075	0.055	0.071	0.060	0.060	0.069	0.060	0.052
Saturday	13	0.075	0.072	0.060	0.063	0.082	0.064	0.070	0.067	0.058	0.055	0.081	0.069	0.075	0.072	0.060	0.054	0.070	0.059	0.067	0.057	0.047	
Saturday	14	0.075	0.068	0.055	0.065	0.081	0.062	0.070	0.064	0.054	0.057	0.076	0.062	0.075	0.068	0.055	0.055	0.065	0.058	0.067	0.057	0.044	
Saturday	15	0.075	0.068	0.052	0.067	0.080	0.054	0.069	0.061	0.049	0.060	0.074	0.062	0.075	0.068	0.052	0.055	0.065	0.056	0.067	0.057	0.044	
Saturday	16	0.072	0.070	0.047	0.071	0.081	0.051	0.068	0.057	0.045	0.056	0.070	0.058	0.072	0.070	0.047	0.057	0.065	0.052	0.068	0.054	0.038	
Saturday	17	0.066	0.063	0.040	0.062	0.072	0.037	0.064	0.051	0.040	0.055	0.061	0.057	0.066	0.063	0.040	0.056	0.053	0.047	0.066	0.054	0.035	
Saturday	18	0.058	0.052	0.031	0.062	0.053	0.032	0.056	0.042	0.033	0.051	0.049	0.052	0.058	0.052	0.031	0.052	0.044	0.042	0.060	0.049	0.032	
Saturday	19	0.047	0.041	0.026	0.059	0.040	0.029	0.048	0.034	0.027	0.049	0.038	0.045	0.047	0.041	0.026	0.049	0.039	0.052	0.044	0.044	0.030	
Saturday	20	0.038	0.031	0.020	0.051	0.032	0.021	0.041	0.029	0.024	0.042	0.031	0.038	0.038	0.031	0.020	0.043	0.031	0.035	0.046	0.040	0.028	
Saturday	21	0.031	0.025	0.016	0.047	0.026	0.023	0.037	0.024	0.021	0.037	0.023	0.031	0.031	0.025	0.016	0.038	0.025	0.029	0.042	0.035	0.025	
Saturday	22	0.025	0.020	0.018	0.037	0.019	0.020	0.031	0.020	0.019	0.031	0.017	0.026	0.025	0.020	0.018	0.030	0.017	0.026	0.036	0.030	0.023	
Saturday	23	0.016	0.013	0.018	0.028	0.014	0.021	0.023	0.016	0.017	0.023	0.012	0.019	0.016	0.013	0.018	0.023	0.011	0.020	0.026	0.024	0.024	
Holiday	0	0.008	0.011	0.020	0.010	0.004	0.009	0.013	0.020	0.027	0.024	0.008	0.015	0.008	0.011	0.020	0.024	0.008	0.014	0.028	0.028	0.038	
Holiday	1	0.005	0.009	0.018	0.014	0.004	0.008	0.009	0.017	0.025	0.027	0.008	0.012	0.005	0.009	0.018	0.022	0.009	0.008	0.008	0.024	0.033	
Holiday	2	0.003	0.010	0.018	0.010	0.003	0.014	0.007	0.015	0.024	0.024	0.008	0.012	0.003	0.010	0.018	0.024	0.007	0.005	0.005	0.026	0.033	
Holiday	3	0.004	0.010	0.021	0.014	0.005	0.012	0.007	0.016	0.026	0.029	0.010	0.013	0.004	0.010	0.021	0.024	0.009	0.004	0.004	0.025	0.034	
Holiday	4	0.005	0.012	0.020	0.014	0.006	0.017	0.011	0.020	0.029	0.029	0.012	0.014	0.005	0.012	0.020	0.031	0.019	0.008	0.008	0.025	0.035	
Holiday	5	0.009	0.018	0.031	0.019	0.018	0.028	0.019	0.028	0.033	0.031	0.016	0.017	0.009	0.018	0.031	0.033	0.029	0.017	0.017	0.030	0.040	
Holiday	6	0.018	0.023	0.038	0.028	0.034	0.042	0.027	0.035	0.038	0.037	0.025	0.023	0.018	0.023	0.038	0.038	0.042	0.024	0.024	0.036	0.044	
Holiday	7	0.029	0.031	0.043	0.039	0.045	0.052	0.035	0.042	0.042	0.038	0.033	0.031	0.029	0.031	0.043	0.040	0.044	0.037	0.030	0.042	0.049	
Holiday	8	0.041	0.044	0.056	0.041	0.051	0.059	0.040	0.048	0.046	0.040	0.049	0.040	0.041	0.044	0.056	0.037	0.050	0.041	0.039	0.047	0.049	
Holiday	9	0.058	0.057	0.075	0.044	0.057	0.066	0.048	0.055	0.050	0.043	0.062	0.054	0.058	0.057	0.075	0.046	0.057	0.048	0.048	0.055	0.057	
Holiday	10	0.076	0.083	0.087	0.050	0.069	0.075	0.059	0.064	0.055	0.050	0.076	0.060	0.076	0.083	0.087	0.048	0.066	0.056	0.060	0.060	0.056	
Holiday	11	0.084	0.086	0.088	0.056	0.072	0.077	0.065	0.070	0.060	0.047	0.084	0.068	0.084	0.086	0.088	0.055	0.077	0.063	0.066	0.064	0.055	
Holiday	12	0.085	0.087	0.089	0.058	0.080	0.078	0.069	0.072	0.061	0.053	0.083	0.070	0.085	0.087	0.089	0.052	0.074	0.065	0.068	0.063	0.060	
Holiday	13	0.083	0.081	0.078	0.063	0.077	0.069	0.071	0.071	0.061	0.062	0.091	0.067	0.083	0.081	0.078	0.055	0.071	0.069	0.069	0.062	0.055	
Holiday	14	0.080	0.074	0.068	0.068	0.083	0.067	0.072	0.069	0.059	0.059	0.087	0.069	0.080	0.074	0.068	0.050	0.071	0.067	0.071	0.060	0.055	
Holiday	15	0.078	0.074	0.060	0.071	0.082	0.064	0.073	0.068	0.058	0.057	0.079	0.065	0.078	0.074	0.060	0.061	0.068	0.068	0.071	0.064	0.054	
Holiday	16	0.078	0.072	0.049	0.075	0.083	0.061	0.073	0.065	0.055	0.056	0.072	0.062	0.078	0.072	0.049	0.062	0.069	0.068	0.068	0.057	0.046	
Holiday	17	0.071	0.066	0.041	0.072	0.076	0.044	0.070	0.057	0.050	0.056	0.058	0.060	0.071	0.066	0.041	0.058	0.062	0.058	0.067	0.055	0.041	
Holiday	18	0.057	0.049	0.033	0.054	0.048	0.040	0.060	0.046	0.044	0.053	0.044	0.058	0.057	0.049	0.033	0.054	0.050	0.049	0.061	0.042	0.038	
Holiday	19	0.043	0.040	0.022	0.056	0.036	0.029	0.050	0.036	0.039	0.048	0.029	0.049	0.043	0.040	0.022	0.049	0.037	0.053	0.053	0.037	0.029	
Holiday	20	0.033	0.026	0.013	0.049	0.025	0.029	0.042	0.029	0.034	0.044	0.024	0.045	0.033	0.026	0.013	0.046	0.032	0.043	0.049	0.029	0.024	
Holiday	21	0.024	0.018	0.011	0.040	0.019	0.023	0.034	0.023	0.030	0.040	0.019	0.040	0.024	0.018	0.011	0.040	0.025	0.038	0.042	0.028	0.024	
Holiday	22	0.017	0.012	0.009	0.029	0.012	0.018	0.027	0.017	0.028	0.031	0.014	0.030	0.017	0.012	0.009	0.031	0.016	0.032	0.035	0.022	0.025	
Holiday	23	0.010	0.008	0.010	0.025	0.010	0.019	0.018	0.014	0.026	0.024	0.009	0.024	0.010	0.008	0.010	0.020	0.008	0.023	0.023	0.018	0.026	

Day of Week	Hour	Nevada			Orange			Placer			Plumas			Riverside			Sacramento			San Benito		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Sunday	0	0.013	0.020	0.031	0.023	0.045	0.061	0.013	0.020	0.031	0.015	0.010	0.015	0.022	0.036	0.050	0.019	0.031	0.044	0.019	0.010	0.029
Sunday	1	0.008	0.016	0.028	0.015	0.032	0.049	0.008	0.016	0.028	0.010	0.006	0.011	0.015	0.028	0.044	0.013	0.025	0.039	0.020	0.008	0.023
Sunday	2	0.006	0.013	0.026	0.011	0.025	0.041	0.006	0.013	0.026	0.004	0.004	0.012	0.011	0.023	0.040	0.009	0.021	0.036	0.020	0.007	0.021
Sunday	3	0.005	0.012	0.025	0.007	0.019	0.034	0.005	0.012	0.025	0.006	0.004	0.012	0.009	0.020	0.036	0.007	0.019	0.034	0.020	0.007	0.019
Sunday	4	0.005	0.012	0.025	0.007	0.018	0.031	0.005	0.012	0.025	0.006	0.005	0.017	0.009	0.020	0.035	0.008	0.020	0.034	0.024	0.012	0.019
Sunday	5	0.008	0.015	0.027	0.011	0.022	0.034	0.008	0.015	0.027	0.010	0.011	0.029	0.012	0.023	0.036	0.011	0.023	0.034	0.026	0.017	0.021
Sunday	6	0.013	0.020	0.030	0.013	0.029	0.038	0.013	0.020	0.030	0.016	0.017	0.037	0.019	0.029	0.039	0.019	0.027	0.037	0.029	0.024	0.026
Sunday	7	0.022	0.028	0.034	0.026	0.036	0.041	0.022	0.028	0.034	0.023	0.029	0.051	0.026	0.035	0.041	0.025	0.033	0.039	0.031	0.030	0.034
Sunday	8	0.034	0.041	0.040	0.037	0.046	0.046	0.034	0.041	0.040	0.033	0.043	0.071	0.036	0.045	0.044	0.035	0.042	0.043	0.035	0.038	0.040
Sunday	9	0.048	0.055	0.046	0.050	0.058	0.051	0.048	0.055	0.046	0.047	0.063	0.091	0.049	0.054	0.047	0.049	0.052	0.049	0.038	0.049	0.049
Sunday	10	0.064	0.068	0.052	0.059	0.065	0.052	0.064	0.068	0.052	0.057	0.075	0.084	0.057	0.061	0.047	0.060	0.060	0.049	0.041	0.057	0.057
Sunday	11	0.075	0.075	0.055	0.065	0.067	0.052	0.075	0.075	0.055	0.067	0.083	0.079	0.064	0.065	0.048	0.066	0.063	0.049	0.047	0.068	0.061
Sunday	12	0.082	0.079	0.058	0.068	0.066	0.049	0.082	0.079	0.058	0.074	0.090	0.070	0.067	0.066	0.047	0.072	0.066	0.049	0.051	0.074	0.063
Sunday	13	0.084	0.079	0.058	0.069	0.064	0.046	0.084	0.079	0.058	0.078	0.089	0.061	0.069	0.065	0.045	0.074	0.067	0.049	0.053	0.073	0.065
Sunday	14	0.084	0.077	0.057	0.068	0.059	0.043	0.084	0.077	0.057	0.079	0.081	0.057	0.069	0.063	0.044	0.074	0.064	0.047	0.059	0.078	0.065
Sunday	15	0.082	0.073	0.057	0.068	0.055	0.040	0.082	0.073	0.057	0.080	0.079	0.045	0.068	0.060	0.042	0.071	0.061	0.046	0.061	0.078	0.066
Sunday	16	0.079	0.068	0.055	0.067	0.051	0.038	0.079	0.068	0.055	0.079	0.075	0.045	0.067	0.056	0.041	0.071	0.059	0.045	0.064	0.074	0.060
Sunday	17	0.072	0.062	0.053	0.064	0.047	0.036	0.072	0.062	0.053	0.075	0.066	0.043	0.064	0.052	0.040	0.068	0.056	0.043	0.063	0.068	0.053
Sunday	18	0.060	0.052	0.049	0.060	0.041	0.034	0.060	0.052	0.049	0.066	0.054	0.039	0.061	0.047	0.039	0.061	0.049	0.041	0.064	0.060	0.049
Sunday	19	0.050	0.043	0.045	0.055	0.036	0.033	0.050	0.043	0.045	0.055	0.042	0.037	0.057	0.042	0.039	0.053	0.042	0.040	0.060	0.052	0.046
Sunday	20	0.041	0.035	0.042	0.052	0.034	0.034	0.041	0.035	0.042	0.045	0.031	0.030	0.053	0.037	0.039	0.048	0.038	0.039	0.055	0.043	0.041
Sunday	21	0.031	0.026	0.039	0.045	0.032	0.036	0.031	0.026	0.039	0.035	0.022	0.024	0.044	0.032	0.038	0.040	0.032	0.039	0.050	0.034	0.037
Sunday	22	0.021	0.019	0.036	0.034	0.028	0.038	0.021	0.019	0.036	0.023	0.013	0.018	0.042	0.024	0.038	0.029	0.027	0.038	0.039	0.022	0.031
Sunday	23	0.013	0.015	0.033	0.022	0.024	0.042	0.013	0.015	0.033	0.014	0.008	0.015	0.021	0.018	0.038	0.019	0.023	0.039	0.030	0.016	0.025
Monday	0	0.008	0.014	0.027	0.010	0.016	0.024	0.008	0.014	0.027	0.006	0.002	0.006	0.011	0.018	0.028	0.009	0.018	0.028	0.023	0.006	0.009
Monday	1	0.005	0.012	0.025	0.006	0.012	0.021	0.005	0.012	0.025	0.004	0.002	0.007	0.008	0.016	0.026	0.005	0.015	0.026	0.024	0.007	0.009
Monday	2	0.004	0.012	0.025	0.005	0.012	0.021	0.004	0.012	0.025	0.003	0.002	0.010	0.007	0.016	0.027	0.004	0.015	0.026	0.025	0.009	0.010
Monday	3	0.006	0.014	0.027	0.006	0.013	0.022	0.006	0.014	0.027	0.003	0.004	0.012	0.011	0.020	0.030	0.006	0.018	0.028	0.025	0.011	0.014
Monday	4	0.011	0.019	0.030	0.015	0.022	0.029	0.011	0.019	0.030	0.007	0.009	0.021	0.024	0.033	0.038	0.013	0.026	0.033	0.033	0.023	0.019
Monday	5	0.023	0.030	0.036	0.034	0.041	0.043	0.023	0.030	0.036	0.018	0.024	0.037	0.040	0.049	0.045	0.029	0.040	0.040	0.039	0.042	0.024
Monday	6	0.042	0.047	0.043	0.054	0.060	0.054	0.042	0.047	0.043	0.041	0.051	0.055	0.055	0.059	0.049	0.052	0.057	0.048	0.044	0.060	0.031
Monday	7	0.060	0.061	0.048	0.066	0.073	0.060	0.060	0.061	0.048	0.078	0.069	0.066	0.059	0.064	0.051	0.071	0.066	0.051	0.041	0.056	0.038
Monday	8	0.059	0.062	0.050	0.064	0.073	0.061	0.059	0.062	0.050	0.067	0.077	0.077	0.056	0.062	0.052	0.066	0.064	0.052	0.043	0.058	0.045
Monday	9	0.056	0.061	0.050	0.056	0.065	0.058	0.056	0.061	0.050	0.057	0.071	0.080	0.053	0.059	0.051	0.056	0.059	0.052	0.045	0.063	0.053
Monday	10	0.058	0.064	0.051	0.052	0.061	0.055	0.058	0.064	0.051	0.057	0.071	0.077	0.052	0.058	0.051	0.052	0.057	0.052	0.046	0.065	0.059
Monday	11	0.062	0.066	0.053	0.052	0.060	0.055	0.062	0.066	0.053	0.060	0.074	0.073	0.053	0.058	0.052	0.053	0.058	0.053	0.050	0.066	0.061
Monday	12	0.066	0.068	0.054	0.053	0.060	0.054	0.066	0.068	0.054	0.063	0.072	0.071	0.055	0.058	0.051	0.056	0.059	0.053	0.052	0.068	0.065
Monday	13	0.067	0.067	0.054	0.055	0.059	0.053	0.067	0.067	0.054	0.063	0.072	0.068	0.057	0.059	0.051	0.057	0.059	0.053	0.056	0.069	0.063
Monday	14	0.070	0.069	0.055	0.060	0.061	0.054	0.070	0.069	0.055	0.067	0.077	0.064	0.061	0.060	0.051	0.062	0.060	0.053	0.057	0.070	0.065
Monday	15	0.073	0.069	0.055	0.064	0.061	0.053	0.073	0.069	0.055	0.078	0.080	0.056	0.065	0.061	0.050	0.070	0.064	0.052	0.058	0.070	0.066
Monday	16	0.075	0.067	0.054	0.067	0.060	0.052	0.075	0.067	0.054	0.086	0.077	0.049	0.067	0.059	0.049	0.076	0.063	0.051	0.059	0.067	0.060
Monday	17	0.073	0.061	0.052	0.068	0.057	0.050	0.073	0.061	0.052	0.087	0.062	0.041	0.066	0.054	0.047	0.073	0.057	0.048	0.058	0.062	0.057
Monday	18	0.056	0.046	0.045	0.060	0.044	0.042	0.056	0.046	0.045	0.051	0.038	0.030	0.056	0.043	0.042	0.056	0.044	0.043	0.055	0.049	0.048
Monday	19	0.040	0.031	0.039	0.047	0.029	0.034	0.040	0.031	0.039	0.036	0.024	0.024	0.044	0.034	0.037	0.040	0.031	0.037	0.045	0.029	0.048
Monday	20	0.031	0.022	0.035	0.037	0.020	0.028	0.031	0.022	0.035	0.026	0.018	0.023	0.035	0.023	0.033	0.032	0.024	0.033	0.041	0.022	0.045
Monday	21	0.025	0.017	0.032	0.032	0.017	0.026	0.025	0.017	0.032	0.020	0.012	0.021	0.030	0.017	0.031	0.028	0.019	0.030	0.035	0.017	0.039
Monday	22	0.017	0.012	0.030	0.024	0.013	0.025	0.017	0.012	0.030	0.013	0.007	0.017	0.020	0.015	0.029	0.021	0.015	0.028	0.026	0.011	0.035
Monday	23	0.012	0.009	0.030	0.015	0.010	0.026	0.012	0.009	0.030	0.008	0.004	0.015	0.016	0.009	0.028	0.014	0.011	0.027	0.020	0.007	0.033
Tues/Wed/Thurs	0	0.008	0.014	0.029	0.009	0.015	0.026	0.008	0.014	0.029	0.006	0.003	0.010	0.010	0.017	0.030	0.008	0.018	0.031	0.020	0.006	0.023
Tues/Wed/Thurs	1	0.004	0.011	0.027	0.005	0.012	0.024	0.004	0.011	0.027	0.003	0.002	0.011	0.007	0.015	0.030	0.005	0.015	0.030	0.022	0.006	0.021
Tues/Wed/Thurs	2	0.004	0.011	0.027	0.004	0.012	0.023	0.004	0.011	0.027	0.003	0.002	0.013	0.006	0.015	0.029	0.004	0.015	0.029	0.023	0.007	0.021
Tues/Wed/Thurs	3	0.005	0.013	0.029	0.005	0.013	0.025	0.005	0.013	0.029	0.003	0.003	0.015	0.010	0.019	0.032	0.006	0.017	0.031	0.025	0.010	0.022



Day of Week	Hour	Nevada			Orange			Placer			Plumas			Riverside			Sacramento			San Benito		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Tues/Wed/Thurs	4	0.010	0.018	0.031	0.022	0.033	0.040	0.031	0.018	0.031	0.006	0.008	0.022	0.022	0.032	0.040	0.012	0.024	0.036	0.030	0.019	0.024
Tues/Wed/Thurs	5	0.022	0.029	0.037	0.033	0.040	0.045	0.022	0.029	0.037	0.017	0.024	0.037	0.039	0.048	0.047	0.027	0.038	0.043	0.037	0.037	0.029
Tues/Wed/Thurs	6	0.042	0.047	0.044	0.054	0.061	0.057	0.042	0.047	0.044	0.041	0.053	0.054	0.053	0.060	0.051	0.052	0.057	0.050	0.043	0.057	0.038
Tues/Wed/Thurs	7	0.060	0.061	0.050	0.063	0.073	0.062	0.060	0.061	0.050	0.077	0.069	0.066	0.059	0.064	0.053	0.071	0.066	0.053	0.042	0.057	0.046
Tues/Wed/Thurs	8	0.060	0.062	0.051	0.065	0.073	0.062	0.060	0.062	0.051	0.066	0.071	0.077	0.056	0.062	0.053	0.066	0.063	0.053	0.045	0.062	0.050
Tues/Wed/Thurs	9	0.055	0.060	0.050	0.057	0.066	0.059	0.055	0.060	0.050	0.057	0.071	0.080	0.052	0.059	0.052	0.056	0.059	0.053	0.046	0.063	0.055
Tues/Wed/Thurs	10	0.056	0.061	0.051	0.052	0.061	0.056	0.056	0.061	0.051	0.056	0.071	0.077	0.051	0.058	0.052	0.051	0.057	0.053	0.047	0.061	0.058
Tues/Wed/Thurs	11	0.059	0.064	0.052	0.052	0.061	0.054	0.059	0.064	0.052	0.058	0.071	0.074	0.051	0.058	0.051	0.052	0.057	0.053	0.049	0.065	0.060
Tues/Wed/Thurs	12	0.061	0.065	0.053	0.053	0.060	0.053	0.061	0.065	0.053	0.062	0.070	0.069	0.053	0.058	0.051	0.054	0.058	0.053	0.051	0.066	0.060
Tues/Wed/Thurs	13	0.064	0.066	0.053	0.055	0.060	0.052	0.064	0.066	0.053	0.063	0.073	0.067	0.056	0.056	0.051	0.056	0.059	0.052	0.054	0.069	0.059
Tues/Wed/Thurs	14	0.068	0.068	0.053	0.059	0.061	0.052	0.068	0.068	0.053	0.066	0.076	0.063	0.060	0.061	0.050	0.061	0.061	0.051	0.058	0.072	0.059
Tues/Wed/Thurs	15	0.073	0.069	0.053	0.063	0.061	0.051	0.073	0.069	0.053	0.079	0.080	0.056	0.064	0.061	0.048	0.070	0.064	0.050	0.059	0.072	0.057
Tues/Wed/Thurs	16	0.075	0.067	0.052	0.065	0.059	0.049	0.075	0.067	0.052	0.087	0.076	0.045	0.066	0.060	0.047	0.075	0.063	0.048	0.060	0.070	0.053
Tues/Wed/Thurs	17	0.074	0.063	0.050	0.066	0.055	0.046	0.074	0.063	0.050	0.088	0.062	0.040	0.066	0.055	0.044	0.073	0.057	0.044	0.058	0.063	0.051
Tues/Wed/Thurs	18	0.059	0.048	0.044	0.060	0.044	0.040	0.059	0.048	0.044	0.054	0.039	0.031	0.058	0.045	0.040	0.059	0.046	0.041	0.052	0.044	0.046
Tues/Wed/Thurs	19	0.043	0.034	0.038	0.049	0.030	0.032	0.043	0.034	0.038	0.036	0.026	0.023	0.046	0.032	0.032	0.034	0.026	0.031	0.049	0.032	0.041
Tues/Wed/Thurs	20	0.035	0.025	0.034	0.040	0.021	0.027	0.035	0.025	0.034	0.028	0.019	0.021	0.038	0.024	0.032	0.034	0.026	0.031	0.043	0.024	0.037
Tues/Wed/Thurs	21	0.029	0.019	0.031	0.035	0.017	0.025	0.029	0.019	0.031	0.021	0.013	0.020	0.033	0.018	0.029	0.030	0.021	0.029	0.038	0.018	0.034
Tues/Wed/Thurs	22	0.020	0.013	0.029	0.026	0.013	0.024	0.020	0.013	0.029	0.014	0.007	0.016	0.025	0.012	0.027	0.022	0.016	0.027	0.029	0.011	0.030
Tues/Wed/Thurs	23	0.013	0.009	0.028	0.016	0.010	0.025	0.013	0.009	0.028	0.009	0.004	0.013	0.017	0.008	0.026	0.015	0.012	0.027	0.022	0.008	0.026
Friday	0	0.007	0.014	0.032	0.010	0.017	0.029	0.007	0.014	0.032	0.007	0.003	0.011	0.011	0.018	0.031	0.009	0.019	0.034	0.020	0.006	0.022
Friday	1	0.005	0.011	0.030	0.006	0.014	0.026	0.005	0.011	0.030	0.004	0.003	0.012	0.007	0.016	0.030	0.005	0.016	0.032	0.020	0.006	0.021
Friday	2	0.004	0.011	0.030	0.005	0.013	0.025	0.004	0.011	0.030	0.004	0.003	0.015	0.007	0.016	0.030	0.004	0.016	0.031	0.020	0.007	0.021
Friday	3	0.005	0.012	0.030	0.006	0.014	0.026	0.005	0.012	0.030	0.004	0.004	0.017	0.009	0.019	0.033	0.006	0.017	0.033	0.024	0.009	0.022
Friday	4	0.008	0.016	0.033	0.013	0.021	0.032	0.008	0.016	0.033	0.006	0.007	0.024	0.020	0.030	0.041	0.011	0.024	0.037	0.028	0.018	0.024
Friday	5	0.017	0.026	0.038	0.029	0.038	0.045	0.017	0.026	0.038	0.015	0.022	0.039	0.034	0.045	0.048	0.024	0.036	0.044	0.035	0.033	0.029
Friday	6	0.033	0.040	0.045	0.048	0.057	0.057	0.033	0.040	0.045	0.035	0.045	0.055	0.046	0.055	0.052	0.045	0.053	0.051	0.041	0.050	0.038
Friday	7	0.049	0.054	0.050	0.061	0.070	0.063	0.049	0.054	0.050	0.058	0.063	0.064	0.053	0.061	0.054	0.059	0.061	0.054	0.039	0.049	0.046
Friday	8	0.051	0.057	0.052	0.059	0.070	0.063	0.051	0.057	0.052	0.058	0.072	0.074	0.051	0.059	0.054	0.059	0.061	0.055	0.041	0.056	0.050
Friday	9	0.050	0.057	0.052	0.054	0.064	0.060	0.050	0.057	0.052	0.052	0.068	0.075	0.050	0.058	0.053	0.052	0.058	0.054	0.045	0.058	0.055
Friday	10	0.054	0.061	0.054	0.052	0.062	0.058	0.054	0.061	0.054	0.055	0.071	0.074	0.051	0.059	0.053	0.050	0.057	0.054	0.047	0.062	0.059
Friday	11	0.060	0.066	0.055	0.054	0.062	0.057	0.060	0.066	0.054	0.060	0.074	0.074	0.053	0.060	0.053	0.053	0.059	0.054	0.050	0.067	0.060
Friday	12	0.063	0.067	0.055	0.055	0.062	0.056	0.063	0.067	0.055	0.063	0.072	0.069	0.055	0.061	0.053	0.056	0.060	0.053	0.051	0.067	0.060
Friday	13	0.066	0.068	0.054	0.057	0.062	0.055	0.066	0.068	0.054	0.065	0.076	0.069	0.058	0.061	0.052	0.058	0.060	0.052	0.056	0.071	0.062
Friday	14	0.070	0.070	0.054	0.060	0.062	0.053	0.070	0.070	0.054	0.069	0.078	0.063	0.061	0.062	0.050	0.063	0.062	0.051	0.060	0.075	0.059
Friday	15	0.073	0.070	0.052	0.061	0.060	0.051	0.073	0.070	0.052	0.078	0.080	0.055	0.062	0.061	0.048	0.070	0.063	0.049	0.060	0.074	0.060
Friday	16	0.074	0.067	0.050	0.063	0.057	0.048	0.074	0.067	0.050	0.085	0.075	0.047	0.063	0.058	0.046	0.072	0.060	0.046	0.060	0.070	0.055
Friday	17	0.072	0.063	0.047	0.063	0.053	0.044	0.072	0.063	0.047	0.082	0.061	0.039	0.062	0.053	0.043	0.069	0.055	0.043	0.060	0.064	0.049
Friday	18	0.063	0.051	0.042	0.058	0.042	0.036	0.063	0.051	0.042	0.059	0.041	0.029	0.058	0.045	0.039	0.060	0.046	0.039	0.054	0.049	0.044
Friday	19	0.050	0.039	0.035	0.050	0.031	0.030	0.050	0.039	0.035	0.042	0.028	0.024	0.050	0.035	0.034	0.046	0.035	0.033	0.050	0.036	0.040
Friday	20	0.041	0.029	0.030	0.042	0.024	0.024	0.041	0.029	0.030	0.032	0.021	0.021	0.043	0.026	0.030	0.038	0.026	0.028	0.045	0.038	0.037
Friday	21	0.037	0.023	0.028	0.038	0.018	0.022	0.037	0.023	0.028	0.027	0.015	0.020	0.039	0.020	0.027	0.035	0.022	0.026	0.038	0.021	0.032
Friday	22	0.030	0.017	0.026	0.033	0.015	0.021	0.030	0.017	0.026	0.021	0.011	0.016	0.032	0.014	0.024	0.029	0.018	0.024	0.031	0.015	0.029
Friday	23	0.019	0.011	0.024	0.024	0.015	0.021	0.019	0.011	0.024	0.014	0.007	0.014	0.022	0.009	0.021	0.020	0.013	0.023	0.023	0.010	0.026
Saturday	0	0.013	0.019	0.038	0.017	0.030	0.049	0.013	0.019	0.038	0.012	0.007	0.021	0.017	0.027	0.047	0.016	0.027	0.046	0.023	0.011	0.030
Saturday	1	0.008	0.015	0.034	0.011	0.022	0.041	0.008	0.015	0.034	0.008	0.005	0.016	0.012	0.021	0.042	0.011	0.022	0.042	0.025	0.010	0.027
Saturday	2	0.006	0.014	0.032	0.009	0.019	0.037	0.006	0.014	0.032	0.006	0.004	0.020	0.010	0.019	0.040	0.008	0.020	0.039	0.025	0.009	0.026
Saturday	3	0.006	0.013	0.031	0.007	0.016	0.034	0.006	0.013	0.031	0.005	0.004	0.022	0.009	0.019	0.039	0.007	0.019	0.038	0.027	0.011	0.024
Saturday	4	0.007	0.014	0.032	0.009	0.018	0.036	0.007	0.014	0.032	0.006	0.008	0.024	0.012	0.021	0.041	0.009	0.022	0.039	0.031	0.020	0.025
Saturday	5	0.011	0.018	0.034	0.011	0.026	0.042	0.011	0.018	0.034	0.012	0.017	0.039	0.018	0.029	0.042	0.014	0.027	0.042	0.038	0.034	0.030
Saturday	6	0.019	0.026	0.039	0.026	0.037	0.050	0.019	0.026	0.039	0.021	0.028	0.049	0.028	0.039	0.050	0.023	0.035	0.046	0.038	0.047	0.040
Saturday	7	0.032	0.038	0.046	0.037	0.049	0.058	0.032	0.038	0.046	0.034	0.041	0.058	0.039	0.048	0.055	0.034	0.044	0.050	0.042	0.047	0.046

Day of Week	Hour	Nevada			Orange			Placer			Plumas			Riverside			Sacramento			San Benito		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Saturday	8	0.045	0.051	0.064	0.048	0.060	0.064	0.045	0.051	0.052	0.045	0.057	0.067	0.047	0.056	0.056	0.045	0.052	0.053	0.043	0.055	0.050
Saturday	9	0.057	0.062	0.056	0.055	0.065	0.064	0.057	0.062	0.056	0.054	0.068	0.074	0.054	0.062	0.057	0.054	0.059	0.055	0.047	0.062	0.055
Saturday	10	0.067	0.071	0.060	0.059	0.068	0.064	0.067	0.071	0.060	0.063	0.080	0.073	0.058	0.064	0.056	0.061	0.063	0.055	0.047	0.067	0.062
Saturday	11	0.074	0.076	0.061	0.062	0.069	0.062	0.074	0.076	0.061	0.068	0.082	0.071	0.062	0.066	0.054	0.066	0.065	0.055	0.049	0.068	0.063
Saturday	12	0.075	0.075	0.060	0.064	0.068	0.058	0.075	0.075	0.060	0.074	0.083	0.068	0.063	0.066	0.052	0.068	0.065	0.053	0.055	0.071	0.060
Saturday	13	0.075	0.074	0.057	0.064	0.064	0.053	0.075	0.074	0.057	0.074	0.079	0.062	0.064	0.064	0.050	0.068	0.064	0.051	0.054	0.070	0.059
Saturday	14	0.074	0.071	0.048	0.064	0.061	0.048	0.074	0.071	0.055	0.074	0.076	0.057	0.064	0.062	0.047	0.068	0.061	0.048	0.055	0.066	0.058
Saturday	15	0.072	0.068	0.051	0.064	0.057	0.044	0.072	0.068	0.051	0.073	0.074	0.052	0.064	0.059	0.044	0.067	0.059	0.045	0.055	0.065	0.056
Saturday	16	0.070	0.064	0.048	0.064	0.053	0.039	0.070	0.064	0.048	0.073	0.067	0.045	0.063	0.056	0.041	0.067	0.056	0.042	0.057	0.065	0.052
Saturday	17	0.066	0.057	0.044	0.062	0.048	0.034	0.066	0.057	0.044	0.069	0.058	0.039	0.061	0.051	0.034	0.064	0.052	0.039	0.056	0.053	0.047
Saturday	18	0.056	0.047	0.038	0.057	0.041	0.028	0.056	0.047	0.038	0.058	0.047	0.034	0.056	0.043	0.033	0.057	0.045	0.034	0.052	0.044	0.042
Saturday	19	0.046	0.037	0.033	0.050	0.032	0.022	0.046	0.037	0.033	0.046	0.036	0.029	0.049	0.035	0.028	0.048	0.037	0.030	0.049	0.039	0.039
Saturday	20	0.040	0.030	0.028	0.044	0.027	0.018	0.040	0.030	0.028	0.040	0.028	0.024	0.044	0.030	0.024	0.040	0.029	0.025	0.043	0.031	0.027
Saturday	21	0.035	0.025	0.025	0.042	0.026	0.018	0.035	0.025	0.025	0.036	0.022	0.023	0.042	0.026	0.022	0.040	0.029	0.025	0.038	0.025	0.029
Saturday	22	0.028	0.019	0.023	0.040	0.025	0.018	0.028	0.019	0.023	0.029	0.016	0.017	0.037	0.022	0.020	0.036	0.026	0.024	0.030	0.017	0.026
Saturday	23	0.020	0.014	0.021	0.030	0.021	0.019	0.020	0.014	0.021	0.020	0.011	0.017	0.029	0.017	0.018	0.026	0.020	0.022	0.023	0.011	0.020
Holiday	0	0.010	0.016	0.028	0.015	0.023	0.030	0.010	0.016	0.028	0.010	0.004	0.012	0.015	0.023	0.032	0.013	0.023	0.024	0.024	0.008	0.016
Holiday	1	0.006	0.013	0.027	0.009	0.018	0.027	0.006	0.013	0.027	0.006	0.004	0.011	0.010	0.018	0.030	0.008	0.019	0.030	0.022	0.009	0.015
Holiday	2	0.004	0.012	0.026	0.007	0.015	0.025	0.004	0.012	0.026	0.004	0.003	0.012	0.008	0.018	0.029	0.006	0.018	0.030	0.024	0.007	0.015
Holiday	3	0.005	0.013	0.027	0.006	0.015	0.025	0.005	0.013	0.027	0.004	0.005	0.015	0.009	0.020	0.031	0.006	0.019	0.030	0.024	0.009	0.017
Holiday	4	0.008	0.016	0.029	0.010	0.019	0.029	0.008	0.016	0.029	0.007	0.009	0.024	0.016	0.027	0.035	0.010	0.023	0.033	0.031	0.019	0.019
Holiday	5	0.014	0.023	0.032	0.023	0.032	0.038	0.014	0.023	0.032	0.014	0.020	0.037	0.026	0.036	0.041	0.019	0.032	0.037	0.033	0.029	0.024
Holiday	6	0.025	0.033	0.036	0.038	0.047	0.047	0.025	0.033	0.036	0.030	0.036	0.047	0.035	0.044	0.044	0.031	0.041	0.043	0.038	0.042	0.030
Holiday	7	0.036	0.044	0.042	0.047	0.057	0.053	0.036	0.044	0.042	0.044	0.052	0.061	0.041	0.049	0.046	0.042	0.049	0.046	0.040	0.044	0.037
Holiday	8	0.046	0.053	0.048	0.047	0.058	0.053	0.046	0.053	0.048	0.052	0.066	0.075	0.046	0.054	0.049	0.048	0.054	0.049	0.037	0.050	0.041
Holiday	9	0.054	0.059	0.050	0.050	0.060	0.054	0.054	0.059	0.050	0.053	0.071	0.081	0.051	0.057	0.050	0.052	0.057	0.051	0.046	0.057	0.048
Holiday	10	0.065	0.069	0.053	0.055	0.064	0.056	0.065	0.069	0.053	0.059	0.076	0.081	0.056	0.061	0.051	0.057	0.060	0.052	0.048	0.066	0.056
Holiday	11	0.074	0.074	0.057	0.059	0.067	0.058	0.074	0.074	0.057	0.066	0.076	0.071	0.061	0.065	0.053	0.063	0.065	0.054	0.055	0.077	0.063
Holiday	12	0.077	0.074	0.056	0.061	0.068	0.057	0.077	0.074	0.056	0.071	0.078	0.074	0.063	0.066	0.053	0.067	0.065	0.054	0.052	0.074	0.065
Holiday	13	0.076	0.074	0.058	0.062	0.067	0.057	0.076	0.074	0.058	0.071	0.076	0.065	0.064	0.066	0.053	0.068	0.066	0.055	0.055	0.071	0.069
Holiday	14	0.075	0.073	0.056	0.064	0.066	0.055	0.075	0.073	0.056	0.070	0.078	0.060	0.064	0.064	0.052	0.069	0.065	0.053	0.050	0.071	0.067
Holiday	15	0.074	0.070	0.055	0.065	0.062	0.052	0.074	0.070	0.055	0.075	0.075	0.053	0.064	0.061	0.050	0.070	0.063	0.052	0.061	0.068	0.068
Holiday	16	0.072	0.066	0.054	0.064	0.057	0.049	0.072	0.066	0.054	0.079	0.070	0.044	0.064	0.058	0.048	0.069	0.060	0.049	0.062	0.069	0.058
Holiday	17	0.068	0.059	0.051	0.064	0.051	0.045	0.068	0.059	0.051	0.074	0.064	0.041	0.064	0.053	0.045	0.066	0.054	0.046	0.058	0.062	0.058
Holiday	18	0.057	0.049	0.045	0.058	0.042	0.040	0.057	0.049	0.045	0.058	0.044	0.034	0.059	0.046	0.043	0.058	0.046	0.042	0.054	0.050	0.049
Holiday	19	0.047	0.036	0.041	0.052	0.032	0.034	0.047	0.036	0.041	0.047	0.033	0.026	0.052	0.036	0.038	0.049	0.036	0.037	0.049	0.037	0.047
Holiday	20	0.039	0.029	0.037	0.046	0.025	0.030	0.039	0.029	0.037	0.038	0.025	0.025	0.045	0.029	0.036	0.043	0.030	0.034	0.046	0.032	0.043
Holiday	21	0.030	0.020	0.033	0.041	0.021	0.029	0.030	0.020	0.033	0.030	0.018	0.021	0.039	0.022	0.032	0.037	0.024	0.031	0.040	0.025	0.038
Holiday	22	0.023	0.015	0.031	0.035	0.018	0.029	0.023	0.015	0.031	0.024	0.011	0.017	0.029	0.016	0.030	0.029	0.019	0.029	0.031	0.016	0.032
Holiday	23	0.015	0.010	0.029	0.023	0.014	0.030	0.015	0.010	0.029	0.014	0.007	0.014	0.021	0.011	0.028	0.020	0.014	0.029	0.020	0.008	0.028

Day of Week	Hour	San Bernardino			San Diego			San Francisco			San Joaquin			San Luis Obispo			San Mateo			Santa Barbara		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Sunday	0	0.024	0.030	0.035	0.019	0.033	0.051	0.026	0.032	0.056	0.016	0.024	0.039	0.017	0.009	0.017	0.021	0.029	0.049	0.020	0.017	0.032
Sunday	1	0.017	0.025	0.031	0.012	0.029	0.044	0.019	0.030	0.050	0.010	0.017	0.034	0.017	0.006	0.012	0.013	0.029	0.047	0.021	0.015	0.026
Sunday	2	0.014	0.022	0.028	0.009	0.026	0.040	0.017	0.030	0.048	0.007	0.015	0.031	0.018	0.005	0.009	0.010	0.028	0.045	0.020	0.012	0.022
Sunday	3	0.011	0.020	0.027	0.007	0.023	0.036	0.011	0.028	0.042	0.006	0.014	0.030	0.019	0.004	0.011	0.006	0.029	0.043	0.019	0.010	0.022
Sunday	4	0.012	0.020	0.027	0.007	0.023	0.034	0.009	0.028	0.040	0.008	0.015	0.030	0.019	0.005	0.009	0.007	0.029	0.041	0.023	0.014	0.023
Sunday	5	0.015	0.022	0.028	0.011	0.026	0.035	0.011	0.029	0.039	0.011	0.018	0.031	0.022	0.009	0.012	0.010	0.030	0.040	0.023	0.017	0.029
Sunday	6	0.021	0.026	0.030	0.018	0.030	0.037	0.018	0.032	0.040	0.017	0.022	0.033	0.026	0.015	0.017	0.015	0.031	0.039	0.029	0.024	0.031
Sunday	7	0.027	0.031	0.033	0.026	0.035	0.040	0.024	0.032	0.039	0.023	0.027	0.036	0.030	0.024	0.025	0.022	0.033	0.038	0.031	0.029	0.031
Sunday	8	0.036	0.038	0.037	0.037	0.041	0.043	0.033	0.036	0.040	0.032	0.036	0.040	0.037	0.037	0.039	0.032	0.036	0.038	0.036	0.042	0.037
Sunday	9	0.046	0.046	0.041	0.050	0.048	0.047	0.047	0.047	0.043	0.045	0.048	0.046	0.043	0.056	0.050	0.047	0.040	0.039	0.042	0.054	0.047
Sunday	10	0.055	0.055	0.047	0.062	0.055	0.050	0.060	0.049	0.044	0.056	0.059	0.050	0.051	0.072	0.068	0.062	0.046	0.042	0.046	0.065	0.055
Sunday	11	0.060	0.060	0.050	0.068	0.059	0.050	0.065	0.053	0.045	0.063	0.067	0.054	0.054	0.079	0.080	0.069	0.052	0.042	0.049	0.072	0.059
Sunday	12	0.064	0.066	0.053	0.072	0.061	0.051	0.067	0.056	0.043	0.068	0.071	0.056	0.058	0.089	0.088	0.072	0.056	0.043	0.055	0.078	0.062
Sunday	13	0.067	0.067	0.054	0.072	0.062	0.049	0.067	0.056	0.041	0.071	0.074	0.055	0.059	0.085	0.081	0.073	0.057	0.042	0.057	0.074	0.057
Sunday	14	0.068	0.066	0.054	0.071	0.059	0.046	0.067	0.056	0.040	0.073	0.073	0.054	0.062	0.085	0.075	0.072	0.058	0.041	0.060	0.072	0.051
Sunday	15	0.066	0.063	0.053	0.071	0.057	0.043	0.066	0.056	0.039	0.073	0.071	0.053	0.065	0.081	0.066	0.070	0.059	0.041	0.061	0.070	0.051
Sunday	16	0.065	0.060	0.052	0.070	0.056	0.042	0.065	0.057	0.038	0.073	0.068	0.050	0.067	0.076	0.063	0.070	0.060	0.041	0.063	0.066	0.049
Sunday	17	0.063	0.056	0.051	0.067	0.053	0.040	0.063	0.057	0.038	0.072	0.063	0.049	0.065	0.070	0.064	0.068	0.060	0.043	0.064	0.059	0.049
Sunday	18	0.060	0.051	0.049	0.061	0.048	0.038	0.058	0.054	0.038	0.067	0.055	0.044	0.063	0.058	0.055	0.059	0.055	0.041	0.061	0.054	0.046
Sunday	19	0.056	0.045	0.047	0.054	0.043	0.036	0.053	0.048	0.037	0.061	0.047	0.041	0.057	0.046	0.044	0.052	0.049	0.040	0.059	0.046	0.043
Sunday	20	0.052	0.042	0.047	0.047	0.039	0.036	0.049	0.044	0.038	0.054	0.040	0.039	0.053	0.037	0.035	0.049	0.045	0.041	0.053	0.040	0.043
Sunday	21	0.044	0.036	0.045	0.039	0.035	0.036	0.045	0.038	0.039	0.044	0.031	0.036	0.045	0.026	0.032	0.045	0.039	0.041	0.045	0.031	0.046
Sunday	22	0.034	0.030	0.042	0.029	0.030	0.037	0.037	0.032	0.041	0.031	0.024	0.035	0.034	0.016	0.025	0.034	0.030	0.040	0.035	0.023	0.045
Sunday	23	0.023	0.025	0.041	0.019	0.027	0.040	0.025	0.025	0.044	0.019	0.019	0.036	0.023	0.010	0.024	0.021	0.022	0.040	0.026	0.017	0.042
Monday	0	0.015	0.017	0.023	0.009	0.018	0.023	0.012	0.020	0.031	0.010	0.010	0.022	0.018	0.004	0.008	0.009	0.016	0.025	0.016	0.005	0.012
Monday	1	0.011	0.015	0.022	0.005	0.017	0.022	0.007	0.021	0.030	0.006	0.010	0.021	0.017	0.003	0.008	0.004	0.018	0.026	0.015	0.004	0.014
Monday	2	0.010	0.015	0.022	0.004	0.017	0.023	0.005	0.021	0.031	0.006	0.010	0.022	0.018	0.003	0.010	0.003	0.019	0.028	0.016	0.005	0.016
Monday	3	0.014	0.018	0.024	0.005	0.018	0.024	0.005	0.022	0.035	0.011	0.015	0.025	0.020	0.006	0.014	0.003	0.020	0.029	0.018	0.007	0.019
Monday	4	0.025	0.028	0.029	0.012	0.022	0.028	0.010	0.025	0.035	0.029	0.028	0.033	0.024	0.011	0.019	0.007	0.022	0.031	0.020	0.013	0.028
Monday	5	0.041	0.044	0.038	0.031	0.034	0.037	0.023	0.031	0.040	0.043	0.043	0.042	0.031	0.027	0.029	0.020	0.026	0.034	0.028	0.025	0.038
Monday	6	0.052	0.053	0.044	0.055	0.050	0.047	0.045	0.040	0.046	0.053	0.052	0.048	0.040	0.048	0.041	0.044	0.035	0.041	0.037	0.048	0.045
Monday	7	0.061	0.065	0.052	0.068	0.066	0.057	0.064	0.057	0.055	0.061	0.059	0.053	0.046	0.065	0.053	0.071	0.058	0.057	0.048	0.071	0.046
Monday	8	0.056	0.056	0.047	0.063	0.062	0.058	0.064	0.064	0.057	0.055	0.057	0.053	0.049	0.066	0.057	0.071	0.070	0.064	0.054	0.083	0.052
Monday	9	0.051	0.051	0.045	0.055	0.056	0.054	0.059	0.054	0.054	0.051	0.056	0.055	0.051	0.069	0.064	0.065	0.059	0.058	0.054	0.078	0.055
Monday	10	0.050	0.050	0.045	0.051	0.055	0.054	0.055	0.053	0.054	0.051	0.058	0.056	0.051	0.070	0.073	0.057	0.052	0.053	0.053	0.069	0.060
Monday	11	0.052	0.052	0.046	0.052	0.056	0.055	0.053	0.053	0.055	0.052	0.060	0.058	0.054	0.070	0.074	0.052	0.050	0.051	0.056	0.072	0.066
Monday	12	0.054	0.054	0.049	0.054	0.058	0.057	0.053	0.054	0.053	0.054	0.061	0.058	0.055	0.070	0.070	0.051	0.053	0.051	0.060	0.073	0.069
Monday	13	0.055	0.057	0.051	0.056	0.059	0.057	0.053	0.056	0.053	0.056	0.063	0.057	0.058	0.071	0.070	0.052	0.055	0.051	0.062	0.072	0.064
Monday	14	0.059	0.062	0.055	0.063	0.062	0.057	0.059	0.059	0.052	0.063	0.068	0.058	0.064	0.076	0.067	0.056	0.059	0.052	0.065	0.075	0.062
Monday	15	0.064	0.065	0.058	0.072	0.065	0.057	0.063	0.061	0.051	0.069	0.072	0.059	0.068	0.083	0.061	0.063	0.065	0.055	0.070	0.077	0.060
Monday	16	0.063	0.066	0.060	0.075	0.065	0.057	0.065	0.061	0.049	0.072	0.071	0.056	0.068	0.079	0.053	0.070	0.070	0.057	0.067	0.067	0.052
Monday	17	0.064	0.065	0.060	0.073	0.062	0.055	0.067	0.068	0.049	0.070	0.065	0.052	0.064	0.065	0.047	0.074	0.077	0.059	0.058	0.046	0.041
Monday	18	0.054	0.050	0.052	0.058	0.046	0.044	0.062	0.056	0.042	0.055	0.045	0.041	0.051	0.041	0.040	0.067	0.059	0.048	0.050	0.034	0.037
Monday	19	0.042	0.035	0.043	0.041	0.033	0.034	0.050	0.039	0.033	0.041	0.031	0.033	0.041	0.031	0.026	0.036	0.035	0.035	0.045	0.025	0.035
Monday	20	0.035	0.028	0.038	0.033	0.026	0.029	0.039	0.030	0.027	0.033	0.023	0.028	0.037	0.018	0.030	0.037	0.029	0.028	0.036	0.017	0.033
Monday	21	0.031	0.023	0.036	0.028	0.022	0.025	0.036	0.024	0.024	0.027	0.017	0.026	0.031	0.014	0.027	0.033	0.022	0.024	0.030	0.013	0.034
Monday	22	0.025	0.018	0.033	0.020	0.017	0.023	0.031	0.018	0.023	0.021	0.013	0.023	0.024	0.009	0.026	0.025	0.016	0.022	0.024	0.024	0.032
Monday	23	0.018	0.013	0.030	0.013	0.015	0.022	0.021	0.013	0.025	0.014	0.010	0.022	0.017	0.005	0.021	0.015	0.011	0.020	0.016	0.007	0.030
Tues/Wed/Thurs	0	0.013	0.016	0.024	0.007	0.017	0.025	0.012	0.019	0.032	0.009	0.011	0.024	0.016	0.004	0.017	0.008	0.016	0.026	0.016	0.005	0.022
Tues/Wed/Thurs	1	0.010	0.014	0.023	0.004	0.016	0.024	0.007	0.019	0.031	0.006	0.010	0.023	0.016	0.003	0.014	0.003	0.017	0.027	0.015	0.004	0.022
Tues/Wed/Thurs	2	0.010	0.015	0.024	0.003	0.016	0.024	0.005	0.020	0.032	0.005	0.010	0.023	0.016	0.003	0.014	0.002	0.018	0.028	0.015	0.004	0.021
Tues/Wed/Thurs	3	0.013	0.018	0.018	0.004	0.017	0.026	0.005	0.021	0.032	0.010	0.014	0.026	0.018	0.004	0.017	0.003	0.019	0.029	0.017	0.006	0.024

R3

Day of Week	Hour	San Bernardino			San Diego			San Francisco			San Joaquin			San Luis Obispo			San Mateo			Santa Barbara		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Tues/Wed/Thurs	4	0.024	0.027	0.031	0.010	0.022	0.029	0.009	0.024	0.036	0.026	0.034	0.021	0.009	0.022	0.007	0.021	0.021	0.031	0.019	0.012	0.033
Tues/Wed/Thurs	5	0.041	0.044	0.040	0.029	0.033	0.038	0.024	0.029	0.041	0.043	0.042	0.030	0.023	0.032	0.020	0.020	0.025	0.034	0.026	0.025	0.045
Tues/Wed/Thurs	6	0.053	0.053	0.046	0.055	0.050	0.049	0.047	0.040	0.049	0.054	0.051	0.049	0.041	0.049	0.046	0.046	0.035	0.042	0.039	0.051	0.052
Tues/Wed/Thurs	7	0.062	0.065	0.054	0.063	0.067	0.059	0.065	0.058	0.057	0.062	0.059	0.048	0.066	0.057	0.073	0.059	0.073	0.058	0.051	0.072	0.052
Tues/Wed/Thurs	8	0.056	0.057	0.050	0.063	0.064	0.059	0.064	0.067	0.059	0.056	0.057	0.049	0.067	0.060	0.071	0.072	0.072	0.066	0.056	0.083	0.056
Tues/Wed/Thurs	9	0.050	0.051	0.046	0.055	0.056	0.055	0.058	0.054	0.055	0.055	0.055	0.050	0.066	0.065	0.064	0.058	0.058	0.056	0.079	0.057	0.057
Tues/Wed/Thurs	10	0.049	0.049	0.046	0.050	0.054	0.054	0.053	0.051	0.053	0.049	0.056	0.052	0.064	0.067	0.054	0.050	0.052	0.054	0.054	0.070	0.060
Tues/Wed/Thurs	11	0.050	0.051	0.047	0.051	0.056	0.055	0.051	0.052	0.054	0.050	0.058	0.053	0.067	0.071	0.050	0.049	0.050	0.050	0.057	0.072	0.064
Tues/Wed/Thurs	12	0.052	0.053	0.049	0.053	0.058	0.056	0.051	0.053	0.053	0.052	0.059	0.056	0.069	0.067	0.049	0.051	0.050	0.060	0.071	0.072	0.062
Tues/Wed/Thurs	13	0.054	0.056	0.051	0.055	0.059	0.055	0.052	0.055	0.052	0.062	0.056	0.060	0.071	0.065	0.050	0.054	0.051	0.063	0.072	0.072	0.060
Tues/Wed/Thurs	14	0.058	0.062	0.054	0.063	0.062	0.056	0.058	0.059	0.051	0.062	0.068	0.063	0.076	0.064	0.056	0.058	0.052	0.064	0.075	0.075	0.058
Tues/Wed/Thurs	15	0.062	0.065	0.057	0.072	0.065	0.055	0.060	0.061	0.050	0.069	0.074	0.069	0.084	0.058	0.062	0.065	0.054	0.067	0.076	0.076	0.052
Tues/Wed/Thurs	16	0.064	0.067	0.059	0.074	0.065	0.055	0.064	0.062	0.047	0.072	0.074	0.057	0.070	0.081	0.050	0.070	0.056	0.064	0.065	0.065	0.044
Tues/Wed/Thurs	17	0.064	0.066	0.058	0.073	0.063	0.054	0.065	0.070	0.047	0.070	0.067	0.053	0.063	0.067	0.045	0.072	0.081	0.056	0.045	0.045	0.036
Tues/Wed/Thurs	18	0.055	0.052	0.050	0.061	0.047	0.043	0.062	0.059	0.041	0.056	0.048	0.041	0.053	0.044	0.039	0.067	0.065	0.050	0.036	0.036	0.034
Tues/Wed/Thurs	19	0.044	0.037	0.041	0.044	0.033	0.033	0.052	0.041	0.032	0.043	0.033	0.044	0.044	0.029	0.034	0.053	0.041	0.044	0.044	0.026	0.031
Tues/Wed/Thurs	20	0.038	0.029	0.037	0.036	0.026	0.028	0.042	0.031	0.026	0.034	0.025	0.028	0.038	0.021	0.028	0.039	0.029	0.037	0.019	0.029	0.029
Tues/Wed/Thurs	21	0.033	0.023	0.033	0.031	0.021	0.025	0.039	0.024	0.024	0.028	0.019	0.025	0.032	0.016	0.026	0.035	0.022	0.031	0.015	0.031	0.031
Tues/Wed/Thurs	22	0.027	0.017	0.029	0.022	0.017	0.022	0.034	0.018	0.023	0.021	0.014	0.021	0.025	0.010	0.023	0.027	0.015	0.025	0.011	0.027	0.027
Tues/Wed/Thurs	23	0.020	0.012	0.025	0.014	0.014	0.021	0.023	0.012	0.024	0.015	0.010	0.021	0.018	0.006	0.019	0.017	0.010	0.018	0.008	0.008	0.026
Friday	0	0.014	0.016	0.025	0.008	0.018	0.027	0.014	0.020	0.034	0.008	0.012	0.025	0.016	0.004	0.016	0.009	0.016	0.016	0.006	0.006	0.024
Friday	1	0.011	0.014	0.024	0.005	0.017	0.026	0.008	0.020	0.033	0.006	0.010	0.024	0.016	0.003	0.014	0.005	0.017	0.016	0.005	0.022	0.022
Friday	2	0.010	0.014	0.024	0.004	0.017	0.027	0.007	0.020	0.033	0.005	0.010	0.024	0.016	0.003	0.014	0.003	0.018	0.016	0.005	0.021	0.021
Friday	3	0.013	0.017	0.026	0.005	0.018	0.028	0.006	0.022	0.034	0.009	0.013	0.027	0.017	0.004	0.017	0.003	0.020	0.016	0.006	0.025	0.025
Friday	4	0.021	0.024	0.030	0.009	0.021	0.031	0.009	0.024	0.036	0.022	0.023	0.034	0.020	0.007	0.022	0.019	0.025	0.020	0.020	0.011	0.033
Friday	5	0.035	0.037	0.038	0.026	0.032	0.040	0.022	0.029	0.042	0.036	0.036	0.042	0.027	0.018	0.031	0.019	0.025	0.025	0.022	0.043	0.043
Friday	6	0.046	0.046	0.044	0.048	0.047	0.050	0.043	0.039	0.048	0.046	0.045	0.048	0.038	0.042	0.045	0.042	0.034	0.038	0.046	0.050	0.050
Friday	7	0.055	0.056	0.050	0.061	0.060	0.058	0.060	0.059	0.056	0.053	0.052	0.053	0.044	0.058	0.044	0.053	0.056	0.046	0.068	0.051	0.051
Friday	8	0.052	0.052	0.048	0.057	0.058	0.058	0.060	0.059	0.058	0.049	0.051	0.054	0.048	0.061	0.059	0.068	0.060	0.053	0.079	0.056	0.056
Friday	9	0.049	0.048	0.046	0.052	0.055	0.056	0.055	0.052	0.056	0.046	0.052	0.055	0.049	0.064	0.065	0.061	0.054	0.054	0.079	0.062	0.062
Friday	10	0.050	0.050	0.047	0.051	0.055	0.056	0.052	0.055	0.056	0.048	0.055	0.057	0.052	0.068	0.070	0.054	0.050	0.053	0.071	0.063	0.063
Friday	11	0.052	0.053	0.050	0.054	0.058	0.058	0.052	0.055	0.056	0.050	0.058	0.059	0.054	0.070	0.072	0.053	0.051	0.058	0.074	0.066	0.066
Friday	12	0.054	0.055	0.051	0.056	0.060	0.058	0.053	0.055	0.054	0.061	0.061	0.058	0.056	0.072	0.070	0.052	0.054	0.059	0.073	0.061	0.061
Friday	13	0.056	0.058	0.053	0.059	0.061	0.057	0.054	0.057	0.053	0.058	0.065	0.058	0.060	0.074	0.068	0.053	0.057	0.064	0.073	0.058	0.058
Friday	14	0.059	0.063	0.056	0.066	0.063	0.056	0.058	0.060	0.052	0.065	0.070	0.059	0.064	0.079	0.066	0.059	0.062	0.066	0.073	0.056	0.056
Friday	15	0.060	0.066	0.058	0.071	0.065	0.055	0.060	0.063	0.050	0.069	0.075	0.059	0.067	0.083	0.058	0.064	0.070	0.067	0.074	0.052	0.052
Friday	16	0.061	0.066	0.058	0.070	0.064	0.054	0.062	0.064	0.047	0.071	0.073	0.057	0.068	0.078	0.051	0.069	0.073	0.064	0.062	0.045	0.045
Friday	17	0.060	0.064	0.056	0.068	0.060	0.050	0.062	0.067	0.046	0.069	0.069	0.053	0.062	0.064	0.047	0.069	0.079	0.057	0.046	0.038	0.038
Friday	18	0.055	0.053	0.050	0.060	0.048	0.041	0.059	0.056	0.039	0.061	0.052	0.041	0.056	0.048	0.039	0.064	0.063	0.050	0.036	0.035	0.035
Friday	19	0.048	0.043	0.043	0.048	0.035	0.031	0.052	0.043	0.031	0.050	0.038	0.031	0.047	0.033	0.032	0.052	0.043	0.046	0.028	0.031	0.031
Friday	20	0.043	0.035	0.038	0.039	0.027	0.025	0.042	0.032	0.025	0.042	0.029	0.026	0.042	0.029	0.028	0.039	0.031	0.038	0.022	0.029	0.029
Friday	21	0.039	0.029	0.034	0.035	0.023	0.021	0.039	0.025	0.021	0.035	0.022	0.022	0.036	0.019	0.024	0.034	0.023	0.032	0.017	0.029	0.029
Friday	22	0.033	0.022	0.029	0.029	0.020	0.019	0.039	0.020	0.020	0.028	0.017	0.019	0.028	0.014	0.021	0.024	0.017	0.027	0.014	0.026	0.026
Friday	23	0.025	0.016	0.024	0.020	0.016	0.017	0.031	0.014	0.020	0.020	0.017	0.017	0.021	0.009	0.017	0.023	0.011	0.019	0.010	0.024	0.024
Saturday	0	0.020	0.024	0.034	0.015	0.026	0.043	0.022	0.026	0.048	0.014	0.021	0.037	0.018	0.007	0.027	0.017	0.024	0.022	0.013	0.039	0.039
Saturday	1	0.015	0.020	0.031	0.010	0.023	0.039	0.015	0.025	0.045	0.009	0.016	0.032	0.020	0.006	0.022	0.010	0.024	0.021	0.010	0.032	0.032
Saturday	2	0.013	0.019	0.029	0.007	0.022	0.037	0.013	0.025	0.043	0.007	0.014	0.031	0.020	0.006	0.020	0.008	0.024	0.022	0.009	0.030	0.030
Saturday	3	0.013	0.018	0.029	0.006	0.020	0.035	0.009	0.025	0.041	0.007	0.015	0.031	0.021	0.005	0.021	0.006	0.025	0.022	0.010	0.032	0.032
Saturday	4	0.015	0.020	0.030	0.007	0.022	0.036	0.008	0.026	0.039	0.011	0.018	0.033	0.022	0.007	0.023	0.007	0.026	0.024	0.014	0.040	0.040
Saturday	5	0.021	0.025	0.033	0.014	0.026	0.039	0.013	0.028	0.041	0.018	0.025	0.037	0.025	0.013	0.024	0.011	0.028	0.028	0.028	0.041	0.046
Saturday	6	0.030	0.032	0.038	0.024	0.032	0.045	0.021	0.031	0.044	0.027	0.033	0.042	0.032	0.024	0.039	0.019	0.031	0.035	0.035	0.053	0.053
Saturday	7	0.039	0.040	0.043	0.036	0.040	0.051	0.031	0.036	0.047	0.036	0.042	0.048	0.038	0.041	0.051	0.031	0.035	0.040	0.048	0.054	0.054

Day of Week	Hour	San Bernardino			San Diego			San Francisco			San Joaquin			San Luis Obispo			San Mateo			Santa Barbara		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Saturday	8	0.046	0.047	0.048	0.048	0.056	0.051	0.043	0.041	0.051	0.050	0.054	0.047	0.053	0.055	0.043	0.039	0.046	0.046	0.059	0.057	
Saturday	9	0.052	0.052	0.050	0.056	0.054	0.059	0.052	0.046	0.052	0.054	0.059	0.050	0.067	0.062	0.054	0.043	0.045	0.050	0.050	0.068	0.060
Saturday	10	0.056	0.056	0.053	0.062	0.058	0.060	0.059	0.051	0.053	0.061	0.067	0.054	0.078	0.069	0.062	0.050	0.050	0.053	0.070	0.059	0.059
Saturday	11	0.059	0.060	0.055	0.066	0.061	0.060	0.062	0.055	0.052	0.065	0.071	0.063	0.059	0.084	0.067	0.056	0.056	0.057	0.073	0.059	0.059
Saturday	12	0.061	0.063	0.057	0.068	0.063	0.058	0.063	0.057	0.051	0.067	0.072	0.062	0.060	0.082	0.070	0.068	0.059	0.059	0.074	0.056	0.056
Saturday	13	0.062	0.063	0.055	0.068	0.062	0.055	0.062	0.058	0.048	0.067	0.070	0.059	0.061	0.079	0.064	0.067	0.060	0.061	0.070	0.051	0.051
Saturday	14	0.062	0.063	0.055	0.068	0.061	0.051	0.062	0.059	0.046	0.067	0.068	0.052	0.060	0.074	0.061	0.067	0.061	0.061	0.068	0.048	0.048
Saturday	15	0.062	0.062	0.054	0.068	0.059	0.047	0.063	0.059	0.043	0.067	0.065	0.052	0.062	0.072	0.053	0.067	0.062	0.061	0.068	0.045	0.045
Saturday	16	0.061	0.060	0.052	0.067	0.057	0.043	0.063	0.059	0.042	0.066	0.061	0.048	0.061	0.066	0.050	0.067	0.062	0.059	0.059	0.041	0.041
Saturday	17	0.059	0.057	0.049	0.064	0.054	0.039	0.061	0.059	0.039	0.063	0.055	0.043	0.059	0.063	0.044	0.067	0.061	0.057	0.053	0.036	0.036
Saturday	18	0.055	0.051	0.044	0.057	0.047	0.033	0.058	0.056	0.036	0.057	0.045	0.036	0.053	0.050	0.037	0.061	0.055	0.052	0.046	0.033	0.033
Saturday	19	0.048	0.042	0.039	0.048	0.040	0.027	0.051	0.047	0.031	0.049	0.036	0.030	0.048	0.038	0.031	0.049	0.046	0.045	0.036	0.029	0.029
Saturday	20	0.043	0.037	0.035	0.042	0.035	0.023	0.044	0.040	0.028	0.043	0.030	0.026	0.043	0.032	0.029	0.042	0.039	0.041	0.031	0.024	0.024
Saturday	21	0.041	0.034	0.033	0.039	0.033	0.022	0.044	0.034	0.026	0.040	0.026	0.023	0.037	0.027	0.025	0.042	0.035	0.035	0.027	0.024	0.024
Saturday	22	0.037	0.029	0.030	0.034	0.031	0.021	0.045	0.032	0.027	0.035	0.023	0.021	0.028	0.018	0.021	0.040	0.030	0.029	0.023	0.023	0.023
Saturday	23	0.030	0.023	0.026	0.025	0.027	0.020	0.036	0.025	0.026	0.025	0.017	0.019	0.021	0.013	0.017	0.029	0.022	0.023	0.019	0.021	0.021
Holiday	0	0.018	0.020	0.026	0.013	0.023	0.029	0.021	0.023	0.035	0.012	0.015	0.027	0.018	0.006	0.012	0.014	0.020	0.020	0.010	0.020	0.020
Holiday	1	0.014	0.018	0.024	0.008	0.021	0.027	0.013	0.022	0.033	0.008	0.013	0.025	0.019	0.004	0.009	0.008	0.021	0.021	0.008	0.020	0.020
Holiday	2	0.012	0.017	0.024	0.006	0.020	0.027	0.010	0.024	0.033	0.006	0.012	0.025	0.019	0.003	0.011	0.005	0.022	0.031	0.019	0.006	0.018
Holiday	3	0.013	0.018	0.026	0.005	0.020	0.027	0.007	0.025	0.033	0.008	0.014	0.026	0.022	0.005	0.013	0.004	0.024	0.021	0.008	0.023	0.023
Holiday	4	0.019	0.024	0.029	0.008	0.023	0.030	0.008	0.028	0.035	0.015	0.020	0.030	0.022	0.008	0.015	0.006	0.025	0.022	0.012	0.028	0.028
Holiday	5	0.029	0.032	0.034	0.019	0.029	0.034	0.016	0.031	0.039	0.023	0.028	0.035	0.028	0.017	0.021	0.014	0.029	0.027	0.023	0.037	0.037
Holiday	6	0.036	0.038	0.037	0.035	0.040	0.042	0.028	0.036	0.044	0.031	0.035	0.039	0.034	0.030	0.031	0.027	0.035	0.031	0.034	0.042	0.042
Holiday	7	0.043	0.045	0.041	0.046	0.048	0.049	0.039	0.042	0.047	0.036	0.040	0.043	0.041	0.044	0.040	0.044	0.043	0.042	0.060	0.045	0.045
Holiday	8	0.047	0.048	0.043	0.048	0.050	0.050	0.046	0.049	0.050	0.041	0.045	0.047	0.046	0.055	0.046	0.053	0.048	0.048	0.073	0.051	0.051
Holiday	9	0.049	0.050	0.045	0.052	0.053	0.053	0.051	0.049	0.053	0.047	0.051	0.050	0.050	0.065	0.062	0.055	0.050	0.051	0.075	0.059	0.059
Holiday	10	0.053	0.053	0.047	0.057	0.058	0.056	0.057	0.054	0.054	0.055	0.061	0.056	0.052	0.076	0.072	0.058	0.052	0.053	0.071	0.058	0.058
Holiday	11	0.057	0.059	0.052	0.062	0.063	0.059	0.061	0.057	0.056	0.063	0.069	0.061	0.052	0.082	0.088	0.062	0.060	0.057	0.076	0.066	0.066
Holiday	12	0.060	0.063	0.053	0.065	0.065	0.060	0.063	0.059	0.055	0.066	0.072	0.062	0.058	0.086	0.085	0.062	0.060	0.059	0.079	0.070	0.070
Holiday	13	0.062	0.064	0.055	0.066	0.066	0.059	0.065	0.062	0.057	0.068	0.074	0.062	0.061	0.081	0.082	0.065	0.062	0.061	0.072	0.056	0.056
Holiday	14	0.063	0.066	0.056	0.068	0.065	0.058	0.067	0.063	0.055	0.070	0.073	0.060	0.059	0.076	0.075	0.067	0.066	0.060	0.073	0.060	0.060
Holiday	15	0.062	0.066	0.057	0.070	0.064	0.057	0.065	0.064	0.053	0.071	0.072	0.058	0.064	0.077	0.065	0.068	0.067	0.064	0.072	0.055	0.055
Holiday	16	0.062	0.063	0.057	0.069	0.060	0.053	0.063	0.062	0.048	0.071	0.068	0.054	0.068	0.072	0.057	0.069	0.067	0.060	0.061	0.050	0.050
Holiday	17	0.062	0.061	0.056	0.066	0.055	0.048	0.061	0.058	0.045	0.068	0.061	0.050	0.062	0.063	0.046	0.069	0.063	0.059	0.047	0.037	0.037
Holiday	18	0.056	0.053	0.052	0.058	0.045	0.042	0.057	0.052	0.040	0.060	0.050	0.042	0.053	0.044	0.039	0.060	0.053	0.053	0.038	0.036	0.036
Holiday	19	0.048	0.043	0.046	0.049	0.037	0.035	0.049	0.042	0.032	0.051	0.040	0.037	0.047	0.035	0.037	0.050	0.044	0.049	0.029	0.036	0.036
Holiday	20	0.043	0.034	0.041	0.043	0.030	0.030	0.044	0.034	0.029	0.044	0.031	0.032	0.041	0.027	0.028	0.045	0.033	0.040	0.024	0.032	0.032
Holiday	21	0.037	0.027	0.037	0.037	0.025	0.027	0.042	0.028	0.024	0.037	0.025	0.029	0.035	0.019	0.023	0.042	0.027	0.036	0.020	0.038	0.038
Holiday	22	0.031	0.021	0.033	0.030	0.022	0.025	0.040	0.021	0.025	0.029	0.019	0.026	0.027	0.014	0.022	0.033	0.020	0.028	0.017	0.034	0.034
Holiday	23	0.023	0.015	0.030	0.020	0.018	0.024	0.028	0.016	0.026	0.020	0.013	0.024	0.021	0.010	0.020	0.023	0.014	0.021	0.013	0.031	0.031

Day of Week	Hour	Santa Clara			Santa Cruz			Shasta			Sierra			Siskiyou			Solano			Sonoma		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Sunday	0	0.018	0.036	0.052	0.011	0.032	0.036	0.013	0.008	0.016	0.013	0.020	0.031	0.019	0.009	0.017	0.017	0.037	0.059	0.019	0.038	0.053
Sunday	1	0.011	0.034	0.046	0.006	0.031	0.036	0.013	0.006	0.013	0.008	0.016	0.028	0.021	0.007	0.014	0.011	0.032	0.052	0.012	0.034	0.047
Sunday	2	0.008	0.032	0.042	0.003	0.030	0.037	0.012	0.006	0.011	0.006	0.013	0.026	0.022	0.006	0.013	0.009	0.030	0.048	0.008	0.031	0.043
Sunday	3	0.005	0.032	0.039	0.002	0.034	0.035	0.012	0.005	0.011	0.005	0.012	0.025	0.022	0.005	0.013	0.007	0.027	0.044	0.006	0.030	0.040
Sunday	4	0.005	0.032	0.037	0.003	0.035	0.038	0.015	0.007	0.013	0.005	0.012	0.025	0.023	0.006	0.013	0.007	0.028	0.042	0.006	0.029	0.038
Sunday	5	0.008	0.033	0.036	0.006	0.035	0.035	0.018	0.012	0.018	0.008	0.015	0.027	0.025	0.008	0.016	0.010	0.029	0.042	0.010	0.031	0.038
Sunday	6	0.014	0.035	0.037	0.013	0.036	0.036	0.021	0.019	0.026	0.013	0.020	0.032	0.028	0.014	0.024	0.016	0.032	0.042	0.016	0.033	0.039
Sunday	7	0.021	0.037	0.039	0.022	0.038	0.039	0.029	0.030	0.039	0.022	0.028	0.034	0.030	0.022	0.034	0.021	0.035	0.043	0.023	0.036	0.040
Sunday	8	0.032	0.040	0.040	0.034	0.036	0.040	0.037	0.043	0.053	0.034	0.041	0.040	0.033	0.036	0.048	0.031	0.041	0.045	0.033	0.040	0.042
Sunday	9	0.047	0.046	0.044	0.051	0.043	0.043	0.043	0.055	0.067	0.048	0.055	0.046	0.036	0.048	0.062	0.059	0.053	0.046	0.048	0.046	0.044
Sunday	10	0.061	0.051	0.047	0.064	0.044	0.047	0.053	0.071	0.079	0.064	0.068	0.052	0.040	0.071	0.075	0.059	0.053	0.045	0.062	0.051	0.045
Sunday	11	0.068	0.053	0.047	0.071	0.047	0.046	0.060	0.077	0.080	0.075	0.075	0.055	0.044	0.082	0.086	0.067	0.055	0.044	0.067	0.053	0.046
Sunday	12	0.073	0.054	0.046	0.073	0.046	0.043	0.064	0.084	0.077	0.082	0.079	0.058	0.049	0.089	0.088	0.069	0.055	0.041	0.070	0.054	0.046
Sunday	13	0.075	0.055	0.045	0.076	0.047	0.041	0.066	0.083	0.070	0.084	0.079	0.058	0.054	0.090	0.080	0.070	0.055	0.038	0.073	0.055	0.050
Sunday	14	0.075	0.055	0.044	0.078	0.052	0.047	0.067	0.085	0.065	0.084	0.077	0.057	0.058	0.089	0.072	0.071	0.053	0.036	0.073	0.055	0.047
Sunday	15	0.075	0.053	0.041	0.081	0.054	0.051	0.072	0.083	0.061	0.082	0.073	0.057	0.063	0.087	0.069	0.071	0.052	0.035	0.072	0.052	0.041
Sunday	16	0.073	0.053	0.041	0.082	0.055	0.051	0.073	0.080	0.058	0.079	0.068	0.055	0.064	0.081	0.059	0.071	0.051	0.033	0.072	0.052	0.039
Sunday	17	0.071	0.051	0.040	0.080	0.058	0.052	0.068	0.066	0.056	0.072	0.062	0.053	0.065	0.066	0.051	0.070	0.051	0.033	0.070	0.050	0.038
Sunday	18	0.064	0.047	0.039	0.069	0.051	0.048	0.065	0.056	0.049	0.060	0.052	0.049	0.066	0.055	0.044	0.066	0.048	0.033	0.063	0.047	0.036
Sunday	19	0.057	0.044	0.038	0.058	0.051	0.047	0.058	0.043	0.041	0.050	0.043	0.045	0.062	0.043	0.036	0.060	0.046	0.034	0.056	0.044	0.035
Sunday	20	0.050	0.040	0.037	0.048	0.047	0.044	0.048	0.031	0.032	0.041	0.035	0.042	0.057	0.032	0.028	0.055	0.043	0.035	0.051	0.041	0.036
Sunday	21	0.041	0.034	0.038	0.036	0.039	0.039	0.041	0.023	0.026	0.031	0.026	0.039	0.049	0.022	0.023	0.045	0.039	0.039	0.042	0.038	0.037
Sunday	22	0.029	0.029	0.040	0.022	0.033	0.036	0.031	0.016	0.021	0.021	0.019	0.036	0.041	0.015	0.019	0.032	0.033	0.043	0.030	0.032	0.039
Sunday	23	0.018	0.024	0.044	0.011	0.028	0.032	0.020	0.012	0.017	0.013	0.015	0.033	0.028	0.012	0.016	0.020	0.028	0.049	0.019	0.027	0.043
Monday	0	0.007	0.022	0.028	0.004	0.024	0.033	0.013	0.006	0.012	0.008	0.014	0.027	0.023	0.006	0.013	0.010	0.026	0.035	0.007	0.023	0.029
Monday	1	0.003	0.022	0.027	0.001	0.025	0.031	0.012	0.006	0.011	0.005	0.012	0.025	0.025	0.007	0.011	0.005	0.024	0.034	0.003	0.022	0.028
Monday	2	0.002	0.023	0.028	0.001	0.025	0.034	0.013	0.006	0.011	0.004	0.012	0.025	0.025	0.007	0.011	0.005	0.024	0.034	0.002	0.022	0.029
Monday	3	0.003	0.025	0.030	0.002	0.025	0.034	0.015	0.010	0.012	0.006	0.014	0.027	0.027	0.010	0.012	0.006	0.026	0.035	0.003	0.023	0.030
Monday	4	0.007	0.029	0.033	0.007	0.031	0.038	0.019	0.019	0.015	0.011	0.019	0.030	0.030	0.015	0.012	0.015	0.032	0.040	0.012	0.028	0.035
Monday	5	0.024	0.035	0.040	0.026	0.034	0.038	0.025	0.030	0.021	0.023	0.030	0.036	0.033	0.022	0.018	0.037	0.043	0.046	0.033	0.041	0.042
Monday	6	0.047	0.046	0.049	0.061	0.043	0.049	0.032	0.041	0.024	0.042	0.047	0.043	0.036	0.034	0.024	0.050	0.051	0.050	0.054	0.051	0.048
Monday	7	0.065	0.054	0.057	0.082	0.053	0.056	0.034	0.048	0.032	0.060	0.061	0.048	0.040	0.043	0.030	0.061	0.058	0.053	0.066	0.058	0.053
Monday	8	0.068	0.057	0.060	0.079	0.054	0.059	0.039	0.059	0.039	0.059	0.062	0.050	0.043	0.054	0.039	0.056	0.057	0.055	0.062	0.060	0.055
Monday	9	0.065	0.055	0.055	0.073	0.053	0.053	0.047	0.065	0.046	0.056	0.061	0.050	0.045	0.067	0.048	0.054	0.056	0.055	0.055	0.056	0.054
Monday	10	0.056	0.053	0.054	0.064	0.050	0.052	0.050	0.070	0.053	0.058	0.064	0.051	0.050	0.074	0.054	0.055	0.058	0.056	0.052	0.054	0.053
Monday	11	0.052	0.054	0.054	0.059	0.055	0.054	0.056	0.072	0.055	0.062	0.066	0.053	0.052	0.075	0.059	0.056	0.057	0.055	0.053	0.055	0.054
Monday	12	0.053	0.055	0.054	0.055	0.060	0.059	0.059	0.073	0.055	0.066	0.068	0.054	0.055	0.078	0.059	0.057	0.058	0.054	0.054	0.056	0.054
Monday	13	0.054	0.056	0.053	0.056	0.054	0.052	0.060	0.076	0.058	0.067	0.067	0.054	0.057	0.081	0.060	0.058	0.057	0.052	0.056	0.056	0.054
Monday	14	0.062	0.060	0.054	0.059	0.061	0.057	0.065	0.079	0.059	0.070	0.069	0.055	0.057	0.081	0.065	0.064	0.057	0.051	0.063	0.059	0.056
Monday	15	0.068	0.063	0.055	0.063	0.060	0.051	0.071	0.081	0.062	0.073	0.069	0.055	0.059	0.080	0.063	0.069	0.056	0.048	0.069	0.063	0.058
Monday	16	0.071	0.063	0.054	0.067	0.059	0.051	0.070	0.070	0.063	0.075	0.067	0.054	0.060	0.072	0.064	0.071	0.054	0.044	0.072	0.060	0.052
Monday	17	0.074	0.062	0.052	0.069	0.058	0.047	0.065	0.057	0.066	0.073	0.061	0.052	0.057	0.059	0.066	0.070	0.050	0.040	0.073	0.056	0.047
Monday	18	0.065	0.050	0.042	0.057	0.051	0.040	0.058	0.042	0.064	0.056	0.046	0.045	0.053	0.045	0.063	0.054	0.041	0.035	0.061	0.045	0.039
Monday	19	0.052	0.037	0.031	0.040	0.042	0.034	0.054	0.031	0.059	0.040	0.031	0.039	0.048	0.032	0.060	0.042	0.032	0.028	0.045	0.033	0.031
Monday	20	0.036	0.028	0.025	0.028	0.030	0.025	0.050	0.022	0.054	0.031	0.022	0.035	0.042	0.022	0.054	0.035	0.026	0.025	0.035	0.026	0.026
Monday	21	0.030	0.022	0.022	0.023	0.024	0.020	0.041	0.017	0.051	0.025	0.017	0.032	0.036	0.016	0.046	0.029	0.022	0.023	0.031	0.022	0.024
Monday	22	0.022	0.016	0.020	0.015	0.018	0.017	0.030	0.011	0.043	0.017	0.012	0.030	0.029	0.012	0.039	0.023	0.018	0.023	0.023	0.017	0.023
Monday	23	0.014	0.012	0.022	0.009	0.013	0.017	0.022	0.008	0.034	0.012	0.009	0.030	0.020	0.008	0.031	0.016	0.016	0.028	0.014	0.014	0.025
Tues/Wed/Thurs	0	0.006	0.022	0.029	0.004	0.023	0.029	0.012	0.006	0.017	0.008	0.014	0.029	0.025	0.007	0.018	0.009	0.025	0.037	0.006	0.022	0.031
Tues/Wed/Thurs	1	0.003	0.022	0.029	0.001	0.024	0.032	0.012	0.005	0.015	0.008	0.011	0.027	0.023	0.006	0.015	0.005	0.023	0.036	0.003	0.021	0.030
Tues/Wed/Thurs	2	0.002	0.023	0.029	0.001	0.025	0.032	0.013	0.006	0.014	0.004	0.011	0.027	0.027	0.006	0.013	0.004	0.023	0.036	0.002	0.021	0.030
Tues/Wed/Thurs	3	0.003	0.025	0.031	0.001	0.027	0.034	0.014	0.009	0.015	0.005	0.013	0.029	0.029	0.009	0.013	0.005	0.025	0.037	0.003	0.023	0.031

Day of Week	Hour	Santa Clara			Santa Cruz			Shasta			Sierra			Siskiyou			Solano			Sonoma		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Tues/Wed/Thurs	4	0.007	0.028	0.034	0.006	0.029	0.036	0.018	0.017	0.017	0.010	0.018	0.031	0.032	0.014	0.016	0.013	0.030	0.041	0.011	0.028	0.036
Tues/Wed/Thurs	5	0.025	0.036	0.042	0.026	0.032	0.038	0.023	0.026	0.022	0.022	0.029	0.037	0.035	0.021	0.020	0.035	0.042	0.048	0.034	0.040	0.044
Tues/Wed/Thurs	6	0.050	0.047	0.052	0.065	0.040	0.045	0.030	0.042	0.030	0.042	0.047	0.044	0.038	0.033	0.027	0.050	0.050	0.052	0.056	0.052	0.049
Tues/Wed/Thurs	7	0.067	0.055	0.059	0.084	0.055	0.056	0.038	0.051	0.039	0.060	0.061	0.050	0.040	0.046	0.036	0.061	0.057	0.054	0.068	0.059	0.054
Tues/Wed/Thurs	8	0.069	0.058	0.061	0.080	0.055	0.055	0.042	0.061	0.048	0.060	0.060	0.051	0.042	0.056	0.046	0.056	0.056	0.055	0.063	0.060	0.056
Tues/Wed/Thurs	9	0.065	0.055	0.055	0.074	0.054	0.056	0.047	0.064	0.058	0.055	0.060	0.050	0.044	0.066	0.057	0.053	0.052	0.055	0.055	0.055	0.053
Tues/Wed/Thurs	10	0.051	0.053	0.054	0.062	0.052	0.053	0.051	0.067	0.066	0.056	0.061	0.051	0.045	0.071	0.065	0.052	0.057	0.054	0.051	0.053	0.052
Tues/Wed/Thurs	11	0.051	0.053	0.053	0.057	0.053	0.055	0.054	0.070	0.069	0.059	0.064	0.052	0.047	0.076	0.070	0.052	0.057	0.054	0.050	0.054	0.052
Tues/Wed/Thurs	12	0.051	0.055	0.053	0.054	0.057	0.055	0.058	0.072	0.067	0.061	0.065	0.053	0.050	0.076	0.070	0.054	0.057	0.053	0.052	0.055	0.053
Tues/Wed/Thurs	13	0.054	0.056	0.052	0.054	0.058	0.054	0.061	0.074	0.066	0.066	0.066	0.053	0.052	0.077	0.069	0.057	0.057	0.051	0.054	0.056	0.054
Tues/Wed/Thurs	14	0.061	0.059	0.052	0.058	0.061	0.056	0.065	0.077	0.063	0.068	0.068	0.053	0.057	0.081	0.067	0.064	0.058	0.049	0.062	0.059	0.054
Tues/Wed/Thurs	15	0.067	0.063	0.054	0.062	0.061	0.055	0.070	0.080	0.061	0.073	0.069	0.053	0.058	0.078	0.064	0.070	0.058	0.046	0.067	0.063	0.056
Tues/Wed/Thurs	16	0.070	0.064	0.053	0.065	0.060	0.053	0.072	0.072	0.058	0.075	0.067	0.052	0.057	0.072	0.061	0.073	0.056	0.043	0.070	0.060	0.051
Tues/Wed/Thurs	17	0.072	0.062	0.051	0.067	0.057	0.047	0.065	0.057	0.056	0.074	0.063	0.050	0.056	0.060	0.057	0.072	0.052	0.039	0.071	0.057	0.046
Tues/Wed/Thurs	18	0.065	0.052	0.042	0.058	0.050	0.043	0.060	0.044	0.052	0.059	0.048	0.044	0.053	0.046	0.053	0.058	0.043	0.033	0.062	0.047	0.039
Tues/Wed/Thurs	19	0.053	0.037	0.030	0.041	0.034	0.034	0.053	0.032	0.045	0.043	0.038	0.038	0.048	0.033	0.044	0.046	0.034	0.028	0.048	0.035	0.031
Tues/Wed/Thurs	20	0.038	0.027	0.024	0.029	0.032	0.028	0.047	0.024	0.039	0.035	0.025	0.034	0.045	0.025	0.038	0.038	0.028	0.024	0.038	0.027	0.026
Tues/Wed/Thurs	21	0.032	0.021	0.021	0.024	0.024	0.021	0.042	0.021	0.034	0.029	0.019	0.031	0.038	0.018	0.032	0.032	0.023	0.022	0.033	0.022	0.024
Tues/Wed/Thurs	22	0.023	0.016	0.019	0.017	0.018	0.019	0.031	0.013	0.028	0.020	0.013	0.028	0.029	0.032	0.014	0.026	0.025	0.018	0.024	0.015	0.022
Tues/Wed/Thurs	23	0.014	0.011	0.020	0.009	0.012	0.015	0.022	0.010	0.022	0.013	0.009	0.028	0.025	0.010	0.021	0.016	0.015	0.028	0.015	0.013	0.024
Friday	0	0.007	0.022	0.032	0.005	0.023	0.030	0.013	0.007	0.021	0.007	0.014	0.032	0.021	0.007	0.019	0.009	0.025	0.040	0.008	0.022	0.033
Friday	1	0.004	0.023	0.031	0.002	0.022	0.031	0.012	0.005	0.018	0.005	0.011	0.030	0.023	0.006	0.017	0.006	0.024	0.039	0.004	0.021	0.031
Friday	2	0.003	0.024	0.032	0.001	0.024	0.032	0.012	0.006	0.018	0.004	0.011	0.030	0.024	0.007	0.016	0.005	0.024	0.039	0.003	0.022	0.032
Friday	3	0.003	0.025	0.033	0.002	0.027	0.034	0.014	0.008	0.018	0.005	0.012	0.030	0.026	0.009	0.016	0.005	0.025	0.040	0.004	0.023	0.033
Friday	4	0.007	0.029	0.036	0.005	0.030	0.038	0.016	0.015	0.021	0.008	0.016	0.033	0.032	0.013	0.019	0.011	0.030	0.044	0.010	0.028	0.036
Friday	5	0.022	0.035	0.044	0.022	0.033	0.041	0.023	0.023	0.026	0.017	0.026	0.038	0.040	0.033	0.030	0.032	0.027	0.040	0.030	0.039	0.044
Friday	6	0.044	0.045	0.053	0.054	0.040	0.046	0.029	0.035	0.033	0.033	0.040	0.045	0.045	0.037	0.039	0.050	0.053	0.056	0.050	0.049	0.050
Friday	7	0.060	0.052	0.058	0.075	0.049	0.055	0.034	0.044	0.041	0.049	0.054	0.050	0.040	0.051	0.049	0.048	0.054	0.057	0.063	0.057	0.056
Friday	8	0.063	0.054	0.060	0.071	0.047	0.050	0.039	0.055	0.049	0.051	0.057	0.052	0.040	0.051	0.049	0.050	0.053	0.056	0.059	0.057	0.055
Friday	9	0.060	0.054	0.057	0.068	0.049	0.051	0.042	0.060	0.055	0.050	0.057	0.052	0.045	0.063	0.054	0.048	0.055	0.057	0.053	0.054	0.054
Friday	10	0.054	0.053	0.056	0.061	0.051	0.053	0.049	0.063	0.058	0.054	0.061	0.054	0.048	0.069	0.060	0.052	0.056	0.056	0.051	0.053	0.053
Friday	11	0.053	0.055	0.056	0.061	0.056	0.054	0.052	0.069	0.061	0.060	0.066	0.055	0.049	0.072	0.063	0.056	0.058	0.055	0.053	0.055	0.054
Friday	12	0.055	0.057	0.056	0.058	0.056	0.053	0.057	0.070	0.061	0.063	0.067	0.055	0.052	0.074	0.063	0.059	0.058	0.053	0.056	0.057	0.055
Friday	13	0.058	0.058	0.054	0.060	0.059	0.058	0.057	0.075	0.061	0.066	0.068	0.054	0.054	0.077	0.062	0.063	0.058	0.051	0.058	0.058	0.056
Friday	14	0.064	0.061	0.053	0.064	0.062	0.056	0.065	0.080	0.060	0.070	0.070	0.054	0.059	0.080	0.063	0.067	0.058	0.048	0.064	0.059	0.056
Friday	15	0.067	0.063	0.054	0.065	0.061	0.055	0.070	0.082	0.059	0.073	0.070	0.052	0.063	0.081	0.061	0.069	0.057	0.045	0.066	0.062	0.056
Friday	16	0.069	0.062	0.051	0.065	0.062	0.054	0.072	0.073	0.057	0.074	0.067	0.050	0.058	0.075	0.059	0.070	0.054	0.041	0.067	0.059	0.050
Friday	17	0.069	0.060	0.048	0.064	0.059	0.049	0.065	0.062	0.055	0.072	0.063	0.047	0.059	0.063	0.055	0.067	0.050	0.037	0.067	0.055	0.046
Friday	18	0.063	0.049	0.038	0.056	0.053	0.046	0.061	0.047	0.051	0.063	0.051	0.042	0.054	0.052	0.051	0.061	0.044	0.031	0.060	0.047	0.039
Friday	19	0.053	0.037	0.028	0.044	0.043	0.035	0.059	0.039	0.046	0.050	0.039	0.035	0.050	0.036	0.046	0.054	0.037	0.026	0.049	0.036	0.030
Friday	20	0.039	0.028	0.021	0.032	0.034	0.027	0.051	0.028	0.040	0.041	0.029	0.030	0.046	0.030	0.041	0.047	0.031	0.022	0.040	0.029	0.023
Friday	21	0.033	0.022	0.018	0.027	0.022	0.022	0.045	0.022	0.035	0.037	0.023	0.028	0.040	0.022	0.036	0.039	0.025	0.020	0.035	0.023	0.020
Friday	22	0.028	0.017	0.016	0.023	0.019	0.016	0.037	0.018	0.031	0.030	0.017	0.026	0.031	0.016	0.031	0.030	0.020	0.020	0.030	0.019	0.019
Friday	23	0.021	0.013	0.016	0.015	0.014	0.013	0.026	0.012	0.025	0.019	0.011	0.024	0.025	0.012	0.025	0.021	0.016	0.022	0.022	0.015	0.020
Saturday	0	0.015	0.029	0.046	0.009	0.028	0.038	0.015	0.011	0.021	0.013	0.019	0.038	0.026	0.013	0.020	0.014	0.031	0.057	0.015	0.030	0.044
Saturday	1	0.009	0.028	0.042	0.005	0.028	0.038	0.014	0.008	0.018	0.008	0.015	0.034	0.026	0.008	0.016	0.009	0.028	0.052	0.009	0.027	0.040
Saturday	2	0.007	0.028	0.040	0.003	0.029	0.042	0.014	0.008	0.018	0.006	0.014	0.032	0.027	0.007	0.015	0.007	0.049	0.026	0.006	0.026	0.039

Day of Week	Hour	Santa Clara			Santa Cruz			Shasta			Sierra			Siskiyou			Solano			Sonoma		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Saturday	8	0.045	0.046	0.053	0.049	0.041	0.046	0.040	0.055	0.051	0.045	0.051	0.052	0.041	0.047	0.047	0.044	0.049	0.056	0.046	0.047	0.049
Saturday	9	0.055	0.051	0.056	0.059	0.046	0.046	0.044	0.064	0.061	0.057	0.062	0.056	0.045	0.063	0.059	0.056	0.056	0.054	0.055	0.055	0.051
Saturday	10	0.062	0.054	0.056	0.066	0.047	0.047	0.051	0.071	0.067	0.067	0.071	0.060	0.049	0.073	0.067	0.065	0.057	0.052	0.061	0.054	0.051
Saturday	11	0.067	0.057	0.056	0.068	0.052	0.052	0.058	0.077	0.068	0.074	0.076	0.061	0.050	0.084	0.073	0.068	0.058	0.050	0.065	0.056	0.052
Saturday	12	0.069	0.057	0.054	0.067	0.053	0.050	0.060	0.076	0.067	0.075	0.075	0.060	0.053	0.083	0.071	0.067	0.057	0.047	0.066	0.058	0.055
Saturday	13	0.069	0.057	0.051	0.067	0.055	0.049	0.059	0.073	0.066	0.075	0.074	0.057	0.055	0.081	0.069	0.066	0.056	0.044	0.067	0.059	0.058
Saturday	14	0.069	0.057	0.049	0.069	0.053	0.049	0.065	0.076	0.066	0.074	0.071	0.055	0.057	0.074	0.065	0.066	0.055	0.041	0.067	0.058	0.057
Saturday	15	0.069	0.057	0.045	0.072	0.056	0.049	0.067	0.073	0.064	0.072	0.068	0.051	0.060	0.074	0.062	0.066	0.054	0.038	0.068	0.057	0.051
Saturday	16	0.068	0.055	0.043	0.074	0.055	0.048	0.065	0.069	0.059	0.070	0.064	0.048	0.056	0.070	0.058	0.066	0.053	0.034	0.068	0.056	0.047
Saturday	17	0.067	0.052	0.038	0.074	0.055	0.046	0.064	0.062	0.055	0.066	0.057	0.044	0.055	0.061	0.051	0.055	0.050	0.031	0.067	0.054	0.044
Saturday	18	0.061	0.047	0.034	0.066	0.052	0.040	0.061	0.048	0.050	0.056	0.047	0.038	0.051	0.049	0.052	0.058	0.046	0.029	0.060	0.048	0.036
Saturday	19	0.050	0.040	0.029	0.054	0.045	0.035	0.059	0.041	0.044	0.046	0.037	0.033	0.049	0.038	0.045	0.050	0.040	0.026	0.049	0.041	0.029
Saturday	20	0.042	0.035	0.025	0.044	0.041	0.033	0.050	0.031	0.036	0.040	0.030	0.028	0.042	0.031	0.038	0.045	0.036	0.023	0.043	0.036	0.025
Saturday	21	0.040	0.031	0.023	0.039	0.037	0.032	0.044	0.023	0.030	0.035	0.025	0.025	0.037	0.023	0.031	0.041	0.033	0.023	0.041	0.033	0.024
Saturday	22	0.036	0.027	0.023	0.032	0.031	0.028	0.034	0.017	0.024	0.028	0.019	0.023	0.031	0.017	0.026	0.035	0.029	0.023	0.037	0.029	0.023
Saturday	23	0.026	0.022	0.022	0.020	0.025	0.025	0.026	0.013	0.019	0.020	0.014	0.021	0.024	0.012	0.019	0.026	0.023	0.023	0.028	0.024	0.022
Holiday	0	0.012	0.025	0.032	0.008	0.024	0.031	0.014	0.008	0.015	0.010	0.016	0.028	0.024	0.008	0.015	0.013	0.029	0.038	0.013	0.027	0.034
Holiday	1	0.007	0.025	0.031	0.003	0.025	0.034	0.013	0.007	0.013	0.006	0.013	0.027	0.027	0.008	0.012	0.008	0.027	0.038	0.007	0.026	0.033
Holiday	2	0.004	0.026	0.032	0.002	0.025	0.034	0.013	0.006	0.012	0.004	0.012	0.026	0.024	0.008	0.012	0.005	0.025	0.037	0.004	0.025	0.033
Holiday	3	0.003	0.027	0.032	0.001	0.024	0.029	0.013	0.006	0.012	0.005	0.013	0.027	0.029	0.010	0.013	0.005	0.026	0.037	0.003	0.025	0.033
Holiday	4	0.005	0.029	0.034	0.004	0.030	0.034	0.016	0.013	0.014	0.008	0.016	0.029	0.029	0.012	0.014	0.008	0.028	0.039	0.007	0.029	0.035
Holiday	5	0.014	0.034	0.038	0.012	0.033	0.041	0.020	0.017	0.020	0.014	0.023	0.032	0.031	0.016	0.017	0.018	0.034	0.043	0.017	0.034	0.039
Holiday	6	0.027	0.039	0.044	0.028	0.037	0.045	0.025	0.028	0.026	0.025	0.033	0.036	0.037	0.025	0.023	0.025	0.040	0.046	0.029	0.040	0.044
Holiday	7	0.039	0.043	0.048	0.043	0.035	0.038	0.030	0.037	0.036	0.036	0.044	0.042	0.038	0.033	0.031	0.032	0.045	0.050	0.038	0.045	0.047
Holiday	8	0.050	0.048	0.052	0.052	0.048	0.053	0.036	0.051	0.046	0.046	0.053	0.048	0.040	0.049	0.040	0.041	0.050	0.053	0.045	0.050	0.051
Holiday	9	0.054	0.052	0.054	0.058	0.051	0.053	0.047	0.068	0.056	0.054	0.059	0.050	0.043	0.062	0.054	0.051	0.055	0.055	0.049	0.053	0.052
Holiday	10	0.058	0.055	0.056	0.064	0.049	0.054	0.051	0.068	0.064	0.065	0.069	0.053	0.050	0.076	0.060	0.062	0.060	0.055	0.056	0.056	0.053
Holiday	11	0.061	0.058	0.057	0.069	0.055	0.050	0.059	0.083	0.069	0.074	0.074	0.057	0.047	0.084	0.068	0.068	0.056	0.062	0.062	0.059	0.055
Holiday	12	0.063	0.060	0.057	0.067	0.057	0.059	0.066	0.081	0.071	0.077	0.074	0.056	0.053	0.083	0.070	0.070	0.061	0.054	0.067	0.061	0.056
Holiday	13	0.066	0.062	0.057	0.068	0.069	0.064	0.062	0.084	0.068	0.076	0.074	0.058	0.062	0.091	0.067	0.071	0.062	0.052	0.070	0.062	0.056
Holiday	14	0.069	0.062	0.056	0.073	0.058	0.060	0.069	0.076	0.064	0.075	0.073	0.056	0.059	0.087	0.069	0.072	0.060	0.051	0.073	0.062	0.057
Holiday	15	0.071	0.062	0.054	0.072	0.070	0.056	0.065	0.081	0.061	0.074	0.070	0.055	0.057	0.079	0.065	0.068	0.056	0.046	0.071	0.061	0.054
Holiday	16	0.072	0.060	0.051	0.071	0.059	0.052	0.070	0.068	0.061	0.072	0.066	0.054	0.056	0.072	0.062	0.066	0.054	0.044	0.070	0.057	0.050
Holiday	17	0.071	0.057	0.047	0.070	0.058	0.048	0.068	0.063	0.060	0.068	0.059	0.051	0.056	0.058	0.060	0.064	0.050	0.040	0.067	0.053	0.044
Holiday	18	0.064	0.048	0.039	0.063	0.054	0.045	0.063	0.047	0.055	0.057	0.049	0.045	0.053	0.044	0.058	0.058	0.042	0.034	0.059	0.045	0.038
Holiday	19	0.054	0.038	0.032	0.052	0.035	0.029	0.056	0.035	0.048	0.047	0.036	0.041	0.048	0.029	0.049	0.051	0.037	0.029	0.051	0.036	0.031
Holiday	20	0.045	0.031	0.026	0.043	0.035	0.027	0.050	0.028	0.041	0.039	0.029	0.037	0.044	0.024	0.045	0.047	0.031	0.025	0.046	0.031	0.028
Holiday	21	0.039	0.025	0.024	0.036	0.029	0.026	0.045	0.021	0.035	0.030	0.020	0.033	0.040	0.019	0.040	0.042	0.026	0.024	0.041	0.026	0.026
Holiday	22	0.031	0.019	0.022	0.024	0.021	0.022	0.027	0.013	0.029	0.023	0.015	0.031	0.031	0.014	0.030	0.033	0.022	0.025	0.033	0.021	0.025
Holiday	23	0.020	0.014	0.024	0.015	0.016	0.015	0.022	0.010	0.023	0.015	0.010	0.029	0.024	0.009	0.024	0.022	0.018	0.029	0.021	0.017	0.026



Day of Week	Hour	Stamislauis		Sutter		Tehama		Trinity		Tulare		Tuolumne		Ventura	
		LM	HH	LM	HH	LM	HH	LM	HH	LM	HH	LM	HH	LM	HH
Sunday	0	0.014	0.025	0.037	0.031	0.013	0.008	0.016	0.009	0.017	0.022	0.015	0.017	0.022	0.032
Sunday	1	0.009	0.019	0.032	0.028	0.013	0.006	0.013	0.021	0.014	0.024	0.015	0.017	0.024	0.009
Sunday	2	0.007	0.016	0.029	0.026	0.012	0.006	0.013	0.022	0.013	0.023	0.011	0.008	0.005	0.007
Sunday	3	0.005	0.015	0.028	0.025	0.012	0.005	0.011	0.022	0.005	0.013	0.009	0.010	0.004	0.004
Sunday	4	0.006	0.016	0.028	0.025	0.015	0.007	0.013	0.023	0.008	0.013	0.010	0.018	0.004	0.004
Sunday	5	0.010	0.019	0.029	0.027	0.018	0.012	0.018	0.025	0.008	0.016	0.018	0.025	0.007	0.008
Sunday	6	0.015	0.023	0.031	0.030	0.021	0.019	0.026	0.028	0.014	0.024	0.030	0.031	0.042	0.014
Sunday	7	0.021	0.029	0.035	0.034	0.029	0.030	0.039	0.030	0.022	0.034	0.035	0.050	0.019	0.022
Sunday	8	0.031	0.038	0.040	0.040	0.037	0.043	0.053	0.033	0.036	0.048	0.042	0.052	0.032	0.034
Sunday	9	0.043	0.050	0.047	0.046	0.043	0.055	0.062	0.036	0.052	0.062	0.040	0.057	0.047	0.049
Sunday	10	0.055	0.060	0.051	0.068	0.053	0.071	0.079	0.040	0.071	0.075	0.044	0.066	0.067	0.065
Sunday	11	0.063	0.065	0.054	0.055	0.060	0.077	0.080	0.044	0.082	0.086	0.070	0.055	0.080	0.074
Sunday	12	0.070	0.070	0.055	0.058	0.064	0.084	0.077	0.049	0.089	0.088	0.051	0.076	0.058	0.078
Sunday	13	0.075	0.071	0.056	0.058	0.066	0.083	0.070	0.054	0.090	0.080	0.073	0.070	0.085	0.080
Sunday	14	0.077	0.069	0.055	0.057	0.067	0.085	0.065	0.058	0.089	0.072	0.056	0.068	0.085	0.079
Sunday	15	0.078	0.070	0.053	0.057	0.072	0.083	0.061	0.063	0.087	0.069	0.059	0.071	0.084	0.077
Sunday	16	0.077	0.067	0.052	0.055	0.073	0.080	0.058	0.064	0.081	0.059	0.060	0.066	0.082	0.075
Sunday	17	0.075	0.062	0.049	0.053	0.068	0.066	0.056	0.065	0.066	0.051	0.061	0.063	0.076	0.070
Sunday	18	0.068	0.055	0.046	0.049	0.065	0.056	0.049	0.065	0.055	0.044	0.060	0.052	0.056	0.062
Sunday	19	0.061	0.047	0.042	0.045	0.058	0.043	0.041	0.062	0.043	0.036	0.059	0.050	0.049	0.055
Sunday	20	0.051	0.039	0.040	0.041	0.048	0.031	0.032	0.057	0.032	0.028	0.055	0.037	0.040	0.046
Sunday	21	0.041	0.031	0.038	0.039	0.041	0.023	0.026	0.049	0.022	0.023	0.048	0.029	0.026	0.037
Sunday	22	0.029	0.024	0.036	0.036	0.031	0.016	0.021	0.041	0.015	0.019	0.038	0.018	0.029	0.026
Sunday	23	0.019	0.019	0.037	0.033	0.020	0.012	0.017	0.028	0.012	0.016	0.028	0.014	0.019	0.015
Monday	0	0.011	0.017	0.023	0.027	0.013	0.006	0.012	0.027	0.013	0.022	0.004	0.006	0.006	0.006
Monday	1	0.007	0.015	0.022	0.025	0.012	0.006	0.011	0.023	0.006	0.011	0.023	0.004	0.004	0.003
Monday	2	0.006	0.015	0.022	0.025	0.013	0.006	0.011	0.025	0.007	0.011	0.023	0.004	0.005	0.002
Monday	3	0.009	0.018	0.025	0.027	0.015	0.010	0.012	0.027	0.010	0.011	0.024	0.006	0.011	0.003
Monday	4	0.018	0.027	0.032	0.030	0.019	0.019	0.015	0.030	0.015	0.012	0.027	0.015	0.020	0.008
Monday	5	0.030	0.039	0.039	0.036	0.025	0.030	0.021	0.033	0.022	0.018	0.035	0.035	0.032	0.024
Monday	6	0.044	0.051	0.045	0.048	0.032	0.041	0.024	0.036	0.034	0.024	0.040	0.056	0.050	0.049
Monday	7	0.058	0.058	0.050	0.048	0.034	0.048	0.032	0.040	0.043	0.030	0.044	0.063	0.057	0.075
Monday	8	0.053	0.058	0.051	0.050	0.039	0.059	0.039	0.043	0.054	0.039	0.046	0.071	0.059	0.071
Monday	9	0.051	0.059	0.053	0.051	0.047	0.065	0.046	0.045	0.067	0.048	0.046	0.066	0.060	0.057
Monday	10	0.054	0.062	0.056	0.064	0.050	0.070	0.053	0.050	0.074	0.054	0.049	0.070	0.066	0.053
Monday	11	0.057	0.064	0.057	0.066	0.056	0.072	0.055	0.052	0.075	0.059	0.051	0.070	0.065	0.056
Monday	12	0.060	0.064	0.058	0.068	0.059	0.073	0.055	0.055	0.078	0.059	0.056	0.072	0.066	0.058
Monday	13	0.061	0.064	0.058	0.067	0.060	0.076	0.058	0.057	0.081	0.060	0.055	0.073	0.071	0.058
Monday	14	0.067	0.066	0.058	0.069	0.065	0.079	0.059	0.057	0.081	0.065	0.058	0.073	0.070	0.063
Monday	15	0.072	0.065	0.057	0.073	0.071	0.081	0.062	0.059	0.080	0.063	0.061	0.077	0.074	0.072
Monday	16	0.075	0.063	0.055	0.067	0.070	0.070	0.063	0.060	0.072	0.064	0.061	0.073	0.064	0.078
Monday	17	0.074	0.055	0.051	0.061	0.065	0.057	0.066	0.057	0.059	0.066	0.059	0.059	0.057	0.080
Monday	18	0.055	0.042	0.042	0.045	0.058	0.042	0.064	0.053	0.045	0.063	0.050	0.037	0.047	0.063
Monday	19	0.042	0.031	0.036	0.039	0.054	0.031	0.059	0.048	0.032	0.060	0.040	0.032	0.036	0.042
Monday	20	0.034	0.023	0.031	0.035	0.050	0.022	0.054	0.042	0.022	0.054	0.040	0.017	0.031	0.031
Monday	21	0.027	0.018	0.028	0.032	0.041	0.017	0.051	0.036	0.016	0.046	0.035	0.013	0.023	0.025
Monday	22	0.020	0.014	0.027	0.030	0.030	0.011	0.043	0.029	0.012	0.039	0.029	0.010	0.015	0.016
Monday	23	0.014	0.011	0.025	0.030	0.022	0.008	0.034	0.020	0.008	0.031	0.022	0.006	0.011	0.009
Tues/Wed/Thurs	0	0.008	0.016	0.025	0.029	0.012	0.006	0.017	0.023	0.007	0.018	0.021	0.004	0.009	0.005
Tues/Wed/Thurs	1	0.005	0.014	0.024	0.027	0.010	0.005	0.015	0.025	0.006	0.015	0.021	0.004	0.005	0.002
Tues/Wed/Thurs	2	0.005	0.014	0.025	0.027	0.013	0.006	0.014	0.027	0.006	0.013	0.022	0.004	0.009	0.001
Tues/Wed/Thurs	3	0.008	0.018	0.028	0.033	0.014	0.009	0.015	0.029	0.009	0.013	0.024	0.005	0.012	0.002

Day of Week	Hour	Stamislauis			Sutter			Tehama			Trinity			Tulare			Tuolumne			Ventura		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Tues/Wed/Thurs	4	0.017	0.026	0.034	0.010	0.018	0.031	0.018	0.017	0.017	0.016	0.014	0.028	0.014	0.018	0.006	0.014	0.025	0.007	0.019	0.035	
Tues/Wed/Thurs	5	0.030	0.039	0.042	0.022	0.029	0.037	0.023	0.026	0.022	0.020	0.035	0.035	0.033	0.032	0.018	0.027	0.039	0.022	0.034	0.043	
Tues/Wed/Thurs	6	0.044	0.050	0.047	0.042	0.047	0.044	0.030	0.042	0.030	0.027	0.038	0.040	0.036	0.036	0.041	0.037	0.042	0.049	0.055	0.049	
Tues/Wed/Thurs	7	0.059	0.059	0.052	0.060	0.061	0.050	0.038	0.051	0.039	0.036	0.040	0.046	0.036	0.047	0.053	0.047	0.064	0.075	0.072	0.054	
Tues/Wed/Thurs	8	0.055	0.058	0.052	0.060	0.062	0.051	0.042	0.061	0.048	0.046	0.042	0.056	0.046	0.071	0.063	0.054	0.056	0.071	0.071	0.054	
Tues/Wed/Thurs	9	0.051	0.059	0.054	0.055	0.060	0.050	0.047	0.064	0.058	0.057	0.044	0.066	0.066	0.065	0.068	0.069	0.083	0.057	0.064	0.053	
Tues/Wed/Thurs	10	0.052	0.060	0.056	0.056	0.061	0.051	0.051	0.067	0.066	0.065	0.045	0.071	0.065	0.049	0.069	0.064	0.081	0.052	0.061	0.053	
Tues/Wed/Thurs	11	0.054	0.061	0.057	0.059	0.064	0.052	0.054	0.070	0.069	0.070	0.047	0.076	0.070	0.052	0.071	0.062	0.069	0.054	0.062	0.053	
Tues/Wed/Thurs	12	0.057	0.062	0.057	0.061	0.065	0.053	0.058	0.072	0.067	0.070	0.050	0.076	0.070	0.054	0.069	0.065	0.071	0.056	0.063	0.053	
Tues/Wed/Thurs	13	0.060	0.063	0.056	0.064	0.066	0.053	0.061	0.074	0.066	0.069	0.052	0.077	0.069	0.056	0.072	0.073	0.074	0.057	0.061	0.050	
Tues/Wed/Thurs	14	0.066	0.065	0.056	0.068	0.068	0.053	0.065	0.077	0.063	0.067	0.057	0.081	0.067	0.059	0.074	0.070	0.067	0.063	0.063	0.050	
Tues/Wed/Thurs	15	0.073	0.066	0.055	0.073	0.069	0.053	0.070	0.080	0.061	0.064	0.058	0.078	0.064	0.061	0.080	0.071	0.058	0.071	0.065	0.049	
Tues/Wed/Thurs	16	0.077	0.064	0.053	0.075	0.067	0.052	0.072	0.072	0.058	0.057	0.072	0.061	0.060	0.060	0.072	0.063	0.074	0.078	0.063	0.046	
Tues/Wed/Thurs	17	0.076	0.057	0.049	0.074	0.063	0.050	0.065	0.057	0.056	0.057	0.056	0.060	0.057	0.057	0.059	0.054	0.048	0.079	0.060	0.044	
Tues/Wed/Thurs	18	0.058	0.044	0.041	0.059	0.048	0.044	0.060	0.044	0.052	0.053	0.046	0.046	0.043	0.037	0.043	0.053	0.044	0.065	0.047	0.040	
Tues/Wed/Thurs	19	0.044	0.032	0.034	0.043	0.034	0.038	0.053	0.032	0.045	0.044	0.048	0.033	0.044	0.045	0.025	0.036	0.031	0.044	0.031	0.034	
Tues/Wed/Thurs	20	0.036	0.025	0.030	0.035	0.025	0.034	0.047	0.024	0.039	0.038	0.041	0.025	0.038	0.041	0.019	0.027	0.030	0.034	0.021	0.030	
Tues/Wed/Thurs	21	0.028	0.019	0.026	0.029	0.019	0.031	0.042	0.021	0.034	0.032	0.035	0.018	0.032	0.035	0.014	0.021	0.023	0.028	0.016	0.029	
Tues/Wed/Thurs	22	0.021	0.014	0.025	0.020	0.013	0.029	0.031	0.013	0.028	0.026	0.029	0.014	0.026	0.029	0.010	0.015	0.017	0.018	0.011	0.028	
Tues/Wed/Thurs	23	0.015	0.012	0.023	0.013	0.009	0.028	0.022	0.010	0.022	0.021	0.022	0.010	0.021	0.022	0.006	0.011	0.010	0.010	0.008	0.030	
Friday	0	0.008	0.016	0.027	0.007	0.014	0.032	0.013	0.007	0.021	0.007	0.019	0.007	0.019	0.020	0.004	0.010	0.005	0.006	0.016	0.033	
Friday	1	0.006	0.014	0.025	0.005	0.011	0.030	0.012	0.006	0.018	0.023	0.006	0.017	0.021	0.003	0.003	0.008	0.003	0.003	0.013	0.031	
Friday	2	0.005	0.014	0.026	0.004	0.011	0.030	0.012	0.006	0.018	0.024	0.007	0.016	0.023	0.004	0.008	0.002	0.002	0.002	0.012	0.031	
Friday	3	0.008	0.017	0.029	0.005	0.012	0.030	0.014	0.008	0.018	0.026	0.009	0.016	0.022	0.005	0.013	0.002	0.008	0.003	0.014	0.032	
Friday	4	0.014	0.024	0.035	0.008	0.016	0.033	0.016	0.015	0.021	0.029	0.013	0.019	0.027	0.013	0.020	0.005	0.005	0.007	0.019	0.036	
Friday	5	0.024	0.035	0.042	0.017	0.026	0.038	0.023	0.023	0.026	0.032	0.018	0.023	0.034	0.032	0.033	0.013	0.023	0.020	0.032	0.042	
Friday	6	0.036	0.045	0.047	0.033	0.040	0.045	0.029	0.035	0.033	0.033	0.030	0.032	0.038	0.051	0.057	0.026	0.035	0.043	0.052	0.049	
Friday	7	0.049	0.053	0.052	0.049	0.054	0.050	0.039	0.044	0.041	0.037	0.039	0.039	0.042	0.062	0.063	0.039	0.040	0.064	0.068	0.052	
Friday	8	0.047	0.054	0.053	0.051	0.052	0.052	0.039	0.055	0.049	0.040	0.040	0.051	0.049	0.046	0.070	0.063	0.043	0.064	0.069	0.054	
Friday	9	0.047	0.056	0.055	0.050	0.057	0.052	0.042	0.060	0.055	0.045	0.063	0.054	0.047	0.066	0.063	0.049	0.054	0.054	0.062	0.053	
Friday	10	0.051	0.060	0.058	0.054	0.061	0.054	0.049	0.063	0.058	0.048	0.069	0.060	0.050	0.070	0.066	0.058	0.053	0.061	0.061	0.054	
Friday	11	0.054	0.062	0.060	0.060	0.066	0.055	0.052	0.069	0.061	0.063	0.063	0.063	0.063	0.052	0.071	0.063	0.064	0.057	0.064	0.054	
Friday	12	0.057	0.063	0.060	0.063	0.067	0.055	0.057	0.070	0.061	0.052	0.074	0.063	0.054	0.070	0.067	0.066	0.059	0.059	0.064	0.053	
Friday	13	0.061	0.065	0.059	0.066	0.068	0.054	0.057	0.075	0.061	0.055	0.077	0.062	0.056	0.072	0.067	0.071	0.074	0.061	0.065	0.052	
Friday	14	0.068	0.067	0.058	0.070	0.070	0.054	0.065	0.080	0.060	0.059	0.080	0.072	0.063	0.058	0.074	0.070	0.076	0.065	0.065	0.050	
Friday	15	0.074	0.067	0.056	0.073	0.070	0.052	0.070	0.082	0.059	0.063	0.081	0.061	0.059	0.075	0.068	0.083	0.071	0.071	0.065	0.049	
Friday	16	0.076	0.064	0.053	0.074	0.067	0.050	0.072	0.073	0.057	0.058	0.075	0.059	0.059	0.070	0.059	0.083	0.077	0.075	0.063	0.046	
Friday	17	0.075	0.058	0.048	0.072	0.063	0.047	0.065	0.062	0.055	0.059	0.063	0.055	0.055	0.057	0.055	0.075	0.064	0.074	0.059	0.043	
Friday	18	0.064	0.048	0.040	0.063	0.051	0.042	0.061	0.047	0.051	0.054	0.052	0.051	0.053	0.041	0.043	0.062	0.064	0.064	0.046	0.040	
Friday	19	0.052	0.037	0.032	0.050	0.039	0.035	0.059	0.039	0.046	0.050	0.036	0.046	0.045	0.027	0.036	0.050	0.048	0.048	0.032	0.034	
Friday	20	0.043	0.029	0.026	0.041	0.029	0.030	0.051	0.028	0.040	0.046	0.030	0.041	0.042	0.020	0.026	0.041	0.037	0.037	0.022	0.029	
Friday	21	0.035	0.022	0.022	0.037	0.023	0.028	0.045	0.022	0.035	0.040	0.022	0.036	0.039	0.017	0.019	0.036	0.032	0.032	0.022	0.029	
Friday	22	0.027	0.016	0.020	0.030	0.017	0.026	0.037	0.018	0.031	0.031	0.016	0.031	0.032	0.014	0.015	0.030	0.024	0.024	0.017	0.027	
Friday	23	0.020	0.012	0.018	0.020	0.011	0.024	0.026	0.012	0.025	0.025	0.012	0.025	0.025	0.011	0.010	0.018	0.016	0.016	0.009	0.027	
Saturday	0	0.015	0.026	0.040	0.013	0.019	0.038	0.015	0.011	0.021	0.026	0.013	0.020	0.025	0.010	0.013	0.010	0.015	0.011	0.024	0.043	
Saturday	1	0.010	0.020	0.035	0.008	0.015	0.034	0.014	0.008	0.018	0.026	0.008	0.016	0.025	0.007	0.010	0.007	0.012	0.006	0.018	0.040	
Saturday	2	0.008	0.018	0.032	0.006	0.014	0.032	0.014	0.008	0.018	0.027	0.007	0.015	0.026	0.005	0.011	0.005	0.004	0.004	0.016	0.038	
Saturday	3	0.008	0.019	0.032	0.006	0.013	0.031	0.014	0.007	0.016	0.030	0.007	0.014	0.027	0.009	0.013	0.004	0.010	0.003	0.015	0.037	
Saturday	4	0.011	0.021	0.035	0.007	0.014	0.032	0.017	0.014	0.017	0.029	0.009	0.016	0.029	0.014	0.024	0.005	0.013	0.005	0.017	0.038	
Saturday	5	0.017	0.028	0.039	0.011	0.018	0.034	0.021	0.018	0.021	0.033	0.015	0.019	0.036	0.011	0.019	0.021	0.021	0.011	0.023	0.041	
Saturday	6	0.025	0.036	0.045	0.019	0.026	0.039	0.025	0.027	0.028	0.036	0.023	0.025	0.042	0.056	0.054	0.017	0.028	0.021	0.033	0.045	
Saturday	7	0.034	0.044	0.050	0.032	0.038	0.046	0.032	0.038	0.039	0.038	0.033	0.036	0.041	0.055	0.068	0.029	0.034	0.034	0.046	0.050	

Day of Week	Hour	Stamslaus			Sutter			Tehama			Trinity			Tulare			Tuolumne			Ventura		
		LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH	LD	LM	HH
Saturday	8	0.044	0.053	0.052	0.045	0.051	0.052	0.040	0.055	0.051	0.047	0.047	0.043	0.057	0.069	0.044	0.045	0.060	0.046	0.057	0.053	
Saturday	9	0.054	0.061	0.062	0.057	0.064	0.062	0.044	0.064	0.061	0.063	0.059	0.045	0.061	0.069	0.049	0.059	0.060	0.057	0.065	0.055	
Saturday	10	0.062	0.068	0.063	0.067	0.071	0.060	0.051	0.067	0.067	0.067	0.067	0.048	0.066	0.068	0.049	0.073	0.078	0.065	0.071	0.056	
Saturday	11	0.067	0.071	0.064	0.074	0.076	0.061	0.058	0.077	0.068	0.084	0.073	0.050	0.067	0.068	0.081	0.077	0.083	0.070	0.076	0.056	
Saturday	12	0.069	0.070	0.062	0.075	0.075	0.060	0.060	0.076	0.067	0.083	0.071	0.052	0.068	0.065	0.078	0.077	0.075	0.072	0.074	0.054	
Saturday	13	0.070	0.067	0.058	0.075	0.074	0.057	0.059	0.073	0.066	0.081	0.069	0.053	0.067	0.068	0.075	0.072	0.060	0.072	0.071	0.053	
Saturday	14	0.070	0.064	0.054	0.074	0.071	0.055	0.065	0.076	0.066	0.076	0.065	0.055	0.070	0.070	0.075	0.068	0.055	0.072	0.068	0.050	
Saturday	15	0.069	0.061	0.049	0.072	0.068	0.051	0.067	0.073	0.064	0.062	0.062	0.058	0.077	0.065	0.075	0.068	0.052	0.072	0.063	0.047	
Saturday	16	0.068	0.057	0.045	0.070	0.064	0.048	0.065	0.069	0.059	0.070	0.058	0.057	0.066	0.055	0.072	0.070	0.047	0.072	0.059	0.044	
Saturday	17	0.064	0.051	0.040	0.066	0.057	0.044	0.064	0.062	0.055	0.061	0.057	0.052	0.064	0.050	0.066	0.063	0.040	0.068	0.051	0.040	
Saturday	18	0.056	0.042	0.033	0.056	0.047	0.038	0.061	0.048	0.050	0.049	0.052	0.040	0.039	0.050	0.058	0.052	0.031	0.059	0.041	0.035	
Saturday	19	0.048	0.034	0.027	0.046	0.037	0.033	0.059	0.041	0.044	0.049	0.038	0.045	0.034	0.030	0.047	0.041	0.026	0.048	0.031	0.030	
Saturday	20	0.041	0.029	0.024	0.040	0.030	0.028	0.050	0.031	0.036	0.042	0.031	0.038	0.042	0.027	0.021	0.038	0.020	0.040	0.024	0.027	
Saturday	21	0.037	0.024	0.021	0.035	0.025	0.025	0.044	0.023	0.030	0.037	0.023	0.031	0.038	0.023	0.018	0.031	0.016	0.037	0.022	0.024	
Saturday	22	0.031	0.020	0.019	0.028	0.019	0.023	0.034	0.017	0.024	0.031	0.017	0.026	0.032	0.019	0.011	0.025	0.020	0.031	0.019	0.023	
Saturday	23	0.023	0.016	0.017	0.020	0.014	0.021	0.026	0.013	0.019	0.023	0.012	0.019	0.023	0.014	0.008	0.016	0.013	0.022	0.016	0.022	
Holiday	0	0.013	0.020	0.027	0.010	0.016	0.028	0.014	0.008	0.015	0.024	0.008	0.015	0.024	0.008	0.009	0.008	0.011	0.009	0.019	0.032	
Holiday	1	0.009	0.017	0.025	0.006	0.013	0.027	0.013	0.007	0.013	0.027	0.008	0.012	0.024	0.007	0.010	0.005	0.009	0.005	0.016	0.030	
Holiday	2	0.007	0.015	0.024	0.004	0.012	0.026	0.013	0.006	0.012	0.024	0.008	0.012	0.023	0.006	0.007	0.003	0.010	0.003	0.014	0.029	
Holiday	3	0.007	0.016	0.026	0.005	0.013	0.027	0.013	0.006	0.012	0.029	0.010	0.013	0.023	0.007	0.011	0.004	0.010	0.003	0.015	0.031	
Holiday	4	0.011	0.020	0.029	0.008	0.016	0.029	0.016	0.013	0.014	0.029	0.012	0.014	0.027	0.016	0.017	0.005	0.012	0.007	0.018	0.032	
Holiday	5	0.019	0.028	0.033	0.014	0.023	0.032	0.020	0.017	0.020	0.031	0.016	0.017	0.033	0.030	0.032	0.009	0.018	0.016	0.029	0.038	
Holiday	6	0.027	0.035	0.038	0.025	0.033	0.036	0.025	0.028	0.026	0.037	0.025	0.023	0.035	0.045	0.052	0.018	0.023	0.031	0.042	0.043	
Holiday	7	0.035	0.042	0.042	0.036	0.044	0.042	0.030	0.037	0.036	0.038	0.033	0.031	0.040	0.052	0.064	0.029	0.031	0.047	0.056	0.047	
Holiday	8	0.040	0.048	0.046	0.046	0.053	0.048	0.036	0.051	0.046	0.040	0.049	0.040	0.043	0.065	0.066	0.044	0.044	0.051	0.059	0.049	
Holiday	9	0.048	0.055	0.050	0.054	0.059	0.050	0.047	0.068	0.056	0.043	0.062	0.054	0.045	0.061	0.058	0.058	0.057	0.052	0.061	0.051	
Holiday	10	0.059	0.064	0.055	0.065	0.069	0.053	0.051	0.068	0.064	0.050	0.076	0.060	0.050	0.075	0.055	0.076	0.083	0.059	0.066	0.053	
Holiday	11	0.065	0.070	0.060	0.074	0.074	0.057	0.059	0.083	0.069	0.047	0.084	0.068	0.049	0.076	0.060	0.085	0.086	0.066	0.069	0.054	
Holiday	12	0.069	0.072	0.061	0.077	0.074	0.056	0.066	0.081	0.071	0.053	0.083	0.070	0.058	0.075	0.060	0.085	0.087	0.068	0.072	0.055	
Holiday	13	0.071	0.071	0.061	0.076	0.074	0.058	0.062	0.084	0.068	0.062	0.091	0.067	0.052	0.069	0.068	0.083	0.081	0.070	0.070	0.053	
Holiday	14	0.072	0.069	0.059	0.075	0.073	0.056	0.069	0.076	0.064	0.059	0.087	0.069	0.055	0.069	0.070	0.080	0.074	0.071	0.068	0.053	
Holiday	15	0.073	0.068	0.058	0.074	0.070	0.055	0.065	0.081	0.061	0.057	0.079	0.065	0.062	0.070	0.078	0.078	0.074	0.073	0.064	0.050	
Holiday	16	0.073	0.065	0.055	0.072	0.066	0.054	0.070	0.068	0.061	0.056	0.072	0.062	0.065	0.074	0.069	0.078	0.072	0.073	0.061	0.049	
Holiday	17	0.070	0.057	0.050	0.068	0.059	0.051	0.068	0.063	0.060	0.056	0.058	0.060	0.053	0.057	0.062	0.071	0.066	0.071	0.056	0.046	
Holiday	18	0.060	0.046	0.044	0.057	0.049	0.045	0.063	0.047	0.055	0.053	0.044	0.058	0.051	0.040	0.046	0.057	0.049	0.061	0.045	0.041	
Holiday	19	0.050	0.036	0.039	0.047	0.036	0.041	0.056	0.035	0.048	0.048	0.029	0.049	0.047	0.031	0.041	0.043	0.040	0.049	0.032	0.036	
Holiday	20	0.042	0.029	0.034	0.039	0.029	0.037	0.050	0.028	0.041	0.044	0.024	0.045	0.046	0.027	0.026	0.033	0.026	0.041	0.024	0.033	
Holiday	21	0.034	0.023	0.030	0.030	0.020	0.033	0.045	0.021	0.035	0.040	0.019	0.040	0.040	0.019	0.021	0.024	0.018	0.034	0.019	0.032	
Holiday	22	0.027	0.017	0.028	0.023	0.015	0.031	0.027	0.013	0.029	0.031	0.014	0.030	0.034	0.014	0.014	0.017	0.012	0.025	0.014	0.031	
Holiday	23	0.018	0.014	0.026	0.015	0.010	0.029	0.022	0.010	0.023	0.024	0.009	0.024	0.024	0.011	0.011	0.010	0.008	0.016	0.012	0.032	

Day of Week	Hour	Yolo			Yuba		
		LD	LM	HH	LD	LM	HH
Sunday	0	0.016	0.026	0.044	0.013	0.020	0.031
Sunday	1	0.011	0.019	0.036	0.008	0.016	0.028
Sunday	2	0.008	0.017	0.033	0.006	0.013	0.026
Sunday	3	0.006	0.015	0.030	0.005	0.012	0.025
Sunday	4	0.007	0.016	0.029	0.005	0.012	0.025
Sunday	5	0.011	0.020	0.032	0.008	0.015	0.027
Sunday	6	0.016	0.025	0.034	0.013	0.020	0.030
Sunday	7	0.023	0.031	0.040	0.022	0.028	0.034
Sunday	8	0.034	0.041	0.046	0.034	0.041	0.040
Sunday	9	0.048	0.054	0.051	0.048	0.055	0.046
Sunday	10	0.060	0.063	0.054	0.064	0.068	0.052
Sunday	11	0.067	0.067	0.054	0.075	0.075	0.055
Sunday	12	0.071	0.070	0.053	0.082	0.079	0.058
Sunday	13	0.072	0.070	0.052	0.084	0.079	0.058
Sunday	14	0.073	0.069	0.050	0.084	0.077	0.057
Sunday	15	0.073	0.067	0.047	0.082	0.073	0.057
Sunday	16	0.072	0.063	0.045	0.079	0.068	0.055
Sunday	17	0.070	0.059	0.043	0.072	0.062	0.053
Sunday	18	0.063	0.051	0.041	0.060	0.052	0.049
Sunday	19	0.057	0.044	0.038	0.050	0.043	0.045
Sunday	20	0.051	0.038	0.036	0.041	0.035	0.042
Sunday	21	0.042	0.032	0.037	0.031	0.026	0.039
Sunday	22	0.030	0.025	0.037	0.021	0.019	0.036
Sunday	23	0.019	0.020	0.040	0.013	0.015	0.033
Monday	0	0.010	0.018	0.028	0.008	0.014	0.027
Monday	1	0.006	0.015	0.026	0.005	0.012	0.025
Monday	2	0.005	0.014	0.026	0.004	0.012	0.025
Monday	3	0.007	0.016	0.028	0.006	0.014	0.027
Monday	4	0.016	0.025	0.034	0.011	0.019	0.030
Monday	5	0.032	0.040	0.043	0.023	0.030	0.036
Monday	6	0.048	0.052	0.050	0.042	0.047	0.043
Monday	7	0.066	0.065	0.056	0.060	0.061	0.048
Monday	8	0.064	0.064	0.057	0.059	0.062	0.050
Monday	9	0.057	0.062	0.056	0.056	0.061	0.050
Monday	10	0.055	0.061	0.057	0.058	0.064	0.051
Monday	11	0.056	0.062	0.056	0.062	0.066	0.053
Monday	12	0.058	0.062	0.056	0.066	0.068	0.054
Monday	13	0.059	0.061	0.055	0.067	0.067	0.054
Monday	14	0.062	0.062	0.054	0.070	0.069	0.055
Monday	15	0.068	0.063	0.053	0.073	0.069	0.055
Monday	16	0.073	0.062	0.051	0.075	0.067	0.054
Monday	17	0.072	0.057	0.046	0.073	0.061	0.052
Monday	18	0.053	0.043	0.039	0.056	0.046	0.045
Monday	19	0.039	0.030	0.031	0.040	0.031	0.039
Monday	20	0.032	0.023	0.026	0.031	0.022	0.035
Monday	21	0.027	0.018	0.024	0.025	0.017	0.032
Monday	22	0.021	0.014	0.023	0.017	0.012	0.030
Monday	23	0.014	0.011	0.025	0.012	0.009	0.030
Tues/Wed/Thurs	0	0.009	0.017	0.031	0.008	0.014	0.029
Tues/Wed/Thurs	1	0.006	0.014	0.028	0.004	0.011	0.027
Tues/Wed/Thurs	2	0.005	0.014	0.028	0.004	0.011	0.027
Tues/Wed/Thurs	3	0.006	0.016	0.030	0.005	0.013	0.029


Day of Week	Hour	Yolo			Yuba		
		LD	LM	HH	LD	LM	HH
Tues/Wed/Thurs	4	0.014	0.023	0.036	0.010	0.018	0.031
Tues/Wed/Thurs	5	0.029	0.037	0.044	0.022	0.029	0.037
Tues/Wed/Thurs	6	0.046	0.051	0.052	0.042	0.047	0.044
Tues/Wed/Thurs	7	0.066	0.065	0.057	0.060	0.061	0.050
Tues/Wed/Thurs	8	0.065	0.064	0.057	0.060	0.062	0.051
Tues/Wed/Thurs	9	0.057	0.062	0.057	0.055	0.060	0.050
Tues/Wed/Thurs	10	0.053	0.061	0.057	0.056	0.061	0.051
Tues/Wed/Thurs	11	0.054	0.061	0.057	0.059	0.064	0.052
Tues/Wed/Thurs	12	0.056	0.061	0.056	0.061	0.065	0.053
Tues/Wed/Thurs	13	0.058	0.061	0.055	0.064	0.066	0.053
Tues/Wed/Thurs	14	0.062	0.062	0.053	0.068	0.068	0.053
Tues/Wed/Thurs	15	0.069	0.063	0.051	0.073	0.069	0.053
Tues/Wed/Thurs	16	0.074	0.062	0.048	0.075	0.067	0.052
Tues/Wed/Thurs	17	0.073	0.058	0.044	0.074	0.063	0.050
Tues/Wed/Thurs	18	0.056	0.045	0.037	0.059	0.048	0.044
Tues/Wed/Thurs	19	0.041	0.032	0.030	0.043	0.034	0.038
Tues/Wed/Thurs	20	0.034	0.025	0.025	0.035	0.025	0.034
Tues/Wed/Thurs	21	0.029	0.020	0.023	0.029	0.019	0.031
Tues/Wed/Thurs	22	0.022	0.015	0.022	0.020	0.013	0.029
Tues/Wed/Thurs	23	0.015	0.011	0.023	0.013	0.009	0.028
Friday	0	0.009	0.017	0.032	0.007	0.014	0.032
Friday	1	0.006	0.014	0.030	0.005	0.011	0.030
Friday	2	0.005	0.014	0.030	0.004	0.011	0.030
Friday	3	0.006	0.015	0.032	0.005	0.012	0.030
Friday	4	0.012	0.022	0.037	0.008	0.016	0.033
Friday	5	0.024	0.034	0.044	0.017	0.026	0.038
Friday	6	0.038	0.047	0.052	0.033	0.040	0.045
Friday	7	0.054	0.059	0.058	0.049	0.054	0.050
Friday	8	0.055	0.059	0.059	0.051	0.057	0.052
Friday	9	0.051	0.059	0.058	0.050	0.057	0.052
Friday	10	0.052	0.060	0.058	0.054	0.061	0.054
Friday	11	0.056	0.062	0.058	0.060	0.066	0.055
Friday	12	0.059	0.063	0.056	0.063	0.067	0.055
Friday	13	0.062	0.064	0.055	0.066	0.068	0.054
Friday	14	0.066	0.064	0.053	0.070	0.070	0.054
Friday	15	0.070	0.063	0.050	0.073	0.070	0.052
Friday	16	0.071	0.061	0.046	0.074	0.067	0.050
Friday	17	0.069	0.057	0.041	0.072	0.063	0.047
Friday	18	0.060	0.047	0.037	0.063	0.051	0.042
Friday	19	0.049	0.036	0.029	0.050	0.039	0.035
Friday	20	0.041	0.028	0.024	0.041	0.029	0.030
Friday	21	0.036	0.023	0.021	0.037	0.023	0.028
Friday	22	0.029	0.018	0.019	0.030	0.017	0.026
Friday	23	0.019	0.013	0.019	0.019	0.011	0.024
Saturday	0	0.014	0.024	0.050	0.013	0.019	0.038
Saturday	1	0.009	0.019	0.042	0.008	0.015	0.034
Saturday	2	0.008	0.017	0.039	0.006	0.014	0.032
Saturday	3	0.007	0.016	0.037	0.006	0.013	0.031
Saturday	4	0.009	0.019	0.038	0.007	0.014	0.032
Saturday	5	0.014	0.025	0.043	0.011	0.018	0.034
Saturday	6	0.023	0.033	0.049	0.019	0.026	0.039
Saturday	7	0.034	0.044	0.055	0.032	0.038	0.046

Day of Week	Hour	Yolo			Yuba		
		LD	LM	HH	LD	LM	HH
Saturday	8	0.046	0.055	0.059	0.045	0.051	0.052
Saturday	9	0.057	0.064	0.061	0.057	0.062	0.056
Saturday	10	0.065	0.070	0.063	0.067	0.071	0.060
Saturday	11	0.069	0.071	0.059	0.074	0.076	0.061
Saturday	12	0.069	0.068	0.056	0.075	0.075	0.060
Saturday	13	0.069	0.065	0.052	0.075	0.074	0.057
Saturday	14	0.068	0.063	0.047	0.074	0.071	0.055
Saturday	15	0.067	0.060	0.043	0.072	0.068	0.051
Saturday	16	0.066	0.056	0.039	0.070	0.064	0.048
Saturday	17	0.063	0.052	0.035	0.066	0.057	0.044
Saturday	18	0.057	0.045	0.029	0.056	0.047	0.038
Saturday	19	0.048	0.035	0.025	0.046	0.037	0.033
Saturday	20	0.042	0.030	0.021	0.040	0.030	0.028
Saturday	21	0.039	0.027	0.020	0.035	0.025	0.025
Saturday	22	0.034	0.023	0.020	0.028	0.019	0.023
Saturday	23	0.024	0.018	0.019	0.020	0.014	0.021
Holiday	0	0.012	0.022	0.032	0.010	0.016	0.028
Holiday	1	0.008	0.017	0.029	0.006	0.013	0.027
Holiday	2	0.006	0.015	0.029	0.004	0.012	0.026
Holiday	3	0.006	0.017	0.029	0.005	0.013	0.027
Holiday	4	0.011	0.021	0.032	0.008	0.016	0.029
Holiday	5	0.019	0.030	0.038	0.014	0.023	0.032
Holiday	6	0.027	0.038	0.044	0.025	0.033	0.036
Holiday	7	0.037	0.046	0.050	0.036	0.044	0.042
Holiday	8	0.046	0.054	0.053	0.046	0.053	0.048
Holiday	9	0.053	0.059	0.056	0.054	0.059	0.050
Holiday	10	0.061	0.065	0.058	0.065	0.069	0.053
Holiday	11	0.067	0.069	0.060	0.074	0.074	0.057
Holiday	12	0.069	0.068	0.059	0.077	0.074	0.056
Holiday	13	0.069	0.068	0.057	0.076	0.074	0.058
Holiday	14	0.070	0.066	0.055	0.075	0.073	0.056
Holiday	15	0.069	0.065	0.052	0.074	0.070	0.055
Holiday	16	0.067	0.060	0.049	0.072	0.066	0.054
Holiday	17	0.064	0.055	0.044	0.068	0.059	0.051
Holiday	18	0.057	0.046	0.039	0.057	0.049	0.045
Holiday	19	0.050	0.036	0.033	0.047	0.036	0.041
Holiday	20	0.044	0.029	0.028	0.039	0.029	0.037
Holiday	21	0.039	0.023	0.025	0.030	0.020	0.033
Holiday	22	0.030	0.018	0.024	0.023	0.015	0.031
Holiday	23	0.020	0.014	0.026	0.015	0.010	0.029

### Appendix C: Scaling procedures after DTIM processing


#### C1. Block Diagram of Scaling Process: Idg (gas: heavy- and light-duty; diesel: light-duty)

DTIM has 1 to 12 Source Classification Codes (SCC) that vary by species. For CO, NOx, SOx and PM species, DTIM only uses SCC=1 for the running exhaust emissions regardless of the fuel type and process. However, distribution of the running exhaust emissions according to the fuel type and process is needed. The following diagram explains how to distribute the running exhaust emissions for the light-duty gas. The running exhaust emissions are distributed to the catalyst cold exhaust, catalyst hot exhaust, non-catalyst cold exhaust, non-catalyst hot exhaust, catalyst bus and non-catalyst bus by using the corresponding emissions from EMFAC. Since there are no idle emissions in DTIM, surrogates are needed for the catalyst idle and non-catalyst idle. The surrogates for the catalyst idle and non-catalyst idle are catalyst hot exhaust, and non-catalyst hot exhaust, respectively.



**C2. Block Diagram of Scaling Process: hdd (heavy-duty diesel)**

The following diagram explains how to distribute the running exhaust emissions for heavy-duty diesel. The running exhaust emissions are distributed to the diesel exhaust or diesel bus exhaust depending on the vehicle type by using the corresponding emissions from EMFAC. Since there are no idle emissions in DTIM, a surrogate is used. The surrogate for the diesel idle emissions is diesel exhaust or diesel bus exhaust, depending on the vehicle type.





**Appendix D: Additional temporal profiles**

Temporal profiles developed from the AGTOOL are applied as potential replacements when processing the emissions inventories for modeling using the SMOKE processor. This would apply for agriculturally related emissions with time-invariant temporal distributions, which includes the following emission source categories: food and agricultural processing, pesticides and fertilizers, farming operations, unpaved road dust, fugitive windblown dust, managed burning and disposal, and farming equipment

Table 11 Day of week temporal profiles from the Agricultural Emissions Temporal and Spatial Allocation Tool (AgTool)

Code	M	T	W	TH	F	S	S
201	1	174	248	182	203	97	95
202	1	2	1	0	2	1	993
203	1	117	192	190	229	222	48
204	2	16	13	13	10	928	17
205	3	342	597	25	4	5	24
206	4	100	33	241	105	455	62
207	5	50	284	126	125	315	95
208	6	94	41	40	348	358	112
209	7	203	111	236	340	0	102
210	8	221	225	123	117	80	225
211	9	37	63	667	111	37	77
212	11	2	881	41	40	18	8
213	12	96	105	153	201	425	8
214	13	370	306	90	47	101	73
215	13	368	72	498	2	41	6
216	19	562	125	102	47	39	107
217	22	348	74	115	125	215	102
218	22	292	63	229	65	104	224
219	22	482	41	111	167	93	83
220	25	184	100	136	223	152	182
221	25	192	107	223	278	75	101
222	27	40	51	99	310	58	415
223	29	51	237	127	172	308	77
224	30	219	195	158	222	112	64
225	30	185	151	125	186	120	203
226	35	131	195	172	151	201	114
227	35	146	162	175	157	180	143
228	36	179	200	93	188	186	117
229	37	82	363	208	2	73	235
230	40	211	162	182	160	165	81
231	40	468	0	420	0	72	0
232	41	269	293	118	95	121	62
233	44	56	399	13	268	61	160
234	45	335	72	82	210	180	77
235	46	124	139	148	199	168	177
236	46	207	54	453	54	134	52
237	48	310	346	83	84	91	38
238	52	201	140	196	121	160	132
239	53	134	123	144	206	192	149
240	53	108	150	163	171	207	148
241	57	156	183	117	92	220	175
242	63	105	176	154	148	195	160
243	63	186	136	175	187	134	120

Code	M	T	W	TH	F	S	S
244	64	230	173	136	83	251	63
245	66	249	149	127	105	185	120
246	67	222	278	236	65	129	2
247	70	120	192	168	188	145	116
248	74	95	170	197	157	144	162
249	74	190	108	126	246	116	138
250	77	295	104	187	155	88	93
251	79	135	291	129	86	182	97
252	80	360	9	19	424	79	29
253	81	133	132	125	226	167	135
254	82	136	151	118	160	196	157
255	82	92	125	207	177	153	164
256	85	133	152	145	188	173	124
257	87	295	16	111	47	244	201
258	96	128	104	169	161	224	119
259	104	196	118	155	202	132	94
260	104	111	196	121	181	127	162
261	107	161	70	90	227	243	102
262	107	145	115	203	187	147	95
263	111	171	137	0	297	202	81
264	112	121	144	165	155	172	131
265	113	199	97	132	218	147	94
266	113	167	15	156	399	70	80
267	115	150	128	153	192	139	122
268	115	103	120	138	117	251	156
269	119	125	119	87	144	158	248
270	120	145	130	137	155	166	147
271	125	155	141	108	179	149	142
272	130	140	137	170	93	139	192
273	135	222	191	83	169	110	90
274	136	160	156	162	144	156	86
275	138	109	107	137	227	147	137
276	139	101	117	171	167	171	134
277	143	143	143	143	143	143	143
278	150	230	118	72	144	170	116
279	163	118	106	135	185	112	181
280	199	136	81	163	143	180	99
281	218	8	2	14	6	525	226
282	250	35	290	130	50	109	137
283	255	116	82	103	128	63	252
284	278	182	148	36	105	112	139
285	326	168	189	0	105	0	211
286	0	212	165	131	202	128	161
287	0	289	0	0	356	222	133
288	0	321	93	208	109	81	188
289	0	431	4	160	246	15	144
290	0	515	122	111	48	128	76
291	0	0	0	916	84	0	0
292	0	0	0	0	148	0	852
294	0	0	0	0	1000	0	0

Table 12 Daily temporal profiles from the Agricultural Emissions Temporal and Spatial Allocation Tool (AgTool)

Code	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
201	0	0	0	0	0	10	102	2	26	358	259	134	65	1	26	10	3	2	1	0	0	0	0	0
202	0	0	0	5	3	2	5	59	44	38	28	640	19	21	48	34	21	22	10	1	0	1	0	0
203	1	0	0	0	10	162	64	51	139	270	115	46	61	3	15	16	4	4	12	6	3	1	3	2
204	1	0	0	0	0	1	139	405	79	126	69	54	33	31	13	20	14	14	2	0	0	0	0	0
205	1	3	6	2	3	8	1	2	5	29	73	112	125	115	101	164	46	49	65	68	3	10	5	2
206	2	5	0	4	22	5	6	8	26	31	88	90	66	397	38	28	43	100	34	5	0	0	0	0
207	2	3	0	0	37	177	45	57	167	203	123	102	23	15	8	6	22	6	1	0	0	0	0	1
208	2	0	0	0	0	20	1	498	9	15	28	8	42	6	358	2	0	0	9	0	0	0	0	0
209	2	0	0	12	54	3	41	471	18	105	94	31	7	9	68	33	43	7	0	0	0	0	0	0
210	2	4	2	4	4	3	17	40	60	137	87	178	42	67	82	198	60	6	3	1	1	1	1	1
211	3	2	3	2	0	2	6	12	43	75	220	413	2	199	2	5	4	7	0	0	0	0	0	0
212	4	5	0	0	6	220	16	73	212	321	135	6	0	0	0	0	0	0	3	0	0	0	0	0
213	4	159	11	187	7	0	0	16	71	536	0	1	0	0	0	0	0	0	7	0	0	0	0	0
214	5	5	5	7	6	13	6	91	50	29	237	161	11	37	123	78	76	1	51	1	1	1	1	2
215	8	5	19	15	44	48	35	44	88	109	96	100	58	112	62	44	30	52	13	3	3	3	6	6
216	9	0	0	0	0	10	19	157	83	105	65	92	15	19	73	308	32	6	2	4	1	0	1	0
217	9	9	6	7	10	84	13	35	113	187	138	63	57	58	25	40	44	45	30	4	5	4	3	13
218	10	3	6	5	7	11	17	61	30	44	61	73	88	56	119	265	18	3	108	3	1	3	3	6
219	0	0	0	0	0	393	374	26	0	139	0	4	11	1	2	15	33	2	0	0	0	0	0	0
220	11	11	8	2	25	16	144	131	173	251	106	55	56	4	1	4	1	0	0	0	0	0	0	0
221	13	13	15	25	32	11	8	12	8	123	19	135	6	47	157	65	26	96	154	7	6	6	6	8
222	9	9	2	19	3	19	7	16	76	20	39	156	44	277	29	52	176	37	2	2	2	1	1	2
223	5	5	3	4	13	23	108	64	68	61	92	278	59	38	56	34	32	22	14	5	1	1	2	5
224	1	1	10	4	8	32	50	118	64	72	75	123	130	51	72	63	61	24	8	2	16	2	11	1
225	4	4	8	12	25	22	33	74	62	76	86	114	72	84	86	92	80	33	12	7	3	4	3	4
226	4	4	8	11	12	26	26	46	37	85	114	231	83	67	91	57	12	4	4	1	2	3	2	2
227	7	7	9	10	19	39	25	45	61	92	97	102	73	120	66	66	72	45	19	7	5	5	5	5
228	4	4	8	9	28	20	30	24	34	58	53	180	122	60	128	104	67	29	22	3	2	4	4	3
229	10	10	15	14	18	171	37	47	47	41	38	40	45	22	27	57	13	3	305	4	6	5	5	20
230	19	19	40	29	38	80	48	119	50	39	31	35	75	49	84	80	64	27	22	21	12	10	9	1
231	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42	42
232	0	0	0	0	0	2	20	24	22	21	37	146	32	41	17	219	406	5	4	4	0	1	0	0
233	0	0	0	0	0	0	0	0	512	0	0	0	0	0	488	0	0	0	0	0	0	0	0	0
234	9	9	7	5	9	32	20	58	39	80	110	105	136	66	131	41	89	12	16	9	9	0	7	1
235	2	2	2	2	6	31	48	95	72	51	41	460	48	29	19	20	34	17	9	8	1	0	0	0
236	11	11	23	12	20	28	23	22	28	64	96	55	75	53	105	105	146	58	13	11	8	10	14	9
237	18	18	12	10	15	7	11	24	20	49	77	80	54	38	59	177	120	20	10	35	38	44	39	26
238	1	1	1	4	1	20	52	86	79	118	93	120	71	56	132	73	42	27	8	4	2	3	3	1
239	2	2	1	3	2	42	31	82	79	87	78	85	78	76	67	142	38	15	4	1	2	2	2	1
240	0	0	0	0	19	27	55	26	23	26	51	112	162	192	85	60	22	8	1	12	6	0	0	1
241	3	3	7	34	3	37	32	238	35	45	66	70	64	43	166	68	52	16	4	5	1	1	4	0
242	3	3	2	35	6	40	47	69	76	97	85	95	80	78	105	42	48	56	12	4	1	15	2	0
243	0	0	0	0	18	6	70	47	130	146	115	21	62	64	247	42	22	4	2	0	0	0	1	0
244	22	22	18	16	38	65	86	87	74	83	68	64	61	34	32	51	105	25	17	10	2	2	6	12
245	6	6	5	7	16	30	26	53	78	126	75	74	33	44	63	118	131	12	8	2	68	8	8	4
246	0	0	0	0	7	426	80	147	29	25	23	109	2	29	53	6	45	0	0	0	0	17	0	0
247	0	0	5	175	1	6	0	37	49	13	4	11	250	0	1	0	439	0	0	0	0	0	0	0
248	4	4	12	8	64	229	105	285	61	59	32	42	10	71	3	4	8	0	0	0	0	0	0	0

Code	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
249	0	0	0	0	1	6	51	4	11	34	153	492	8	40	7	15	167	8	0	1	0	0	0	0
250	8	8	8	1	1	4	4	4	368	389	188	12	1	1	1	1	1	0	0	0	0	0	0	0
251	17	17	7	68	22	64	11	227	26	299	87	17	4	4	60	15	0	0	0	1	2	25	15	12
252	0	0	0	0	0	3	2	1	2	2	958	9	3	3	2	3	3	8	2	0	0	0	0	0
253	0	2	0	0	0	2	60	212	153	137	76	138	58	47	61	25	13	7	9	1	0	0	0	0
254	0	6	0	0	0	178	73	63	226	62	12	58	9	7	39	21	80	15	0	0	0	0	0	0
255	0	17	356	0	0	149	0	213	0	2	258	0	0	0	0	0	0	0	4	0	0	0	0	0
256	0	0	0	0	0	244	44	98	70	1	0	538	2	0	0	0	0	2	0	0	0	0	0	0
257	0	0	0	0	0	0	11	38	8	77	89	690	18	14	14	10	21	2	8	0	0	0	0	0
258	0	0	0	0	0	1	217	54	47	60	119	231	0	82	0	54	17	0	0	0	0	0	0	0
259	0	0	0	0	0	8	312	108	95	177	227	73	0	0	0	0	0	0	0	0	0	0	0	0
260	0	0	0	0	0	77	0	1	18	74	134	241	121	48	8	11	0	23	0	1	0	0	0	0
261	0	0	0	0	0	1	10	58	48	373	106	114	34	70	38	15	0	0	0	0	0	58	0	76
262	0	0	0	0	0	3	2	20	7	113	26	792	4	5	9	4	10	5	0	0	0	0	0	0
263	0	0	0	0	0	72	919	0	0	9	0	0	0	0	0	0	0	0	0	0	0	0	0	0
264	0	0	0	0	0	75	0	618	307	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
265	0	0	0	0	0	89	14	0	0	0	0	897	0	0	0	0	0	0	0	0	0	0	0	0
266	0	0	0	0	0	92	0	263	71	187	123	70	50	6	19	4	10	85	19	0	0	0	0	0
267	0	0	0	0	0	377	95	0	0	32	0	495	0	0	0	0	0	0	0	0	0	0	0	0
268	0	0	0	0	0	772	22	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
269	0	0	0	0	0	795	121	7	1	16	9	22	5	3	7	8	4	0	0	0	0	0	0	0
270	0	0	0	0	0	0	67	0	9	371	397	127	26	3	1	0	0	0	0	0	1	0	0	0
271	0	0	0	0	0	0	495	0	31	269	0	0	0	144	0	61	0	0	0	0	0	0	0	0
272	0	0	0	0	0	0	929	34	0	0	0	37	0	0	0	0	0	0	0	0	0	0	0	0
273	0	0	0	0	0	0	0	1	0	0	0	997	0	1	0	0	0	0	0	0	0	0	0	0
274	0	0	0	0	0	0	0	6	24	368	49	198	25	32	42	95	45	58	56	1	0	0	0	0
275	0	0	0	0	0	0	0	46	483	33	11	12	7	17	50	4	336	0	0	0	0	0	0	0
276	0	0	0	0	0	0	0	864	0	0	0	136	0	0	0	0	0	0	0	0	0	0	0	0
277	0	0	0	0	0	0	0	0	42	75	167	483	0	233	0	0	0	0	0	0	0	0	0	0
278	0	0	0	0	0	0	0	0	0	84	93	823	0	0	0	0	0	0	0	0	0	0	0	0
279	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
280	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1000	0	0	0	0	0	0	0
281	0	0	0	0	0	0	0	0	0	0	0	1000	0	0	0	0	0	0	0	0	0	0	0	0
282	0	0	0	0	0	0	0	1000	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
283	0	0	0	0	0	0	1000	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
284	0	0	0	0	0	1000	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

## APPENDIX C

### VMT Offset Demonstration for the 2008 Ozone Standard As Required by Clean Act Section 182(d)(1)(A)

## Appendix C VMT Offset Demonstration

### Appendix C.1 Background

In 1979, United States Environmental Protection Agency (USEPA) established a primary health-based national ambient air quality standard (NAAQS) for ozone at 0.12 parts per million (ppm) averaged over a 1-hour period (44 FR 8220). The Clean Air Act (CAA), as amended in 1990, classified areas that had not yet attained that standard, based on the severity of their ozone problem, ranging from Marginal to Extreme. Extreme areas were provided the most time to attain, until November 15, 2010, but were also subject to the most stringent requirements. In particular, Severe and Extreme areas were subject to CAA Section 182(d)(1)(A), which requires state implementation plans to adopt “specific enforceable transportation control strategies and transportation control measures to offset any growth in vehicle miles traveled or numbers of vehicle trips in such area....” USEPA designated the Sacramento Federal Nonattainment Area (SFNA) as Severe on April 25, 1995 (60 FR 20237), and thus the Sacramento Metropolitan area was subject to this requirement. The USEPA has historically interpreted this provision of the CAA (now called “Vehicle Miles Traveled (V T) emissions offset requirement”) to allow areas to meet the requirement by demonstrating that emissions from motor vehicles decline each year through the attainment year (57 FR 13521)

In 1997, USEPA replaced the 1-hour ozone standard with an 8-hour standard of 0.08 ppm (62 FR 38856). The USEPA promulgated rules implementing this standard with the “Phase 1” rule issued on April 30, 2004 (69 FR 23951), and the Phase 2 rule issued on November 29, 2005 (70 FR 71612). These implementation rules required that areas classified as Severe or Extreme under the 1997 8-hour standard would also be subject to the VMT offset requirement.

In 2008, USEPA revised the 8-hour ozone NAAQS to a level of 0.075 parts per million (73 FR16436). The SFNA was designated as non-attainment for the 2008 standard on May 21, 2012 and classified as severe (77 FR 30087), making the SFNA subject to the requirements of CAA Section 182(d)(1)(A) for the 2008 8-hour ozone NAAQS.

### Appendix C.2 USEPA Guidance on VMT Offset Requirement

In August 2012, USEPA issued guidance titled “Implementing Clean Air Act Section 182(d)(1)(A): Transportation Control Measures and Transportation Control Strategies to Offset Growth in Emissions due to Growth in Vehicle Miles Travelled”. Among other things, USEPA’s guidance states that both “transportation control measures” and “transportation control strategies” (TCS) are eligible to offset growth in emissions due to growth in VMT. The USEPA’s guidance indicates that TCSs, which are not defined in the CAA or USEPA regulation, include technology improvements such as vehicle technology improvements, motor vehicle fuels, motor vehicle inspection and

maintenance programs, and other control strategies that are transportation-related. USEPA's revised guidance sets forth a method to calculate the actual growth in VOC emissions due to growth in VMT. Essentially, the state would compare projected attainment year emissions assuming no new control measures and no VMT growth with projected actual attainment year emissions (including new control measures and VMT growth). If the first number is higher than the second, the new TCSs and TCS's are sufficient and no additional transportation control measures or strategies would be required. If the first number is lower, additional transportation control measures and transportation control strategies are required. As a practical matter, the state must add the measures and re-calculate emissions until it demonstrates that the TCSs and TCS's are sufficient to offset the growth in VMT. The new measures must be clearly identified and distinguished from the measures included in the initial calculations for the base year.

In addition, the guidance recommends that the base year used in the demonstration should be the base year used in the attainment demonstration for the ozone standard. To address USEPA's guidance, 2012 is used in this demonstration as the base year for the 2008 8-hour standard.

### **Appendix C.3 Transportation Control Strategies and Transportation Control Measures**

By listing them separately, the CAA Section 182(d)(1)(A) differentiates between transportation control strategies (TCS) and transportation control measures (TCM), and thus provides for a wide range of strategies and measures as options to offset growth in emissions from VMT growth. In addition, the example TCMs listed in CAA Section 108(f)(1)(A) include measures that reduce emissions by reducing VMT, reducing tailpipe emissions, and removing dirtier vehicles from the fleet. California's motor vehicle control program includes a variety of strategies and measures, including new engine standards and in-use programs (e.g., smog check, vehicle scrap, fleet rules, and idling restrictions). There were no local TCMs built into the transportation model used to generate emissions. The only TCM moving forward is spare the air. That program was also not built into the transportation model, but since it is implemented by the air district's, they opted to include it in the SIP.

Based on the provisions in CAA Section 182(d)(1)(A) and the clarifications provided in the USEPA guidance, any combination of transportation control strategies and TCMs may be used to meet the requirement to offset growth in emissions resulting from VMT growth. Since 1990 when this requirement was established, California has adopted more than sufficient enforceable transportation strategies and measures to meet the requirement to offset the growth in emissions from VMT growth. A list of the state's mobile source control program adopted since 1990 is provided as part of the Reasonably Available Control Measure Evaluation (Appendix E). Section 7.2, State and

Federal Control Measures, discusses how state and federal regulations will produce increasing emission reduction benefits from now until 2024 and beyond, as the regulated fleets are retrofitted, and as older and dirtier portions of the fleets are replaced with newer and cleaner models at an accelerated pace.

#### **Appendix C.4 Emissions due To VMT Growth**

As discussed above, the USEPA guidance provides a recommended calculation methodology to determine if sufficient TCSs and TCMs have been adopted and implemented to offset the growth in emissions due to growth in VMT. Any increase in emissions solely from VMT increases in the future attainment year from the base year would need to be offset. In addition, the EPA guidance recommends that the analysis include a calculation showing the emissions levels if VMT had remained constant from the base year to the future attainment year. As discussed earlier, the analysis compares the projected attainment year emissions assuming no new control measures and no VMT growth with projected actual attainment year emissions (including new control measures and VMT growth). If the second number is lower than the first, the new measures are adequate and no additional transportation control measures or strategies would be required.

#### **Appendix C.5 Methodology**

The following calculations are based on the USEPA guidance recommended calculation methodology. The attainment demonstration for the 8-hour ozone standard uses 2012 as the base year and 2024 is the attainment year.

##### Analysis Tool

This analysis uses California's approved motor vehicle emissions model, EMFAC.

The EMFAC model estimates the emissions from two combustion processes: running exhaust and start exhaust, and four evaporative processes: hot soak, running losses, diurnal, and resting losses.

Emissions from running exhaust, start exhaust, hot soak, and running losses are a function of how much a vehicle is driven. Emissions from these processes are directly related to VMT, trips, and starts. These processes are included in the calculation of the emissions levels used in the VMT offset demonstration. Emissions from resting loss and diurnal loss processes are not related to VMT, trips or vehicle starts and are not included in the analysis because these emissions occur whether or not the vehicle makes a trip (i.e., a start).

EMFAC combines trip-based VMT from the regional transportation planning agencies, starts data based on household travel surveys, and vehicle population data from the



California Department of Motor Vehicles with corresponding emission rates to calculate emissions<sup>1</sup>.

With the EMFAC model, the calculation of emissions growth and whether it is offset is simplified to a comparison of future year emissions with “no growth” in V T or new control strategies to future emissions with VMT growth and new control strategies. This follows USEPA’s 2012 guidance.

Analysis Using 2012 as the Base Year for the 2008 8-hour Ozone Standard with Attainment Year of 2024.

**Step 1. Provide the emissions level for the base year.**

The following table shows the VOC emissions, VMT, starts, and vehicle population for calendar year 2012 from the EMFAC2014 model.

**Summary of 2012 Base Year**

	VMT (thousand miles/day)	Starts (thousands/day)	Vehicle Population (thousands)	VOC Emissions* (tons/day)
2012 Base Year	60,570	11,739	1,849	28

\* Does not include diurnal or resting loss emissions.

**Step 2. Calculate three emissions levels in the attainment year.**

For the attainment year,

- (1) Calculate emissions level with the motor vehicle control program frozen at 2012 levels and with projected VMT, starts, and vehicle population for the attainment year. This represents what the emissions in the attainment year would have been if transportation control strategies and TCMs had not been implemented after 2012;
- (2) Calculate emissions level with the motor vehicle control program frozen at 2012 levels and assuming VMT, starts, and vehicle population do not increase from 2012 levels; and
- (3) Calculate an emissions level that represents emissions with full implementation of all transportation control strategies and TCMs pre- and post-2012 and the projected VMT, starts, and vehicle population for the attainment year.

---

<sup>1</sup> More information on data sources can be found in the EMFAC technical document which is located on the web at: <https://www.arb.ca.gov/msei/downloads/emfac2014/emfac2014-vol3-technical-documentation-052015.pdf>

**Calculation 1. Calculate the emissions in the attainment year assuming no new measures since the base year, and including growth in VMT, starts, and vehicle population.**

To perform this calculation, California Air Resources Board (CARB) staff identified the on-road motor vehicle control programs adopted since 2012 and adjusted EMFAC2014 to reflect the VOC emissions levels in 2024 without the benefits of the post-2012 control programs. The projected VOC emissions are 16 tons/day.

**Calculation 2. Calculate the emissions with no growth in VMT, starts, or vehicle population.**

In this calculation, the VOC emission levels in calendar year 2024 without benefit of the post 2012 control program are calculated. EMFAC2014 allows a user to input different VMT, starts, and vehicle population than default. For this calculation, EMFAC2014 was run without the benefit of the post 2012 control program for calendar year 2024 with the 2012 level of VMT of 60,569,748 miles per day, the 2012 level of starts at 11,739,339 per day, and the 2012 level of population at 1,849,178 vehicles. The VOC emissions associated with 2012 VMT, starts, and vehicle population in calendar year 2024 are 15 tons/day.

**Calculation 3. Calculate emission reductions with full Implementation of Transportation Control Strategies & TCMs.**

The VOC emission levels for 2024 assuming the benefits of the post-2012 motor vehicle control program and the projected VMT, starts, and vehicle population in 2024 are calculated using EMFAC2014. The projected VOC emissions level is 11 tons/day. VOC emissions for the three sets of calculations described above are summarized in the following table.

**Summary of 2024 Attainment Year Emissions Levels**

	<b>Description</b>	<b>VMT* (miles/day, thousands)</b>	<b>Starts (thousands/day)</b>	<b>Vehicle Population (thousands)</b>	<b>VOC Emissions** (tons/day)</b>
<b>(1)</b>	<b>Emissions with Motor Vehicle Control Program Frozen at 2012 Levels.</b> (VMT, starts and vehicle population at 2024 levels.)	69,579	11,965	1,939	16
<b>(2)</b>	<b>Emissions with Motor Vehicle Control Program Frozen at 2012 Levels.</b> (VMT, starts, and vehicle population at 2012 levels)	60,570	11,739	1,849	15
<b>(3)</b>	<b>Emissions with Full Motor Vehicle Control Program in Place.</b> (VMT, starts and vehicle population at 2024 levels)	69,579	11,965	1,939	11

\* CY 2024 VMT based on the SACOG 2016 MTP

\*\* Does not include diurnal or resting loss emissions.

As provided in the USEPA guidance, to determine compliance with the provisions of CAA Section 182(d)(1)(A), the emissions levels calculated in Calculation 3 should be less than the emissions levels in Calculation 2:


$$\text{VOC: } 11 < 15 \text{ tons/day}$$

**Appendix C.6 Summary**

The previous sections provide an analysis to demonstrate compliance with the provisions of CAA Section 182(d)(1)(A). To further illustrate the demonstration, Figure C-1 below shows graphically the emissions benefits of the motor vehicle control programs in offsetting VOC emissions due to increased VMT, starts, and vehicle population in the SFNA for the 2008 8-hour ozone standard (2012 base year). The left

bar (in purple) shows the emissions in the base year with base year controls. The three bars on the right in each figure show the emissions levels in the attainment year for the three calculations identified above: the red bar shows attainment year emissions with base year controls and attainment year VMT, starts, and vehicle population at 2024 levels (calculation 1), the green bar shows attainment year emissions with base year controls, VMT, starts, and vehicle population at 2012 levels (calculation 2), and the blue bar shows attainment year emissions with attainment year controls, VMT, starts, and vehicle population at 2024 levels (calculation 3). Based on the USEPA guidance, if the blue bar (calculation 3) is lower than the green bar (calculation 2), then the identified transportation control strategies and TCMs are sufficient to offset the growth in emissions.

**Figure C-1. VOC Emissions\* from On-Road Mobile Sources in the SFNA (2012 Base Year)**



## APPENDIX D

### Reasonable Further Progress Calculation

**Appendix D Reasonable Further Progress Calculation**

<b>Calculation of Reasonable Further Progress Demonstrations<sup>A</sup></b>				
<b>Sacramento Nonattainment Area</b>				
<b>Year</b>	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>
1. VOC (with existing measures from CEPAM 1.04) <sup>B</sup>	110.2	91.0	86.8	84.4
2. VOC ERCs <sup>C</sup>		5	5	5
3. VOC plus ERCs (Line 1+Line2)	110.2	96.0	91.8	89.4
4. Required % change since previous milestone year (VOC or Nox)		18%	9%	9%
5. Required % change since 2012 (VOC or Nox)		18%	27%	36%
6. Target VOC levels ((1-Line4)*previous milestone year Line6 (except 110.2 for 2018))		90.3	82.2	74.8
7. Shortfall (-)/Surplus (+) in VOC reductions needed to meet target (Line3 - Line6)		-5.7	-9.6	-14.6
8. Shortfall (-)/Surplus (+) in VOC reductions needed to meet target, % ((Line7)/110.2*100%)		-5.2%	-8.7%	-13.2%
9. VOC reductions since 2012 used for contingency in this milestone year, %		0%	0%	0%
10. VOC reductions shortfall previously provided by Nox substitution, % (sum of previous milestone year Line 20)		0%	5.2%	8.7%
11. Actual VOC reduction Shortfall (-)/Surplus (+), % (Line8 + Line10)		-5.2%	-3.5%	-4.5%
<b>Year</b>	<b>2012</b>	<b>2018</b>	<b>2021</b>	<b>2024</b>
12. NOx (with existing measures from CEPAM 1.04) <sup>B</sup>	101.1	69.4	58.4	48.8
13. NOx ERCs <sup>C</sup>		4	4	4
14. NOx Safety Margin - Transportation Conformity Emissions Budgets <sup>D</sup>		0	0.5	0
15. NOx plus ERCs and Safety Margin (Line 12+Line13+Line14)	101.1	73.4	62.4	52.8
16. Change in Nox since 2012 (101.1 - Line15)		27.7	38.8	48.4
17. Change in Nox since 2012, % (Line16/101.1*100%)		27.4%	38.4%	47.8%
18. NOx reductions since 2012 already used for VOC substitution and contingency through last milestone year, % (previous milestone year(Line18+Line20+Line21))		0%	8.2%	11.7%
19. NOx reductions since 2012 available for VOC substitution and contingency in this milestone year, % (Line17 - Line18)		27.4%	30.2%	36.2%
20. NOx reductions since 2012 used for VOC substitution in this milestone year, % (0-Line11)		5.2%	3.5%	4.5%
21. NOx reductions since 2012 used for contingency in this milestone year, %		3%	0%	0%
22. NOx reductions since 2012 surplus after meeting VOC substitution and contingency needs in this miles year, % (Line19 - Line20 - Line21)		19.2%	26.7%	31.6%
23. RFP shortfall (-) in reductions needed to meeet target, if any, %		0%	0%	0%
24. Total shortfall (-) for RFP and Contingency, if any, %		0%	0%	0%
25. RFP Met?		YES	YES	YES
26. Contingency Met?		YES	YES	YES
<sup>A</sup> CARB RFP write-up September 8, 2016, email transmittal to SMAQMD with safety margin of 0.5 tpd NOx in 2021 for Transportation Conformity.				
<sup>B</sup> VOC and NOx are from CEPAM 2016 Ozone SIP forecast for SFNA, Version 1.04 with approved external adjustments.				
<sup>C</sup> ERCs from Chapter 5, Section 5.6: VOC= 5 tpd, NOx = 4 tpd.				
<sup>D</sup> Safety Margin of 0.5 tpd NOx in 2021 for Transportation Conformity Emissions Budgets is from Table 10-1.				

## APPENDIX E

### Reasonably Available Control Measure Analysis

## Appendix E Reasonably Available Control Measure Analysis

### Appendix E.1 RACM requirements

This Appendix describes the Reasonably Available Control Measure (RACM) analysis that was conducted for the Sacramento Federal Nonattainment Area (SFNA). This analysis complies with Clean Air Act (CAA) Section 172(c)(1) which requires a nonattainment plan to:

*“provide for the implementation of all reasonably available control measures as expeditiously as practicable (including such reductions in emissions from existing sources in the area as may be obtained through the adoption, at a minimum, of reasonably available control technology) and shall provide for attainment of the national primary ambient air quality standards.”*

United States Environmental Protection Agency’s (USEPA) RAC policy (40 FR 122 2-12283; USEPA, 1999) indicates that nonattainment areas “should consider all available measures that are potentially reasonably available”. Sources of potentially reasonable measures include measures adopted in other nonattainment areas and measures that the USEPA has identified in guidelines or other documents.

Areas should consider all reasonably available measures for implementation in light of local circumstances. However, areas are only required to adopt measures if they are economically and technologically feasible and (alone or cumulatively) will advance the attainment date by one year or more, or are necessary for reasonable further progress (RFP)(80 FR 12282). EPA “does not believe that Congress intended the RACM requirement to compel the adoption of measures that are absurd, unenforceable, or impracticable.” (57 FR 1349 )

### Appendix E.2 Process of identifying RACM

To identify all RACM, District staff reviewed multiple sources of control measure information, including:

- Control measures included in the attainment plan for the 1997 8-hour National Ambient Air Quality Standard (NAAQS)(SMAQMD, et al, 2013)
- Rules adopted or amended between January 2006 and July 2013 in the Bay Area Air Quality Management District (BAAQMD), South Coast Air Quality Management District (SCAQMD), San Diego Air Pollution Control District (SDAPCD), San Joaquin Valley Unified Air Pollution Control District (SJVUAPCD), and Ventura County Air Pollution Control District (VCAPCD);
- USEPA’s Reasonably Available Control Technology (RACT)/ Best Available Control Technology (BACT)/ Lowest achievable Emission Rate (LAER) Clearinghouse;



- California Air Resources Board's (CARB's) BACT Clearinghouse;
- BAAQ's 2010 Clean Air Plan;
- SCAQ's 2012 Air Quality Management Plan; and
- Rules from other areas of the nation with similar nonattainment status, including Houston-Galveston-Brazoria, TX; Dallas-Fort Worth, TX; and Baltimore, MD.

Staff from each of the five air districts in the SFNA performed the RACM analysis for the stationary and areawide sources in their jurisdictions. For each potential RACM measure, the emissions inventory, emissions reductions, and cost effectiveness were estimated.

### Appendix E.3 Conclusion

The District evaluated and analyzed all reasonable control measures that were currently available for inclusion in the plan. The control measures evaluated for inclusion in this plan also include mobile source measures provided by CARB, and Transportation Control Measures (TCMs) provided by Sacramento Area Council of Governments (SACOG).<sup>1</sup>

The RACMs collectively would not advance the attainment date or contribute to RFP for the Sacramento region because of the insufficient or non-quantifiable amount of emission reductions that they may potentially generate. Tables E1 through E5 contain a list of the measures evaluated by each of the five air districts and a brief discussion of the conclusions. The RACM demonstration for transportation control measures was prepared by the SACOG and is included in Table E6. CARB analyzed mobile measures for RACM purposes and included a written description of that analysis on page E-39.

---

<sup>1</sup> A RACM analysis for TC's (Sierra Research, 2015) was completed by SACOG. This analysis summarized the ozone SIP requirements, documents the TCM identification process, and also provided preliminary RACM determination specific to SACOG. This is also discussed in Chapter 7 (Control Measures).

**Appendix E.4 Sacramento Metropolitan Air Quality Management District  
(SMAQMD)**

Table E-1 SMAQMD Stationary/Area Source Control Measures Considered

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
460	Adhesives and Sealants	VOC limits on adhesives and sealants	Reduce VOC limits on adhesives and sealants similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
442	Architectural Coatings	VOC limits on coatings	Reduce the VOC limits on architectural coatings similar to the rule adopted by SCAQMD	Not Recommended - Technical feasibility not demonstrated outside of SCAQMD
	Asphaltic Concrete	None	Establish NO <sub>x</sub> emission standards for aggregate dryers similar to the rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
459	Automotive Refinishing	VOC limits on coatings	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
411	Boilers	NO <sub>x</sub> limits on boiler/steam generators with a rated heat input capacity of 1 mmBtu/hr or greater	Reduce NO <sub>x</sub> limits similar to SCAQMD and SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Brandy and Wine Aging	None	Establish VOC emissions standards to reduce evaporative VOC emissions from the fermentation process at distilleries and wineries similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
452	Can Coating	VOC limits on coatings	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Commercial Cooking	VOC emission standards for large commercial bread bakeries	Establish standards to control VOC emissions from commercial charbroilers similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Composting Operations	None	Establish work practice requirements to reduce VOC emissions from green waste composting similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
496	Confined Animal Facilities	Implement VOC emission mitigation measures from a menu of options	Reduce animal-count applicability thresholds; increase number of mitigation measures, and control efficiency	Not Recommended - Evaluated for Attainment Advancement
	Flares	None	Establish NO <sub>x</sub> emission standards for flares similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Furnaces (Residential)	None	Establish point-of-sale NO <sub>x</sub> emissions standard for natural gas-fired central furnaces similar to SCAMQD requirements	Not Recommended - Evaluated for Attainment Advancement
	Further Control of High-Emitting Spray Booth Facilities	None	Require additional controls to reduce VOC emissions from spray booths at facilities emitting > 20 tons per year	Not Recommended - No sources
446/447/448	Gasoline Storage, Loading, and Degassing of Tanks and Pipelines	VOC emission standards for organic liquid storage tanks; vapor-recovery requirements for loading at bulk plants and bulk terminals	Reduce VOC emission limits for gasoline loading at bulk plants and bulk terminals to be as stringent as BAAQMD; establish VOC emission standards for degassing storage tanks and pipelines similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Glass Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for glass melting furnaces	Not Recommended - No sources
450	Graphic Arts	VOC limits on inks, coatings, adhesives or use emission control system	Reduce VOC limits for flexographic ink on porous substrates, extreme performance ink, and metallic ink to be as stringent as SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
412	Internal Combustion (IC) Engines	NO <sub>x</sub> emission limits on IC engines located at major stationary sources of NO <sub>x</sub>	Reduce NO <sub>x</sub> limits to be stringent as SCAQMD; expand applicability to include non-major stationary sources of NO <sub>x</sub>	Not Recommended - Evaluated for Attainment Advancement
464	Industrial Wastewater	Requirements for covers and emission control systems for wastewater collection and treatment systems at organic chemical plants	Lower applicability thresholds to require controls on more wastewater streams, increase required efficiency of VOC control devices similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Liquefied Petroleum Gas (LPG) Transfer and Dispensing	None	Establish standards to control VOC emissions from LPG transfer and dispensing similar to the rules adopted by SCAQMD and VCAPCD	Not Recommended - Evaluated for Attainment Advancement
	Metal Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for metal melting furnaces	Not Recommended - No sources
451	Metal Parts and Products Coating	VOC limits on coatings, strippers, cleaning solvents	Reduce VOC limits for general one-component, extreme high gloss, and prefabricated architectural coatings, similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Metal Working Fluids	None	Establish VOC limits on metalworking fluids and direct-contact lubricants similar to the rules adopted by SCAQMD and VCAPCD	Not Recommended - Evaluated for Attainment Advancement
440	Miscellaneous Coatings	None	Establish VOC limits and application method requirements for coating operations not covered by other rules, similar to SCAQMD, SJVUAPCD, VCAPCD, and BAAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Miscellaneous Combustion Sources	None	Establish NO <sub>x</sub> emission limits on miscellaneous combustion equipment including dryers and ovens similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Mold Release Agents	None	Establish VOC limits on mold release agents similar to the control measure proposed by SCAQMD	Not Recommended – Has not yet been implemented in SCAQMD or any other area
485	Municipal Landfill Gas	Landfill gas collection and control systems	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
	Oil and Natural Gas Production	None	Establish requirements to inspect and maintain equipment to reduce fugitive VOC emissions	Not Recommended - Evaluated for Attainment Advancement
407/501	Open Burning	Burning of certain materials prohibited; burn procedures to minimize smoke; burning is not allowed on days declared no-burn day	Reduce the types of allowable agricultural burns similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Paper, Fabric, and Film Coatings	None	Establish VOC limits on coatings similar to rule adopted by SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
444	Petroleum Solvent Dry Cleaning	Emit no more than 3.5 kg of solvent per 100,000 articles dry cleaned or use a solvent recovery dryer	Expand applicability to include all non-halogenated solvents; require closed-loop machines for new installations	Not Recommended - Evaluated for Attainment Advancement
	Plastic Parts Coating	None	Establish VOC limits on plastic parts coatings similar to rule adopted by SCAMQD	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
465	Polyester Resin/Plastic Product Manufacturing	Limits on the monomer content of resin, use of vapor suppressants, use of close-mold systems, or emission capture and control system	Remove low-usage exemption, require non-atomizing equipment, and reduce monomer content similar to rules adopted by BAAQMD, SCAQMD, and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
	Polystyrene /Polymeric Cellular (Foam) Manufacturing	None	Require reduction of VOC emissions from Expanded Polystyrene (EPS) molding using an emission control device	Not Recommended – No sources
	Portland Cement Manufacturing	None	Establish NO <sub>x</sub> limits for Portland cement manufacturing	Not Recommended – No sources
	Semiconductor Manufacturing	None	Establish VOC limits for semiconductor manufacturing	Not Recommended – No sources
443	Synthetic Organic Chemical Manufacturing – Fugitive Leaks	Leak detection and repair program	Reduce VOC leak detection threshold	Not Recommended - Evaluated for Attainment Advancement
	Soil Decontamination	None	Establish VOC emission control standards for soil vapor extraction systems, similar to rules adopted by BAAQMD and VCAPCD; Establish work practices to minimize VOC emissions from soil aeration similar to rule adopted by SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
454/466	Solvent Cleaning	VOC limits on solvents, or use airtight/airless cleaning systems	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
413	Stationary Gas Turbines	NO <sub>x</sub> emission limits on stationary gas turbines	Reduce NO <sub>x</sub> emission limits to be as stringent as SCAQMD	Not Recommended - Evaluated for Attainment Advancement

<b>Measure No.</b>	<b>Title</b>	<b>Current Requirements</b>	<b>Opportunity for Strengthening</b>	<b>Conclusion</b>
	Wastewater Separators	None	Require solid cover, floating pontoon cover; double-deck cover, or vapor recovery system similar to rule adopted by SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
414	Water Heaters and Small Boilers	Point-of-sale NO <sub>x</sub> emission standards on water heaters with rated heat input capacity less than 1 mmBtu/hr	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
463	Wood Products Coatings	VOC limits on coatings	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement

**Appendix E.5 El Dorado County Air Quality Management District (EDCAQMD)**

Table E-2 EDCAQMD Stationary/Area Source Control Measures Considered

<b>Measure No.</b>	<b>Title</b>	<b>Current Requirements</b>	<b>Opportunity for Strengthening</b>	<b>Conclusion</b>
236	Adhesives and Sealants	VOC limits on adhesives and sealants	Reduce VOC limits on adhesives and sealants similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
215	Architectural Coatings	VOC limits on coatings	Reduce the VOC limits on architectural coatings similar to the SCM and rule adopted by SCAQMD	Not Recommended – SCM Evaluated for Attainment Advancement. Technical feasibility of SCAQMD requirements not demonstrated outside of SCAQMD
	Asphaltic Concrete	None	Establish NO <sub>x</sub> emission standards for aggregate dryers similar to the rules adopted by SCAQMD and SJVUAPCD	Not Recommended – No Sources
230	Automotive Refinishing	VOC limits on coatings	Reduce the VOC limits on architectural coatings consistent with the SCM	Not Recommended - Evaluated for Attainment Advancement
229	Boilers	NO <sub>x</sub> limits on boiler/steam generators with a rated heat input capacity of 5 mmBtu/hr or greater	Expand applicability to units ≥ 2 mmBtu/hr and reduce NO <sub>x</sub> limits similar to SCAQMD and SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Brandy and Wine Aging	None	Establish VOC emissions standards to reduce evaporative VOC emissions from the fermentation process at distilleries and wineries similar to SJVUAPCD requirements	Not Recommended – No sources
	Can Coating	None	Establish VOC limits on can coatings similar to rule adopted by SMAQMD	Not Recommended – No sources



Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Commercial Cooking	None	Establish standards to control VOC emissions from commercial charbroilers similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Composting Operations	None	Establish work practice requirements to reduce VOC emissions from green waste composting similar to SCAQMD requirements	Not Recommended - No sources
	Confined Animal Facilities	None	Establish work practice requirements to reduce VOC emissions from confined animal facilities	Not Recommended - No sources
	Flares	None	Establish NO <sub>x</sub> emission standards for flares similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Furnaces (Residential)	None	Establish point-of-sale NO <sub>x</sub> emissions standard for natural gas-fired central furnaces similar to SCAMQD requirements	Not Recommended - Evaluated for Attainment Advancement
	Further Control of High-Emitting Spray Booth Facilities	None	Require additional controls to reduce VOC emissions from spray booths at facilities emitting > 20 tons per year	Not Recommended - No sources
216/244	Organic Liquid Storage, Loading, and Degassing of Tanks and Pipelines, Bulk Plant Terminals	VOC emission standards for organic liquid storage tanks; vapor-recovery requirements for loading at bulk plants and bulk terminals	Reduce VOC emission limits for gasoline loading at bulk plants and bulk terminals to be as stringent as BAAQMD; establish VOC emission standards for degassing storage tanks and pipelines similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Glass Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for glass melting furnaces	Not Recommended - No sources
231	Graphic Arts	VOC limits on inks, coatings, adhesives or use emission control system	Reduce VOC limits for flexographic ink on porous substrates, extreme performance ink, and metallic ink to be as stringent as SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
233	IC Engines	NO <sub>x</sub> limits on IC Engines	Reduce NO <sub>x</sub> limits for IC engines similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
464	Industrial Wastewater	None	Establish emission control standards for wastewater systems	Not Recommended - No sources
	Metal Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for metal melting furnaces	Not Recommended - No sources
	Metal Parts and Products Coating	None	Establish VOC limits on metal parts and products coating similar to SMAQMD and SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Metal Working Fluids	None	Establish VOC limits on metalworking fluids and direct-contact lubricants similar to the rules adopted by SCAQMD and VCAPCD	Not Recommended - Evaluated for Attainment Advancement
	Miscellaneous Coating	None	Establish VOC limits and application method requirements for coating operations not covered by other rules, similar to SCAQMD, SJVUAPCD, VCAPCD, and BAAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Miscellaneous Combustion Sources	None	Establish NO <sub>x</sub> emission limits on miscellaneous combustion equipment including dryers and ovens similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
	Mold Release Agents	None	Establish VOC limits on mold release agents similar to the control measure proposed by SCAQMD	Not Recommended - Has not yet been implemented in SCAQMD or any other area
	Municipal Landfill Gas	None	Establish requirements for landfills including gas collection and control systems	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Oil and Natural Gas Production	None	Establish requirements to inspect and maintain equipment to reduce fugitive VOC emissions	Not Recommended – No sources
300	Open Burning	Burning of certain materials prohibited; burn procedures to minimize smoke; burning is not allowed on days declared no-burn day	Reduce the types of allowable agricultural burns similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Paper, Fabric, and Film Coatings	None	Establish VOC limits on coatings similar to rule adopted by SJVUAPCD	Not Recommended – No sources
218	Petroleum Solvent Dry Cleaning	Emit no more than 0.6 kg of solvent per kg of wet waste or use a system that reduces waste losses below 0.01 kg per kg of clothes	Remove applicability threshold to include all dry cleaning solvents except for perchloroethylene and ban the use of open transfer systems	Not Recommended - Evaluated for Attainment Advancement
	Plastic Parts Coating	None	Establish VOC limits on plastic parts coatings similar to rule adopted by SCAMQD	Not Recommended - Evaluated for Attainment Advancement
	Polyester Resin/Plastic Product Manufacturing	None	Establish VOC standards on monomer content of resins and require vapor suppressants and use of close-mold systems similar to rules adopted by BAAQMD, SCAQMD, and SJVUAPCD	Not Recommended – No sources
	Polystyrene /Polymeric Cellular (Foam) Manufacturing	None	Require reduction of VOC emissions from EPS molding using an emission control device	Not Recommended - Evaluated for Attainment Advancement
	Portland Cement Manufacturing	None	Establish NO <sub>x</sub> limits for Portland cement manufacturing	Not Recommended – No sources
	Semiconductor Manufacturing	None	Establish VOC limits for semiconductor manufacturing	Not Recommended – No sources

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Synthetic Organic Chemical Manufacturing – Fugitive Leaks	None	Establish VOC emissions standards for leak detection and repair program	Not Recommended – No sources
	Soil Decontamination	None	Establish VOC emission control standards for soil vapor extraction systems, similar to rules adopted by BAAQMD and VCAPCD; Establish work practices to minimize VOC emissions from soil aeration similar to rule adopted by SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
225/235	Solvent Cleaning	VOC limits on solvents	Reduce VOC limits of solvents similar to rules adopted by SMAQMD and PCAPCD.	Not Recommended - Evaluated for Attainment Advancement
	Stationary Gas Turbines	None	Establish NO <sub>x</sub> emission limits to be as stringent as SCAQMD	Not Recommended – No sources
	Wastewater Separators	None	Require solid cover, floating pontoon cover; double-deck cover, or vapor recovery system similar to rule adopted by SJVUAPCD	Not Recommended – No sources
239	Water Heaters and Small Boilers	Point-of-sale NO <sub>x</sub> emission standards on water heaters with rated heat input capacity less than 75,000 Btu/hr	Expand point-of-sale emission standards to include units ≥ 75,000 Btu/hr and < 1 mmBtu/hr similar to rule adopted by SMAQMD	Not Recommended - Evaluated for Attainment Advancement
237	Wood Products Coatings	VOC limits on coatings	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement

**Appendix E.6 Feather River Air Quality Management District (FRAQMD)**

**Table E-3 FRAQMD Stationary/Area Source Control Measures Considered**

<b>Measure No.</b>	<b>Title</b>	<b>Current Requirements</b>	<b>Opportunity for Strengthening</b>	<b>Conclusion</b>
	Adhesives and Sealants	None	Establish VOC limits on adhesives and sealants similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
3.15	Architectural Coatings	VOC limits on coatings	Reduce the VOC limits on architectural coatings similar to the rule adopted by SCAQMD	Not Recommended – Technical feasibility not demonstrated outside of SCAQMD
	Asphaltic Concrete	None	Establish NO <sub>x</sub> standards similar to the rules adopted by SCAQMD/SJVUAPCD	Not Recommended – No sources
3.19	Automotive Refinishing	VOC limits on coatings	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
3.21	Boilers	NO <sub>x</sub> limits on boiler/steam generators with a rated heat input capacity of 1 mm Btu/hr or greater	Reduce NO <sub>x</sub> limits similar to SCAQMD and SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Brandy and Wine Aging	None	Establish VOC emissions standards to reduce evaporative VOC emissions from the fermentation process at distilleries and wineries similar to SJVUAPCD requirements	Not Recommended – No sources
	Can Coating	None	Establish VOC limits on can coatings similar to rule adopted by SMAQMD	Not Recommended – No sources
	Commercial Cooking	None	Establish standards to control VOC emissions from commercial charbroilers similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Composting Operations	None	Establish work practice requirements to reduce VOC emissions from green waste composting similar to SCAQMD requirements	Not Recommended – No sources
	Confined Animal Facilities	None	Establish work practice requirements to reduce VOC emissions from confined animal facilities	Not Recommended – No sources
	Flares	None	Establish NO <sub>x</sub> emission standards for flares similar to SJVUAPCD requirements	Not Recommended – No sources
	Furnaces (Residential)	None	Establish point-of-sale NO <sub>x</sub> emissions standard for natural gas-fired central furnaces similar to SCAMQD requirements	Not Recommended - Evaluated for Attainment Advancement
	Further Control of High-Emitting Spray Booth Facilities	None	Require additional controls to reduce VOC emissions from spray booths at facilities emitting > 20 tons per year	Not Recommended – No sources
3.9	Gasoline Storage, Loading, and Degassing of Tanks and Pipelines	VOC emission standards for organic liquid storage tanks; vapor-recovery requirements for loading at bulk plants and bulk terminals	Reduce VOC emission limits for gasoline loading at bulk plants and bulk terminals to be as stringent as BAAQMD; establish VOC emission standards for degassing storage tanks and pipelines similar to SCAQMD requirements	Not Recommended – No sources
	Glass Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for glass melting furnaces	Not Recommended – No sources
	Graphic Arts	None	Establish VOC limits on inks, coatings, or adhesives for graphic arts similar to SJVUAPCD requirements	Not Recommended – No sources
3.22	IC Engines	NO <sub>x</sub> limits on IC Engines	Reduce NO <sub>x</sub> limits for IC engines similar to SCAQMD requirements	Not Recommended – No sources
	Industrial Wastewater	None	Establish emission control standards for wastewater systems	Not Recommended – No sources

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	LPG Transfer and Dispensing	None	Establish standards to control VOC emissions from LPG transfer and dispensing similar to the rules adopted by SCAQMD and VCAPCD	Not Recommended - Evaluated for Attainment Advancement
	Metal Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for metal melting furnaces	Not Recommended - No sources
	Metal Parts and Products Coating	None	Establish VOC limits on metal parts and products coating similar to SMAQMD and SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Metal Working Fluids	None	Establish VOC limits on metalworking fluids and direct-contact lubricants similar to the rules adopted by SCAQMD and VCAPCD	Not Recommended - Evaluated for Attainment Advancement
	Miscellaneous Coating	None	Establish VOC limits and application method requirements for coating operations not covered by other rules, similar to SCAQMD, SJVUAPCD, VCAPCD, and BAAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Miscellaneous Combustion Sources	None	Establish NO <sub>x</sub> emission limits on miscellaneous combustion equipment including dryers and ovens similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
	Mold Release Agents	None	Establish VOC limits on mold release agents similar to the control measure proposed by SCAQMD	Not Recommended - Has not yet been implemented in SCAQMD or any other area
3.18	Municipal Landfill Gas	Landfill gas collection and control systems	No control strategies identified	Not Recommended - No sources
	Oil and Natural Gas Production	None	Establish requirements to inspect and maintain equipment to reduce fugitive VOC emissions	Not Recommended - No sources

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
Reg. II	Open Burning	Burning of certain materials prohibited; burn procedures to minimize smoke; burning is not allowed on days declared no-burn day	Reduce the types of allowable agricultural burns similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Paper, Fabric, and Film Coatings	None	Establish VOC limits on coatings similar to rule adopted by SJVUAPCD	Not Recommended - No sources
	Petroleum Solvent Dry Cleaning	None	Establish VOC limits on solvents used and ban the use of open transfer systems	Not Recommended - No sources
	Plastic Parts Coating	None	Establish VOC limits on plastic parts coatings similar to rule adopted by SCAMQD	Not Recommended - No sources
	Polyester Resin/Plastic Product Manufacturing	None	Establish VOC standards on monomer content of resins and require vapor suppressants and use of close-mold systems similar to rules adopted by BAAQMD, SCAQMD, and SJVUAPCD	Not Recommended - No sources
	Polystyrene /Polymeric Cellular (Foam) Manufacturing	None	Require reduction of VOC emissions from EPS molding using an emission control device	Not Recommended - No sources
	Portland Cement Manufacturing	None	Establish NO <sub>x</sub> limits for Portland cement manufacturing	Not Recommended - No sources
	Semiconductor Manufacturing	None	Establish VOC limits for semiconductor manufacturing	Not Recommended - No sources
	Synthetic Organic Chemical Manufacturing – Fugitive Leaks	None	Establish VOC emissions standards for leak detection and repair program	Not Recommended - Evaluated for Attainment Advancement



Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Soil Decontamination	None	Establish VOC emission control standards for soil vapor extraction systems, similar to rules adopted by BAAQMD and VCAPCD; Establish work practices to minimize VOC emissions from soil aeration similar to rule adopted by SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
3.14	Solvent Cleaning	VOC limits on solvents	Reduce VOC limits of solvents similar to rules adopted by SMAQMD and PCAPCD.	Not Recommended - Evaluated for Attainment Advancement
	Stationary Gas Turbines	None	Establish NO <sub>x</sub> emission limits to be as stringent as SCAQMD	Not Recommended - No sources
	Wastewater Separators	None	Require solid cover, floating pontoon cover; double-deck cover, or vapor recovery system similar to rule adopted by SJVUAPCD	Not Recommended - No sources
3.23	Water Heaters and Small Boilers	None	Establish point-of-sale emission standards for include units < 1 mmBtu/hr similar to rule adopted by SMAQMD	Not Recommended - Evaluated for Attainment Advancement
3.20	Wood Products Coatings	VOC limits on coatings	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement

**Appendix E.7 Placer County Air Pollution Control District (PCAPCD)**

**Table E-4 PCAPCD Stationary/Area Source Control Measures Considered**

<b>Measure No.</b>	<b>Title</b>	<b>Current Requirements</b>	<b>Opportunity for Strengthening</b>	<b>Conclusion</b>
235	Adhesives and Sealants	VOC limits on adhesives and sealants	Reduce VOC limits on adhesives and sealants similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
218	Architectural Coatings	VOC limits on coatings	Reduce the VOC limits on architectural coatings similar to rule adopted by SCAQMD	Not Recommended - Technical feasibility not demonstrated outside of SCAQMD
	Asphaltic Concrete	None	Establish NO <sub>x</sub> emission standards for aggregate dryers similar to the rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
234	Automotive Refinishing	VOC limits on coatings	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
231/247	Boilers	NO <sub>x</sub> limits on boiler/steam generators with a rated heat input capacity of 5 mmBtu/hr or greater	Expand applicability to units ≥ 2 mmBtu/hr and reduce NO <sub>x</sub> limits similar to SCAQMD and SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Brandy and Wine Aging	None	Establish VOC emissions standards to reduce evaporative VOC emissions from the fermentation process at distilleries and wineries similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Can Coating	None	Establish VOC limits on can coatings similar to rule adopted by SMAQMD	Not Recommended - No sources
	Commercial Cooking	None	Establish standards to control VOC emissions from commercial charbroilers similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Composting Operations	None	Establish work practice requirements to reduce VOC emissions from green waste composting similar to SCAQMD requirements	Not Recommended – No sources
	Confined Animal Facilities	None	Establish work practice requirements to reduce VOC emissions from confined animal facilities	Not Recommended – No sources
	Flares	None	Establish NO <sub>x</sub> emission standards for flares similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Furnaces (Residential)	None	Establish point-of-sale NO <sub>x</sub> emissions standard for natural gas-fired central furnaces similar to SCAMQD requirements	Not Recommended - Evaluated for Attainment Advancement
	Further Control of High-Emitting Spray Booth Facilities	None	Require additional controls to reduce VOC emissions from spray booths at facilities emitting > 20 tons per year	Not Recommended – No sources
212/215	Storage of Organic Liquids and Transfer of Gasoline into Tank Trucks, Trailers, and Railroad Tank Cars at Loading Facilities	VOC emission standards for organic liquid storage tanks; vapor-recovery requirements for loading at bulk plants and bulk terminals	Reduce VOC emission limits for gasoline loading at bulk plants and bulk terminals to be as stringent as BAAQMD; establish VOC emission standards for degassing storage tanks and pipelines similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Glass Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for glass melting furnaces	Not Recommended – No sources
239	Graphic Arts	VOC limits on inks, coatings, adhesives or use emission control system	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
242	IC Engines	NO <sub>x</sub> emission limits on IC engines located at stationary sources of NO <sub>x</sub>	Reduce NO <sub>x</sub> limits to be stringent as SCAQMD	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Industrial Wastewater	None	Establish emission control standards for wastewater systems	Not Recommended – No sources
	LPG Transfer and Dispensing	None	Establish standards to control VOC emissions from LPG transfer and dispensing similar to the rules adopted by SCAQMD and VCAPCD	Not Recommended - Evaluated for Attainment Advancement
	Metal Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for metal melting furnaces	Not Recommended – No sources
245	Metal Parts and Products Coating	VOC limits on coatings, strippers, and solvent cleaner	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
	Metal Working Fluids	None	Establish VOC limits on metalworking fluids and direct-contact lubricants similar to the rules adopted by SCAQMD and VCAPCD	Not Recommended - Evaluated for Attainment Advancement
	Miscellaneous Coating	None	Establish VOC limits and application method requirements for coating operations not covered by other rules, similar to SCAQMD, SJVUAPCD, VCAPCD, and BAAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Miscellaneous Combustion Sources	None	Establish NO <sub>x</sub> emission limits on miscellaneous combustion equipment including dryers and ovens similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
	Mold Release Agents	None	Establish VOC limits on mold release agents similar to the control measure proposed by SCAQMD	Not Recommended – Has not yet been implemented in SCAQMD or any other area
	Municipal Landfill Gas	None	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Oil and Natural Gas Production	None	Establish requirements to inspect and maintain equipment to reduce fugitive VOC emissions	Not Recommended – No sources
301-306	Open Burning	Burning of certain materials prohibited; burn procedures to minimize smoke; burning is not allowed on days declared no-burn day	Reduce the types of allowable agricultural burns similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Paper, Fabric, and Film Coatings	None	Establish VOC limits on coatings similar to rule adopted by SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
	Petroleum Solvent Dry Cleaning	None	Establish VOC limits on solvents used and ban the use of open transfer systems	Not Recommended - Evaluated for Attainment Advancement
249	Plastic Parts Coating	VOC limits on coatings	Reduce VOC limits on plastic parts coatings similar to rule adopted by SCAMQD	Not Recommended - Evaluated for Attainment Advancement
243	Polyester Resin/Plastic Product Manufacturing	Limits on the monomer content of resin, use of vapor suppressants	Remove low-usage exemption, require non-atomizing equipment, and reduce monomer content similar to rules adopted by BAAQMD, SCAQMD, and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
	Polystyrene /Polymeric Cellular (Foam) Manufacturing	None	Require reduction of VOC emissions from EPS molding using an emission control device	Not Recommended – No sources
	Portland Cement Manufacturing	None	Establish NO <sub>x</sub> limits for Portland cement manufacturing	Not Recommended – No sources
244	Semiconductor Manufacturing	VOC limits on semiconductor manufacturing	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Synthetic Organic Chemical Manufacturing – Fugitive Leaks	None	Establish VOC emissions standards for leak detection and repair program	Not Recommended – No sources
	Soil Decontamination	None	Establish VOC emission control standards for soil vapor extraction systems, similar to rules adopted by BAAQMD and VCAPCD; Establish work practices to minimize VOC emissions from soil aeration similar to rule adopted by SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
216/240	Solvent Cleaning	VOC limits on solvents	Reduce VOC limits for solvents similar to rule adopted by SCAQMD	Not Recommended - Evaluated for Attainment Advancement
250	Stationary Gas Turbines	NO <sub>x</sub> limits on stationary gas turbines	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
	Wastewater Separators	None	Require solid cover, floating pontoon cover; double-deck cover, or vapor recovery system similar to rule adopted by SJVUAPCD	Not Recommended – No sources
246	Water Heaters and Small Boilers	None	Establish point-of-sale NO <sub>x</sub> emission standards on water heaters with rated heat input capacity less than 1 mmBtu/hr	Not Recommended - Evaluated for Attainment Advancement
236	Wood Products Coatings	VOC limits on coatings	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement

**Appendix E.8 Yolo-Solano Air Quality Management District (YSAQMD)**

Table E-5 YSAQMD Stationary/Area Source Control Measures Considered

<b>Measure No.</b>	<b>Title</b>	<b>Current Requirements</b>	<b>Opportunity for Strengthening</b>	<b>Conclusion</b>
2.33	Adhesives and Sealants	VOC limits on adhesives and sealants	Reduce VOC limits on adhesives and sealants similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
2.14	Architectural Coatings	VOC limits on coatings	Reduce the VOC limits on architectural coatings similar to the rule adopted by SCAQMD	Not Recommended - Technical feasibility not demonstrated outside of SCAQMD
	Asphaltic Concrete	None	Establish NO <sub>x</sub> emission standards for aggregate dryers similar to the rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
2.26	Automotive Refinishing	VOC limits on coatings	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
2.27	Boilers	NO <sub>x</sub> limits on boiler/steam generators with a rated heat input capacity of 5 mmBtu/hr or greater	Expand applicability to units ≥ 2 mmBtu/hr and reduce NO <sub>x</sub> limits similar to SCAQMD and SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
4695	Brandy and Wine Aging	None	Establish VOC emissions standards to reduce evaporative VOC emissions from the fermentation process at distilleries and wineries similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
	Can Coating	VOC limits on coatings	No control strategies identified	Not Recommended - No sources
	Commercial Cooking	None	Establish standards to control VOC emissions from commercial charbroilers similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
	Composting Operations	None	Establish work practice requirements to reduce VOC emissions from green waste composting similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
11.2	Confined Animal Facilities	Implement VOC emission mitigation measures from a menu of options	Reduce animal-count applicability thresholds; increase number of mitigation measures, and control efficiency	Not Recommended - Evaluated for Attainment Advancement
	Flares	None	Establish NO <sub>x</sub> emission standards for flares similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
2.44	Furnaces (Residential)	NO <sub>x</sub> limits from natural gas-fired, fan-type central furnaces	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
	Further Control of High-Emitting Spray Booth Facilities	None	Require additional controls to reduce VOC emissions from spray booths at facilities emitting > 20 tons per year	Not Recommended - No sources
2.21	Gasoline Storage, Loading, and Degassing of Tanks and Pipelines	VOC emission standards for organic liquid storage tanks; vapor-recovery requirements for loading at bulk plants and bulk terminals	Reduce VOC emission limits for gasoline loading at bulk plants and bulk terminals to be as stringent as BAAQMD; establish VOC emission standards for degassing storage tanks and pipelines similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Glass Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for glass melting furnaces	Not Recommended - No sources
2.29	Graphic Arts	VOC limits on inks, coatings, adhesives or use emission control system	Reduce VOC limits for flexographic ink on porous substrates, extreme performance ink, and metallic ink to be as stringent as SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement



Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
2.32	IC Engines	NO <sub>x</sub> limits on IC engines located at stationary sources	Reduce NO <sub>x</sub> limits to be stringent as SCAQMD	Not Recommended - Evaluated for Attainment Advancement
	Industrial Wastewater	None	Establish emission control standards for wastewater systems	Not Recommended - No sources
	LPG Transfer and Dispensing	None	Establish standards to control VOC emissions from LPG transfer and dispensing similar to the rules adopted by SCAQMD and VCAPCD	Not Recommended - Evaluated for Attainment Advancement
	Metal Melting Furnaces	None	Establish NO <sub>x</sub> emission limits for metal melting furnaces	Not Recommended - No sources
2.25	Metal Parts and Products Coating	VOC limits on coatings, strippers, cleaning solvents	Reduce VOC limits for general one-component, extreme high gloss, and prefabricated architectural coatings, similar to SCAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Metal Working Fluids	None	Establish VOC limits on metalworking fluids and direct-contact lubricants similar to the rules adopted by SCAQMD and VCAPCD	Not Recommended - Evaluated for Attainment Advancement
2.25-3	Miscellaneous Coating	None	Establish VOC limits and application method requirements for coating operations not covered by other rules, similar to SCAQMD, SJVUAPCD, VCAPCD, and BAAQMD requirements	Not Recommended - Evaluated for Attainment Advancement
	Miscellaneous Combustion Sources	None	Establish NO <sub>x</sub> emission limits on miscellaneous combustion equipment including dryers and ovens similar to rules adopted by SCAQMD and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
	Mold Release Agents	None	Establish VOC limits on mold release agents similar to the control measure proposed by SCAQMD	Not Recommended - Has not yet been implemented in SCAQMD or any other area

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
2.38	Municipal Landfill Gas	Landfill gas collection and control systems	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
2.23	Oil and Natural Gas Production	Leak detection and repair standards for components used in natural gas production and processing	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
6.0	Open Burning	Burning of certain materials prohibited; burn procedures to minimize smoke; burning is not allowed on days declared no-burn day	Reduce the types of allowable agricultural burns similar to SJVUAPCD requirements	Not Recommended - Evaluated for Attainment Advancement
2.29-2	Paper, Fabric, and Film Coatings	None	Establish VOC limits on coatings similar to rule adopted by SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
9.7	Petroleum Solvent Dry Cleaning	Use of closed-loop machine with primary control system; newer facilities must install close loop with both primary and secondary control systems	Expand applicability to include all non-halogenated solvents	Not Recommended - Evaluated for Attainment Advancement
2.25-2	Plastic Parts Coating	None	Establish VOC limits on plastic parts coatings similar to rule adopted by SCAMQD	Not Recommended - Evaluated for Attainment Advancement
2.30	Polyester Resin/Plastic Product Manufacturing	Limits on the monomer content of resin, use of vapor suppressants, use of close-mold systems, or emission capture and control system	Remove low-usage exemption, require non-atomizing equipment, and reduce monomer content similar to rules adopted by BAAQMD, SCAQMD, and SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement

Measure No.	Title	Current Requirements	Opportunity for Strengthening	Conclusion
2.41	Polystyrene /Polymeric Cellular (Foam) Manufacturing	VOC limits for the manufacturing of expanded polystyrene products	No control strategies identified	Not Recommended – No sources
	Portland Cement Manufacturing	None	Establish NO <sub>x</sub> limits for Portland cement manufacturing	Not Recommended – No sources
	Semiconductor Manufacturing	None	Establish VOC limits for semiconductor manufacturing	Not Recommended – No sources
	Synthetic Organic Chemical Manufacturing – Fugitive Leaks	None	Establish VOC emissions standards for leak detection and repair program	Not Recommended – No sources
	Soil Decontamination	None	Establish VOC emission control standards for soil vapor extraction systems, similar to rules adopted by BAAQMD and VCAPCD; Establish work practices to minimize VOC emissions from soil aeration similar to rule adopted by SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement
2.31	Solvent Cleaning	VOC limits on solvents, or use airtight/airless cleaning systems	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
2.34	Stationary Gas Turbines	NO <sub>x</sub> limits on stationary gas turbines	Reduce NO <sub>x</sub> emission limits to be as stringent as SCAQMD	Not Recommended - Evaluated for Attainment Advancement
	Wastewater Separators	None	Require solid cover, floating pontoon cover; double-deck cover, or vapor recovery system similar to rule adopted by SJVUAPCD	Not Recommended – No sources

<b>Measure No.</b>	<b>Title</b>	<b>Current Requirements</b>	<b>Opportunity for Strengthening</b>	<b>Conclusion</b>
2.37	Water Heaters and Small Boilers	Point-of-sale NO <sub>x</sub> emission standards on water heaters with rated heat input capacity less than 1 mmBtu/hr	No control strategies identified	Not Recommended - Evaluated for Attainment Advancement
2.39	Wood Products Coatings	VOC limits on coatings	Reduce VOC limits on wood coatings similar to rules adopted by SCAQMD/SJVUAPCD	Not Recommended - Evaluated for Attainment Advancement

## Appendix E.9 Sacramento Area Council of Governments (SACOG)

### Transportation Control Measures Considered

Information for this chapter was provided by SACOG based on a TCM RACM analysis (Sierra Research, 2015). A small number of control measures were identified during the TCM review, which have not been implemented in the Sacramento region. These were advanced for further RACM analysis and assessed based on the criteria specified in the 2015 Ozone Implementation Rule and USEPA’s RAC guidance. Factors considered included technical and economic feasibility, enforceability, local applicability, and ability to provide emission reductions before attainment deadline (advancement of attainment). These measures are discussed in more detail below.

Table E-6 RACM Analysis for Transportation Control Measures

Economic Feasibility		
TCM	Measure Description	Justification
Free transit during special events	Provide free alternative transportation to special events	Not cost-effective. SACOG cannot mandate that Transit Agencies provide free service.
Free rail-to-bus/bus-to-rail transfers	Vanpool and shuttle services at non-intermodal centers	Not cost-effective. SACOG cannot mandate that Transit Agencies provide free service.
Close roads for use of non-motorized traffic	Convert roadways to bike/pedestrian paths	Not cost-effective. The same emission reductions could be achieved with Complete Streets planning through road widening to create new bike and pedestrian paths and appropriate landscaping to provide a safe active transportation environment.
Free bikes	Provide free bikes to transit users	Not cost-effective. This voluntary measure does not guarantee emission reductions. Consumers could sell bikes for profit.
Truck Stop Electrification	Self-explanatory	Very costly to implement. May require state or federal subsidies. Cost-effectiveness >\$34,000/ton
Promote business closure on high ozone days	Self-explanatory	Would impact economic activity in the region and would not be socially and economically acceptable.
Cash incentives for carpoolers	Self-explanatory	Not cost-effective. SACOG’s T Funding Program will address this with employers through education and outreach.

<b>Advancement of Attainment</b>		
<b>TCM</b>	<b>Measure Description</b>	<b>Justification</b>
Bus queue jumps	Installing special lanes and signals to allow transit to get ahead in traffic	Due to infrastructure needs, cannot be implemented in time to advance attainment or by 2026.
Reduce idling at drive-throughs, parking lots and in traffic	Self-explanatory	No clear demonstration of air quality benefits; not easily enforceable
Reversible lanes	Change direction of travel during special events or during congestion periods	Will not advance attainment due to minimal emission reductions from this episodic strategy
Central Business District vehicle restrictions	Restrict vehicle use in downtown areas	Minimal air quality benefits that will not advance attainment

<b>Implementation Authority</b>		
<b>TCM</b>	<b>Measure Description</b>	<b>Justification</b>
Bus and carpool lanes on arterials	Provide fixed lanes for buses and carpools on arterial streets	No implementation authority; would require state agency authority and funds (Caltrans and California Transportation Commission (CTC))
Express toll lanes/ high-occupancy toll (HOT) lanes	Construct toll lanes to reduce congestion	No implementation authority; would require state agency authority and funds (Caltrans and CTC)
Mandatory bike racks for worksites	Mandate that employers install bike racks at businesses	No implementation authority; CA Health and Safety Code (HSC) §40717.6 prohibits mandatory employer-based trip reduction programs
Pay-As-You-Drive Insurance	Charge insurance fees based on driving patterns	No implementation authority; would require changes to state insurance practices and regulations
Express Busways/Dedicated Bus Lanes	Construct bus-only lanes	No implementation authority; would require state agency authority and funds (Caltrans and CTC)
Income tax credit to telecommuters	Self-explanatory	No implementation authority; would require changes to California tax law

<b>TCM</b>	<b>Measure Description</b>	<b>Justification</b>
Speed limit reduction	Reduce freeway speed limit to 55mph	No implementation authority; would require changes to California Vehicle Code
Off-peak goods movement	Require trucks to operate during off-peak hours	No implementation authority; would not be economically or socially acceptable
Truck only lanes	Construct or convert lanes for use by heavy-duty trucks only	No authority to implement; would require state agency authority and funds (Caltrans and CTC)
Divert Trucks from Nonattainment Areas	Require pass-through trucks to choose routes away from the Sacramento region	No authority to implement; would require state agency authority and funds (CARB, Caltrans, and CTC)
Satellite Work Centers	Work centers set-up closer to where employers live	No authority to implement; CA HSC §40717.6 prohibits mandatory employer-based trip reduction programs

### Conclusions

Out of the approximately 20 candidate TCMs identified as candidate RACM, no measures were found to meet the criteria for RACM implementation. Based on a comprehensive review of TCM projects in other nonattainment areas, it was determined that the TCMs being implemented in the Sacramento region represent all RACM.

## Appendix E.10 California Air Resource Board (CARB)

### Mobile sources Reasonably Available Control Measure (RACM) Evaluation

#### Introduction

CARB is responsible for measures to reduce emissions from mobile sources needed to attain the NAAQS. This section of the Appendix will discuss how California's mobile source measures meet RACM.

CARB's comprehensive strategy to reduce emissions from mobile sources includes stringent emissions standards for new vehicles, in-use programs to reduce emissions from existing vehicle and equipment fleets, cleaner fuels that minimize emissions, and incentive programs to accelerate the penetration of the cleanest vehicles beyond that achieved by regulations alone. Taken together, California's mobile program meets RACM requirements in the context of ozone nonattainment.

CARB developed its SIP strategy through a multi-step measure development process, including extensive public consultation, to develop and evaluate potential strategies for mobile source categories under CARB's regulatory authority that could contribute to expeditious attainment of the standard. First, CARB developed a series of technology assessments for heavy-duty mobile source applications and the fuels necessary to power them<sup>1</sup> along with ongoing review of advanced vehicle technologies for the light-duty sector in collaboration with USEPA and the National Highway Traffic Safety Administration. CARB staff then used a scenario planning tool to examine the magnitude of technology penetration necessary, as well as how quickly technologies need to be introduced to meet attainment of the standard.

CARB staff released a discussion of draft Mobile Source Strategy<sup>2</sup> for public comment in October 2015. This strategy specifically outlined a coordinated suite of proposed actions to not only meet federal air quality standards, but also achieve greenhouse gas emission reduction targets, reduce petroleum consumption, and decrease health risk from transportation emissions over the next 15 years. CARB staff held a public workshop on October 16, 2015 in Sacramento, and on October 22, 2015, CARB held a public Board meeting to update the Board and solicit public comment on the Mobile Source Strategy in Diamond Bar.

CARB Staff continued to work with stakeholders to refine the measure concepts for incorporation into related planning efforts including the 75 ppb 8-hour ozone SIPs. In May 2016, CARB released an updated Mobile Source Strategy. On May 17, 2016, CARB released the proposed SIP strategy for a 45-day public comment period.

---

<sup>1</sup> Technology and Fuel assessments <http://www.arb.ca.gov/msprog/tech/tech.htm>

<sup>2</sup> 2016 Mobile Source Strategy <http://www.arb.ca.gov/planning/sip/2016sip/2016mobsr.htm>



The current mobile source program and proposed measures included in the SIP Strategy provide attainment of the ozone standard as expeditiously as practicable and meet RFP requirements.

### **Waiver Approvals**

While the CAA preempts states from adopting emission standards and other emission-related requirements for new motor vehicles and engines, it allows states to seek a waiver from the federal preemption to enact emission standards and other emission-related requirements for new motor vehicles and engines that are at least as protective as applicable federal standards, except for locomotives and engines used in farm and construction equipment which are less than 175 horsepower (hp). The CAA also allows California to seek authorization for more stringent standards for new and in-use off-road vehicles and engines, and allows other states to adopt the standards after EPA authorization.

Over the years, California has received waivers and authorizations for over 100 regulations. The most recent California standards and regulations that have received waivers and authorizations are Advanced Clean Cars (including ZEV and LEV III) for light-duty vehicles, and On-Board Diagnostics, Heavy-Duty Idling, Malfunction and Diagnostics System, In-Use Off-Road Diesel Fleets, Large Spark Ignition Fleet, Mobile Cargo Handling Equipment for heavy-duty engines. Other authorizations include Off-Highway Recreational Vehicles and the Portable Equipment Registration Program.

Finally, CARB obtained an authorization from USEPA to enforce adopted emission standards for off-road engines used in yard trucks and two-engine sweepers. CARB adopted the off-road emission standards as part of its “Regulation to Reduce Emissions of Diesel Particulate Matter, Oxides of Nitrogen and Other Criteria Pollutants from In-Use Heavy-Duty Diesel-Fueled Vehicles,” (Truck and Bus Regulation). The bulk of the regulation applies to in-use heavy-duty diesel on-road motor vehicles with a gross vehicle weight rating in excess of 14,000 pounds, which are not subject to preemption under section CAA section 209(a) and do not require a waiver under CAA section 209(b).

### **Light- and Medium-Duty Vehicles**

Light- and medium-duty vehicles are currently regulated under California’s Advanced Clean Cars program including the Low-Emission Vehicle III (LEV III) and Zero-Emission Vehicle (ZEV) programs. Other California programs such as the 2012 Governor Brown Executive Order (B-16-2012) to put 1.5 million zero-emission vehicles on the road by 2025, and California’s Reformulated Gasoline program (CaRFG) will produce substantial and cost-effective emission reductions from gasoline-powered vehicles.

CARB is also active in implementing programs for owners of older dirtier vehicles to retire them early. The “car scrap” programs, like the Enhanced Fleet Modernization Program, and Clean Vehicle Rebate Project provide monetary incentives to replace old vehicles with zero-emission vehicles. The Air Quality Improvement Program (AQIP) is a voluntary incentive program to fund clean vehicles.

Taken together, California’s emission standards, fuel specifications, and incentive programs for on-road light- and medium-duty vehicles represent all measures that are technologically and economically feasible in the context of a RACM assessment.

### Heavy-Duty Vehicles

California’s heavy-duty vehicle emissions control program includes requirements for increasingly tighter new engine standards and address vehicle idling, certification procedures, on-board diagnostics, emissions control device verification, and in-use vehicles. This program is designed to achieve an on-road heavy-duty diesel fleet with 2010 engines emitting 98 percent less NO<sub>x</sub> and PM<sub>2.5</sub> than trucks sold in 1986.

Most recently in the ongoing efforts to go beyond federal standards and achieve further reductions, CARB adopted the Optional Reduced Emissions Standards for Heavy-Duty Engines regulation in 2014 that establishes the new generation of optional NO<sub>x</sub> emission standards for heavy-duty engines.

The recent in-use control measures include On-Road Heavy-Duty Diesel Vehicle (In-Use) Regulation, Drayage (Port or Rail Yard) Regulation, Public Agency and Utilities Regulation, Solid Waste Collection Vehicle Regulation, Heavy-Duty (Tractor-Trailer) Greenhouse Gas Regulation, ATCM to Limit Diesel-Fueled Commercial Motor Vehicle Idling, Heavy-Duty Diesel Vehicle Inspection Program, Periodic Smoke Inspection Program, Fleet Rule for Transit Agencies, Lower-Emission School Bus Program, and Heavy-Duty Truck Idling Requirements. In addition, CARB’s significant investment in incentive programs provides an additional mechanism to achieve maximum emission reductions from this source sector.

Taken together, California’s emission standards, fuel specifications, and incentive programs for heavy-duty vehicles represent all measures that are technologically and economically feasible in the context of a RACM assessment.

### Off-Road Vehicles and Engines

California regulations for off-road equipment include not only increasingly stringent standards for new off-road diesel engines, but also in-use requirements and idling restrictions. The Off-Road Regulation is an extensive program designed to accelerate the penetration of the cleanest equipment into California’s fleets, and impose idling limits on off-road diesel vehicles. The program goes beyond emission standards for new engines through comprehensive in-use requirements for legacy fleets.

Taken together, California's comprehensive suite of emission standards, fuel specifications, and incentive programs for off-road vehicles and engines represent all measures that are technologically and economically feasible in the context of a RACM assessment.

### Other Sources and Fuels

The emission limits established for other mobile source categories, coupled with USEPA waivers and authorization of preemption establish that California's programs for motorcycles, recreational boats, off-road recreational vehicles, cargo handling equipment, and commercial harbor craft sources meet the requirements for RACM.

Cleaner burning fuels also play an important role in reducing emissions from motor vehicles and engines as CARB has adopted a number of more stringent standards for fuels sold in California, including the Reformulated Gasoline program, low sulfur diesel requirements, and the Low Carbon Fuel Standard. These fuel standards, in combination with engine technology requirements, ensure that California's transportation system achieves the most effective emission reductions possible.

Taken together, California's emission standards, fuel specifications, and incentive programs for other mobile sources and fuels represent all measures that are technologically and economically feasible in the context of a RACM assessment.

### Mobile Source Summary

California's long history of comprehensive and innovative emissions control has resulted in the most stringent mobile source control program in the nation. USEPA has previously acknowledged the strength of the program in their approval of CARB's regulations and through the waiver process. Since then, CARB has continued to substantially enhance and accelerate reductions from our mobile source control programs through the implementation of more stringent engine emissions standards, in-use requirements, incentive funding, and other policies and initiatives as described in the preceding sections.

The CARB process for developing the proposed State measures included an extensive public process and is consistent with USEPA RACM guidance. Through this process CARB found that there are no additional RACM that would advance attainment of the 75 ppb 8-hour ozone standard in the SFNA from emissions reductions associated with unused regulatory control measures. As a result, California's mobile source control programs fully meet the requirements for RACM.

### Appendix E.11 References

USEPA (57 FR 13498). *General Preamble for the Implementation of Title I of the Clean Air Act Amendments of 1990*. Federal Register, Volume 57, 16 April, 1992, p.13498. Print.

USEPA. (80 FR 12282 - 12283) *Final Rule to Implement the 2008 8-Hour Ozone National Ambient Air Quality Standard for Ozone: SIP Requirements*. Federal Register, Volume 80, 6 March 2015, p. 12282-12283. Print.

USEPA. *Guidance on the Reasonably Available Control Measures (RACM) Requirement and Attainment Demonstration Submissions for Ozone Nonattainment Areas*. United States Environmental Protection Agency, December [1999.] Print.

Sierra Research. *Reasonably Available Control Measures Analysis for the Sacramento Area Council of Governments*. Sacramento, CA: Sierra Research, 12 November [2015.] Print.

SMAQMD, et al. *Sacramento Regional 8-Hour Ozone Attainment and Reasonable Further Progress Plan (2013 SIP Revision)*. Sacramento, CA: Sacramento Metropolitan Air Quality Management District, 26 September [2013.]

## APPENDIX F

### Federal Clean Air Act Requirements

**Appendix F Federal Clean Air Act Requirements**

**Table F-1 General Nonattainment Plan Requirements**

<b>Required Plan Element</b>	<b>Description</b>	<b>Location in Plan</b>
Reasonably Available Control Measures (RACM) [Section 172(c)(1)]	The plan should provide for the implementation of all reasonably available control measures as expeditiously as practicable, including reduction in emissions from existing sources through the adoption of reasonably available control technology.	Chapter 7 (Control Measures) Appendix E (RACM Analysis)
Reasonable Further Progress (RFP) [Section 172(c)(2)]	The plan should meet reasonable further progress requirements for emission reduction.	Chapter 12 (RFP Demonstration) Appendix D (RFP Progress Demonstrations)
Emissions Inventory [Section 172(c)(3)]	The plan should include a comprehensive, accurate, current inventory of actual emissions from all sources of the relevant pollutant or pollutants in such area, including periodic revisions as the Administrator may determine necessary to assure that the requirements of this part are met.	Chapter 5 (Emissions Inventory) Appendix A (Emissions Inventory)
Identification and Quantification [Section 172(c)(4)]	The plan should identify and quantify the emissions, if any, of any such pollutant or pollutants, which will be allowed, in accordance with section 173(a)(1)(B), from the construction and operation of major new or modified stationary sources in each such area. The plan shall demonstrate to the satisfaction of the United States Environmental Protection Agency (USEPA) that the emissions quantified for this purpose will be consistent with the achievement of reasonable further progress and will not interfere with attainment of the applicable national ambient air quality standard by the applicable attainment date.	Chapter 5, Sections 5.2 (Emissions Inventory Forecasts) Chapter 8 (Attainment Demonstration) Chapter 12 (RFP Demonstration)
Permits for new and modified stationary sources [Section 172(c)(5)]	The plan provisions should require permits for the construction and operation of new or modified major stationary sources anywhere in the nonattainment area, in accordance with section 173.	Chapter 3, Section 3.5 (NSR Review Requirements)
Other Measures [Section 172(c)(6)]	The plan provisions shall should include enforceable emission limitations, and such other control measures, means or techniques (including economic incentives such as fees, marketable permits, and auctions of emission rights), as well as schedules and timetables for compliance, as may be necessary or appropriate to provide for attainment by the applicable date.	Chapter 7 (Control Measures)

Required Plan Element	Description	Location in Plan
Compliance with Section 110(a)(2) [Section 172(c)(7)]	The plan provisions should meet the applicable provisions of section 110(a)(2). Section 110(a)(2) includes reasonable notice and public hearing requirements for plan adoptions.	Chapters 2, Section 2.4.4 (Public Input and Review Process)
Equivalent Techniques [Section 172(c)(8)]	Upon application by any State, the USEPA may allow the use of equivalent modeling, emission inventory, and planning procedures, unless USEPA determines that the proposed techniques are, in the aggregate, less effective than the methods specified by the USEPA.	Not Applicable – Standard methods employed in chapters.
Contingency Measures [Section 172(c)(9)]	The plan should include the implementation of specific measures to be undertaken if the area fails to make reasonable further progress, or to attain the national primary ambient air quality standard by the applicable attainment date. Such measures shall be included in the plan revision as contingency measures to take effect in any such case without further action by the State or the USEPA.	Chapter 8, Section 8.2 (Attainment Demonstrations Evaluation) Chapter 12, Section 12.3 (Contingency Measure Requirement)
Demonstration of attainment of the standard as expeditiously as practicable but not later than 20 years after designation [Section 181(a)]	Each area designated nonattainment for ozone pursuant to section 107(d) shall be classified at the time of such designation, as a Marginal Area, a Moderate Area, a Serious Area, a Severe Area, or an Extreme Area based on the design value for the area. For each area classified under this subsection, the primary standard attainment date for ozone shall be as expeditiously as practicable.	Chapter 3 (Federal Clean Air Act Requirements) Chapter 8 (Attainment Demonstration)

Table F-2 Severe Area Plan Requirements for Ozone Nonattainment Areas

Required Plan Element	Description	Location in Plan
Inventory [Section 182(a)(1)]	Submit a comprehensive, accurate, current inventory of actual emissions from all sources.	Chapter 5 (Emissions Inventory) Appendix A (Emissions Inventory)
Emissions Statement [Section 182(a)(3)(B)]	Within 2 years after the date of the enactment of the CAA Amendments of 1990, the State shall submit a revision to the SIP to require that the owner or operator of each stationary source of oxides of nitrogen or volatile organic compounds provide the State with a statement, in such form as the Administrator may prescribe (or accept an equivalent alternative developed by the State) for classes or categories of sources, showing the actual emissions of oxides of nitrogen and volatile organic compounds from that source.	Chapter 5 (Emissions Inventory)
Reasonably Available Control Technology [Section 182(b)(2)]	Implementation of control technologies for VOC sources covered by control technique guidelines (CTG) documents and all other major stationary sources of VOCs that are located in the area.	Chapter 3, Section 3.9 (RACT Requirements)
Motor Vehicle Inspection and Maintenance [Section 182(b)(4)]	Provide for a vehicle inspection and maintenance program as described in Section 182(a)(2)(B).	Chapter 7, Section 7.3.1.5 (California Enhanced Smog Check Program)
Enhanced Monitoring [Section 182 (c)(1)]	The State shall commence such actions as may be necessary to adopt and implement a program based on enhanced monitoring (Photochemical Assessment Monitoring Stations, PAMS), to improve monitoring for ambient concentrations of ozone, oxides of nitrogen and volatile organic compounds and to improve monitoring of emissions of oxides of nitrogen and volatile organic compounds.	Chapter 4, Section 4.1.1 (Ozone Monitoring Sites)
Attainment demonstration [Section 182(c)(2)(A)]	A demonstration that the plan will provide for attainment of the national ambient air quality standard as expeditiously as practicable by the applicable attainment date. The attainment demonstration must be based on photochemical grid modeling.	Chapter 4 (Air Quality Trends) Chapter 6 (Air Quality Modeling Analysis) Chapter 8 (Attainment Demonstration) Appendix B (Photochemical Modeling)
Reasonable Further Progress demonstration [Section 182(c)(2)(B) and (C)]	A demonstration that the plan will result in VOC emissions (and/or NO <sub>x</sub> emissions) reductions from the baseline emissions of an average of at least three percent each year.	Chapter 12 (RFP demonstration)



<b>Required Plan Element</b>	<b>Description</b>	<b>Location in Plan</b>
Enhanced vehicle inspection and maintenance program [Section 182(c)(3)]	The State shall provide for an enhanced program to reduce hydrocarbon emissions and NO <sub>x</sub> emissions from in-use motor vehicles registered in each urbanized area.	Chapter 7, Section 7.3.1.5 (California Enhanced Smog Check Program) and Chapter 5, Section 5.3.2 (on-road motor vehicle emissions EMFAC2014)
Clean-Fuel Vehicle Programs [Section 182(c)(4)]	The State will develop a Program including all measures necessary to make the use of clean alternative fuels in clean-fuel vehicles (as defined in part C of title II) economic from the standpoint of vehicle owners.	Chapter 7, Section 7.3 (State and Federal Control Measures)
Vehicle Miles Traveled Offset [Section 182(d)(1)(A)]	The Plan shall identify and adopt specific enforceable transportation control strategies and transportation control measures to offset any growth in emissions from growth in vehicle miles traveled or numbers of vehicle trips.	Appendix C (VMT Offset Analysis)
General Offset requirements [Section 182(d)(2)]	The ratio of total emission reductions of volatile organic compounds (VOCs) to total increased emissions of such air pollutant shall be at least 1.3 to 1.	Chapter 3, Section 3.5 (NSR Review Requirements)
Milestones [Section 182(g)]	Provide a report every three years after the designation to determine whether the nonattainment area has achieved a reduction in emissions during the preceding interval equivalent to the total emission reductions required to be achieved by the attainment date given in the plan.	Chapter 3, Section 3.10 (Milestone Reports)